



UNIV. OF
TORONTO
LIBRARY







Digitized by the Internet Archive
in 2009

<http://www.archive.org/details/fieldiana14fiel>

Chicago Natural History
Museum
(PUBLICATIONS

OF

FIELD MUSEUM OF NATURAL
HISTORY)

Feldman. Zoology 4
(ZOOLOGICAL SERIES)

VOLUME XIV



321344
20.11.35

CHICAGO, U. S. A.

1921 - 1925


FIELD MUSEUM OF NATURAL
HISTORY

QL

1

F4

v.14



FIELD MUSEUM OF NATURAL HISTORY

PUBLICATION 246

ZOOLOGICAL SERIES

VOL. XIV, No. 4

CONTENTS
TO
VOLUME XIV
NUMBERS 1 to 3

WILFRED H. OSGOOD
Curator, Department of Zoology
EDITOR



CHICAGO, U. S. A.

1928

CONTENTS

A Monographic Study of the American Marsupial, <i>Caenolestes</i> . By Wilfred H. Osgood.....	1-156
The Brain of <i>Caenolestes obscurus</i> . By C. Judson Herrick.....	157-162
Review of Living <i>Caenolestids</i> with Description of a New Genus from Chile. By Wilfred H. Osgood.....	163-172
The Brains of the South American Marsupials <i>Caenolestes</i> and <i>Orolestes</i> . By Jeannette Brown Obenchain.....	173-232

LIST OF ILLUSTRATIONS

PLATES

- I. *Caenolestes obscurus*.
- II. External characters.
- III. General musculature after removal of superficial fascia.
- IV. Muscles and glands of the head and neck.
- V. Muscles of the legs.
- VI. Muscles of throat and inner side of thigh.
- VII. Parts of alimentary system.
- VIII. Reproductive organs.
- IX. Dissections of male organs.
- X. Parts of reproductive, alimentary and respiratory systems.
- XI. Cervical vertebrae.
- XII. Bones of sternum, hind leg and thoracic region.
- XIII. Scapula, pelvis and lumbar vertebrae.
- XIV. Sacral and caudal vertebrae.
- XV. Humerus, ulna and radius.
- XVI. Manus and pes.
- XVII. Longitudinal section of skull.
- XVIII. Dentition and hard palate.
- XIX. Fig. 1. Mounted skeleton of *Nesogale dobsoni*.
Fig. 2. Mounted skeleton of *Caenolestes obscurus*.
- XX. Skull of *Caenolestes*.
- XXI. Dorsal and lateral surfaces of brain.
- XXII. Ventral view of brain.
- XXIII. Figs. 1, 1a, 1b. Skull of *Caenolestes obscurus*.
Figs. 2, 2a, 2b. Skull of *Orolestes inca*.
Figs. 3, 3a, 3b, 3c. Skull of *Rhyncholestes raphanurus*.
- XXIV. *Orolestes inca*. Dorsal view of brain.
- XXV. *Orolestes inca*. Ventral view of brain.
- XXVI. *Orolestes inca*. Fig. 3. Lateral view of brain. Fig. 4. Median section of brain.
- XXVII. Median surface of hemisphere. Fig. 5. *Orolestes inca*. Fig. 6. *Caenolestes obscurus*.
- XXVIII. Figs. 7-10. Cerebella of *Caenolestes*, *Notoryctes* and *Perameles*. Fig. 11. Development of mammalian "hippocampal figure."
- XXIX. Figs. 12-15. Reconstructions of hemispheres. Fig. 16. Ventral diverticulum of recessus superior.
- XXX. Fig. 17. Dissected diagram of hippocampus of *Caenolestes*. Fig. 18. Temporal hippocampus of *Didelphis*.
- XXXI. Temporal recurving of hippocampus. Fig. 19. *Hypsiprymnus*. Fig. 20. Rabbit. Fig. 21. *Felis leo*. Fig. 22. *Notoryctes*.
- XXXII. Transverse sections of brain of *Caenolestes*.
- XXXIII. Transverse sections of brain of *Caenolestes*.
- XXXIV. Transverse sections of brain of *Caenolestes*.
- XXXV. Transverse sections of brain of *Caenolestes*.
- XXXVI. Transverse sections of brain of *Caenolestes*.

MAP

Records of *Caenolestes* and *Orolestes*.....

Page 16



CAENOLESTES OBSCURUS.
Nearly natural size.

1
FIELD MUSEUM OF NATURAL HISTORY.

PUBLICATION 207.

ZOÖLOGICAL SERIES.

VOL. XIV, No. I.

A MONOGRAPHIC STUDY OF THE
AMERICAN MARSUPIAL, CÆNOLESTES.

BY

WILFRED H. OSGOOD,
Assistant Curator of Mammalogy and Ornithology.

WITH

A DESCRIPTION OF THE BRAIN OF
CÆNOLESTES.

BY

C. JUDSON HERRICK,
Professor of Neurology, University of Chicago.

CHARLES B. CORY,
Curator, Department of Zoölogy.



CHICAGO, U. S. A.
MAY, 1921.

A MONOGRAPHIC STUDY OF THE AMERICAN MARSUPIAL, CÆNOLESTES.

BY WILFRED H. OSGOOD.

CONTENTS.

	PAGE
Introduction	4
History	6
Distribution	15
Habits	17
External characters	19
Measurements	21
Myology	22
Muscles of the skin	22
Muscles of the back	24
Muscles of the head	30
Muscles of the neck	32
Muscles of the thorax	34
Muscles of the abdomen	35
Muscles of the sacrum	36
Muscles of the tail	38
Muscles of the anterior limb	40
Muscles of the forefoot	46
Muscles of the hind limb	47
Muscles of the hind foot	56
Summary of myological characters	59
Urinogenital System	61
Male organs	62
Female organs	66
Alimentary Canal	69
Glands	73
Respiratory System	76
Circulatory System	77
Skeleton	77
Cervical vertebrae	77
Thoracic vertebrae	82
Lumbar vertebrae	84
Sacral and caudal vertebrae	84
Ribs and sternum	86
Pectoral girdle	87
Pelvic girdle	88

Skeleton—Continued.

Arm and forearm	89
Bones of the hand	92
Thigh and leg	93
Bones of the foot	95
Summary of skeletal characters	97
Skull	99
Summary of cranial characters	109
Dentition	111
Number and homologies of teeth	112
Upper incisors	116
Lower incisors	118
Canines	118
Upper premolars	119
Upper molars	120
Lower premolars	126
Lower molars	127
Origin of Diprotodonty	128
Relationships of <i>Wynyardia</i>	136
Relationships of <i>Myrmecoboides</i>	139
Phylogeny and taxonomy	140
Dispersal of marsupials	145
General summary	150
Literature cited	152
The Brain of <i>Cenolestes obscurus</i>	157

INTRODUCTION.

Owing to their many peculiarities of structure and their probable relationship to ancient and long extinct types, marsupials are among the most interesting of mammals. Their present distribution, largely in Australasia but also in South America, adds to the importance they possess for the morphologist and phylogenist. They have long been classified in two large groups or suborders, the Polyprotodontia and the Diprotodontia. These were first recognized by De Blainville in 1816 and later by Owen in 1866. Broadly speaking, the polyprotodonts include the carnivorous or insectivorous forms with small and relatively numerous incisors, while the diprotodonts are the herbivorous forms with the incisors reduced in number and modified much as in rodents. Without exception, the diprotodonts are characterized also by a syndactylous foot, while the polyprotodonts, with the exception of one family (Peramelidae), are eleutherodactylous. This makes slightly different divisions (Diadactyla and Syndactyla) possible, but other considerations have favored the conclusion that the two major groups indicated by the dentition are the most natural ones.

Until recently, it was supposed that all modern diprotodonts and the

majority of polyprotodonts were confined to Australasia, only one family of polyprotodonts being known elsewhere, the Didelphiidae or opossum family of South and Central America. Under these conditions, it was a matter of the greatest interest when a small marsupial, widely different from the opossums, was discovered in South America. This animal, now known by the generic name *Cænolestes*, presented an apparent combination of diprotodont and polyprotodont characters, having a diprotodont dentition and an eleutherodactylous foot. Moreover, it was found to be closely allied to certain extinct forms known from fragmentary remains from the Tertiary of Patagonia, being clearly a surviving member of a highly differentiated group detailed knowledge of which promised to throw new light on problems connected with the origin and dispersal of marsupials. These facts were evident from examination of the few specimens, consisting merely of skulls and skins, which for many years were the only ones available. Such specimens had been obtained through aboriginal sources and the exact habitat of the animal was unknown.

In 1911, while making general collections in the mountains of western Venezuela (Osgood, 1912), I had the good fortune to discover *Cænolestes* living in dense forests at an altitude of about 8000 feet. As a result of assiduous trapping, a total of eleven animals was captured and, despite poor equipment for the care of anatomical material, two entire specimens representing both sexes and two others with some parts mutilated were preserved in formaldehyde and bichloride of mercury. In addition, several dry skeletons and a small number of conventional skins and skulls were saved. This material has formed the basis of the following study. Its importance seemed to demand the fullest possible treatment and, although this has not been accomplished, it is hoped that sufficiently detailed information has been obtained to warrant general conclusions. Study of particular organs and systems by specialists would have been exceedingly desirable, but the nature of the material and the conditions under which it has been studied have not made this feasible except to a limited extent.¹ The coöperation of Dr. C. Judson Herrick has been greatly appreciated and his description of the brain of *Cænolestes* together with the figures drawn under his direction which are incorporated in the present publication will doubtless add greatly to its value.

Comparative material, especially of Australian forms, has been confined almost exclusively to skeletons and skulls. Fortunately a fairly representative collection of these is possessed by Field Museum and this has been supplemented by specimens borrowed from the U. S.

¹ It is especially to be regretted that no study of the organ of Jacobson has been made.

National Museum, the American Museum of Natural History, and the Museum of Comparative Zoölogy. To the officials of these institutions, especially Dr. J. A. Allen and Dr. G. M. Allen, acknowledgment is gratefully made.

The accompanying illustrations are from drawings made by several different artists. The line drawings of muscles and soft parts (Pls. IV-X) are the work of Mr. Robert E. Snodgrass, now of the U. S. Bureau of Entomology; the wash drawings of skeletal parts (Pls. XI-XVIII) were done by Mr. Kenji Toda of the Department of Zoölogy, University of Chicago; the illustrations of the brain (Pls. XXI-XXII) are by Mr. A. B. Streedain formerly of the Department of Anatomy, University of Chicago; and the drawings of external parts (Pls. I-II) are by Mr. L. L. Pray of Field Museum.

HISTORY.

In a brief list of mammals from Ecuador published by R. F. Tomes in 1860, the following note appeared without reference either to genus or species:

"A small animal about the size of the Water Shrew (*Sorex fodiens*), with external characters and incisor teeth so much like those of the *Soricidae* as to have led in the first instance to the belief that it was a placental *Insectivore*, perhaps in some degree resembling the *Solenodon* of Cuba. However, the existence of a small and rudimentary pouch sufficiently attests the implacental nature of the creature, which but for this must certainly, as far as external appearances go, be regarded as one of the *Soricidae*. A more ample account of it will be given on a future occasion."

This is the first published reference to the animal now known as *Cænolestes*. Three years later (Tomes 1863), the single immature specimen was described as to its dentition and external characters and given the name *Hyracodon fuliginosus* without any reference to its affinities within the order of marsupials. Subsequently it received little notice for many years. In 1880, Alston referred to it in the following brief statement, published as a footnote (Alston 1880, p. 195):

"A small Marsupial from Ecuador, named *Hyracodon fuliginosus* by Mr. Tomes (P. Z. S. 1863, p. 51, pl. VIII), may represent a distinct family, but it is still only known from the very unsatisfactory original description. In any case it will require a new title, the name *Hyracodon* having been applied to a genus of fossil ungulates by Professor Leidy in 1856, seven years before its use by Mr. Tomes."

Again in the concluding paragraph of the very important Catalogue of Marsupialia and Monotremata in the British Museum (Thomas 1888) it receives only this slight mention:

"And, finally, one animal, referred to this Order, has been so insufficiently described that no idea of its proper position can be gained, nor have any further specimens of it been collected. This is—

"*Hyracodon fuliginosus*, Tomes, P. Z. S. 1863, p. 50, pl. viii (animal).

"Hab. Ecuador."

So meager were the data regarding it, that Lydekker (1894) completely ignored it in his very comprehensive Hand-book of Marsupials and Monotremes. Thus for more than thirty years after its discovery the subject of this monograph remained almost as unknown as if no specimen existed. In 1895, however, a second specimen, consisting of a skin and skull, was received at the British Museum and in two brief but important papers (Thomas 1895) was promptly described, tentatively classified, and figured. Its obvious affinity with extinct Patagonian forms was at once recognized and its importance in connection with theories of the dispersal of marsupials was noted. The preoccupied name *Hyracodon* was discarded and a new name, *Cænolestes*, was proposed as well as another specific name, *obscurus*, the second specimen having been obtained in the Bogota region and appearing to represent a different species of larger size. The following extracts from these papers of Thomas are of interest:

"In vindication of Mr. Tomes's paper I should like to say, firstly, that his description, hitherto supposed (from our ignorance of any such animal) to be imperfect or incorrect, proves to agree, so far as it goes, very closely with the present specimen; and, secondly, that remarks on the affinities of the animal must have been at that time more easily wanted than given, since even now, with infinitely greater material and the best of advice, I am unable to be at all positive about the exact position and relationships of the little marsupial described by Mr. Tomes."

"The rediscovery of this long-lost genus, whose wide distinction from all other living marsupials its original describer does not appear to have fully appreciated, is one of the most interesting events in mammalogy that has happened for many years. A full description of the animal will be given elsewhere; but it may be here briefly stated (1) that *Cænolestes* represents among the marsupials a family, and, perhaps, a suborder, entirely different from any now living; and (2) that it is closely related to, and evidently a surviving representative of, some of the fossil marsupials from the Santa Cruz beds of Patagonia."

"Apart from this, the survival to the present day of a member of so

ancient a group, otherwise wholly extinct, is a fact of the utmost interest, and one whose discovery will be welcomed by every zoölogist."

"I have proposed for it the name *Cænolestes*, as it is a modern member of an ancient group."

"The affix '*lestes*' is connected in mammalogy with small and ancient fossil marsupials, e. g., *Microlestes*, *Amphilestes*, etc., so that the above name may be considered to represent an existing animal with ancient fossil relatives."

"*Cænolestes*, with its uninteresting exterior, appeals mainly to the technical mammalogist. To him, however, with its intense palæontological and geographical interest, and the added puzzle its structure gives rise to in the general classification of the order, no animal will appear more important or more worthy of close and detailed study. That by the arrival of spirit specimens account of its anatomy may be rendered possible is very much to be hoped."

In these papers of Thomas, *Cænolestes* was classified as a diprotodont and a member of the family Epanorthidae with the reservation that "It is, however possible that, in spite of the resemblance of the teeth of *Cænolestes* to those of certain Australian Diprotodonts, the study of further material, including soft parts, skeleton, and milk-teeth, will bring out differences of such importance as to necessitate its subordinal separation from them." He says further, "It is clearly a diprotodont, as not only does it possess the characteristic development of the lower incisors, but even the molars resemble most closely in structure those of certain members of the family *Phalangeridae*, while being wholly unlike those of the typical Polyprotodonts. From all the existing Diprotodonts, however, apart from its habitat and numerous detailed differences, *Cænolestes* is at once distinguished by not being syndactylous, a character which is always considered of family rank. It forms, therefore, among existing marsupials a peculiar family, and one which in America represents the Diprotodonts of Australia, just as Didelphyidae do the Polyprotodonts."

Although some further material was obtained in later years, it was similar to that used by Thomas and consisted only of a few native-collected skins and skulls. Additional study of these, therefore, added but little to what was already known and opinions varied as to the interpretation of the peculiar characters of the animal.

The next reference to *Cænolestes* was a note by Ameghino (1897, p. 95) from which the following may be quoted:

"M. Thomas est venu à La Plata, rapportant avec lui un crâne de *Cænolestes* que nous avons soigneusement comparé aux formes fossiles de Patagonie et nous avons pu reconnaître qu'il presente plus de rap-

ports avec les Garzonidæ qu'avec les Epanorthidæ. Pourtant il est probable que le *Coenolestes* devra constituer le type d'une famille nouvelle."

Accepting the suggestion of Ameghino and others, Trouessart (1898) gave it the rank of a family, the Cænolestidae, including only the one genus, and assigned it to the suborder Diprotodontia. Ameghino (1900) later treated it as a family of diprotodonts, and still later (1903), in his well known rearrangement of marsupials and rodents, he included in the family the genus *Cænolestes* and also the extinct genus *Zygolestes*. The family was classified with four others, containing only extinct forms, in his suborder Paucituberculata, which in turn was referred to the order Plagiaulacoidea and regarded as directly ancestral to the Australian diprotodonts. The name Diprotodontia was reserved to designate a superorder conceived to embrace not only American and Australian diprotodonts but also the multituberculates and the rodents. Although Ameghino's ideas have not been generally accepted, especially those regarding the derivation of rodents, his strong conviction of the close alliance of American and Australian diprotodonts is noteworthy in the present connection. While differing from him in other respects, Weber (1904) agreed with Ameghino in placing *Cænolestes* in neither the Diprotodontia nor the Polyprotodontia but in a third suborder of marsupials for which Ameghino's name Paucituberculata was available. Meanwhile, Bensley (1903), in his very important paper on the evolution of Australian marsupials, included *Cænolestes* in his "First Neogæic Radiation" and in casual references indicated his belief that its dentition is due to parallel or convergent development rather than to any direct relationship to Australian diprotodonts closer than that of common derivation from a didelphid ancestry.

On the other hand, Sinclair (1905, 1906) and Scott (1913) revert to the opinion of Thomas and Ameghino that *Cænolestes* and allied extinct forms are so closely related to Australian diprotodonts as to furnish strong evidence of a former land connection between South America and Australia. Sinclair includes all South American forms with diprotodont dentition in the family Cænolestidae which he divides into two subfamilies, of which the Cænolestinae embraces the genera *Cænolestes*, *Garzonia*, and *Halmariphus*. Of this family he says (1905): "The Cænolestidae resemble the primitive phalangers in so many respects that it is impossible to escape the conclusion that the two families are related and not merely convergent groups." This statement is somewhat modified in a later paper (1906), as follows: "While substantially the same conclusions are still held, it is proper to point out the evidence in favor of the view that the striking similarity in dental structure

displayed by the two families may be explained by convergence. Until the upper dentition, skull and feet of the *Cænolestidæ*, and especially of the primitive members of the family, are fully known, this must remain an unsettled question. At present the arguments in favor of the alternatives expressed are about equally balanced."

The next author to give special attention to *Cænolestes* was Miss Pauline Dederer (1909), who made a study of the skull and pointed out certain resemblances to polyprotodonts, concluding that "While there is undeniably a series of forms connecting *Cænolestes* with the Diprotodonts in tooth structure, yet *Cænolestes* itself is so generalized in this respect that we may perhaps, in the absence of corroborating characters, question its inclusion within this group. Possibly it may be found to be an offshoot from the Polyprotodonts, as it appears structurally to be more generalized than any Diprotodont, and therefore it might well occupy a separate suborder, as Thomas suggested—the Paucituberculata of Ameghino."

Gregory (1910, p. 211) refers to Miss Dederer's work and concludes that "the detailed characters of the skull show no striking Diprotodont characters and the writer is inclined to regard *Cænolestes* and its allies as an independent suborder, an offshoot of primitive Polyprotodonts, which has paralleled Diprotodonts in certain characters of the dentition." Accordingly he places *Cænolestes* in the suborder Paucituberculata with the parenthetical suggestion that the group might be called the *Cænolestoidæ*. Gregory also discusses the possible relationship of the *cænolestids* to *Propolymastodon* and allied forms regarded as multituberculates by Ameghino. He concludes his treatment of the marsupial group by stating that "The problem of the genetic relations of the Diprotodontia and the Polyprotodontia is complicated to a certain extent by the existence of the *Cænolestoids*; but the opinion may be expressed that probably the resemblance of certain *Cænolestoids* to the Multituberculates is an instance of convergence between related suborders, and that the same is true, but to a less extent, of the resemblances of other *Cænolestoids* to the Diprotodont phalangers." Subsequently (Osborn, 1910, pp. 515-518) the classification proposed by Gregory was somewhat modified and the *cænolestids* were included in the suborder Diprotodontia as a superfamily, the *Cænolestoidæ*.

Broom (1912), in his special paper "On the Affinities of *Cænolestes*," reviewed the evidence adduced by Miss Dederer and pointed out certain resemblances to polyprotodonts not previously noted. He concluded that "as *Cænolestes* differs from the typical Polyprotodonts only in tooth specialization, it should not be removed from the Polyprotodontia, but merely be made the type of a distinct family, or section at most."

Gidley (1915), in a short footnote, expresses an opinion similar to that of Broom, saying: "The Cænolestidae have been placed in this great group [Diprotodontia] apparently on the diprotodont-like development of the lower jaw. However, this may be an entirely independently acquired character. This family more probably belongs with the Polyprotodonts."

With the exception of casual references in general works, nothing else of importance has appeared in reference to the systematic or phylogenetic position of *Cænolestes*. The occurrence of the animal in several different localities has been recorded (Osgood, 1912; Stone, 1914) and the number of species has been increased at least nominally to three, one of which has been made the type of a second genus, *Orolestes* (Thomas, 1917), which, as described, seems only slightly different from the original form.¹

Reviewing the foregoing it is seen that, within two decades and upon a knowledge of its cranial and external characters only, *Cænolestes* has been placed in three different suborders and that various competent authors have disagreed as to its proper position. Thomas, Ameghino and Scott frankly classify it with Australian diprotodonts; Sinclair inclines to do the same; Bensley, Miss Dederer and Gregory believe its diprotodont characters are convergent and place it in a suborder of its own; while Broom and Gidley would rank it merely as an aberrant polyprotodont. Beginning with that of the period prior to the discovery of *Cænolestes*, the following are the various classifications of marsupials concerned in its history, wholly extinct groups being indicated by a dagger:

THOMAS 1888.

Order MARSUPIALIA

Suborder *Diprotodontia*

- Family 1. Macropodidae
- 2. Phalangeridae
- 3. Phascolomyidae

Suborder *Polyprotodontia*

- Family 1. Peramelidae
- 2. Dasyuridae
- 3. Didelphyidae

¹ At the time of writing, the privilege of examining specimens of *Orolestes* is denied me.

THOMAS 1895.

Order MARSUPIALIA

I. Suborder *Diprotodontia*

A. Non-syndactylous. American

- Family 1. Epanorthidae *Cænolestes*
 †*Epanorthus*
 †*Decastis*
 †*Paraepanorthus*

B. Syndactylous. Australian

- Family 2. Phalangeridae
 3. Phascologyidae
 4. Macropodidae

II. Suborder *Polyprotodontia*

A. Syndactylous. Australian

- Family 5. Peramelidae

B. Non-syndactylous. American and Australian

- Family 6. Didelphyidae
 7. Dasyuridae
 8. Notoryctidae

TROUESSART 1898.

Order MARSUPIALIA

I. Suborder *Diprotodontia*

Section 1. *Syndactylia*

- Family 1. Phalangeridae
 2. Phascologyidae
 3. †Diprotodontidae
 4. Macropodidae

Section 2. *Asyndactylia*

- Family 5. †Abderitidae
 6. †Epanorthidae
 7. †Garzonidae
 8. Cænolestidae (*Cænolestes* only)

II. Suborder *Polyprotodontia*.

- Family 9. Peramelidae
 10. †Borhyaenidae
 11. Dasyuridae

- 12. Notoryctidae
- 13. †Microbiotheridae
- 14. Didelphyidae
- 15. †Amphitheridae
- 16. †Triconodontidae
- 17. †Dromatheridae

AMEGHINO 1903.

Subclass MARSUPIALIA

Superorder DIPROTODONTA

Order PLAGIAULACOIDEA

Suborder †*Allotheria*

Family †Plagiaulacidae

†Polydolopidae

†Neoplagiaulacidae

†Promysopidae

†Polymastodontidae

Suborder *Paucituberculata*

Family †Abderitidae

†Epanorthidae

Cænolestidae — *Cænolestes*† — *Zygolestes*

†Garzonidae

†Microlestidae

Order HYPSPRYMNOIDEA

Australian diprotodonts

Order RODENTIA

Rodents. Equivalent to Glires

Superorder POLYPROTODONTA

Australian and American polyprotodonts

WEBER 1904.

Subclass MARSUPIALIA

Order MARSUPIALIA

Suborder *Polyprotodontia*

Family Didelphyidae

Dasyuridae

Notoryctidae

Peramelidae

Suborder *Paucituberculata*

Family Epanorthidae (*Cænolestes et al*)

Suborder *Diprotodontia*

Family Phascolarctidae

Phalangeridae

SINCLAIR 1906.

Suborder *Diprotodontia*

Family *Cænolestidae*

Subfamily Cænolestinae

Genus †Halmarhiphus

†Garzonia

Cænolestes

Subfamily †Palaeothentinae

Genus †Palaeothentes

†Callomenus

†Decastis

Subfamily †Abderitinae

Genus †Abderites

GREGORY 1910.

Infraclass METATHERIA

Order †TRICONODONTA

Order †TRITUBERCULATA

Order MARSUPIALIA

Suborder †*Allotheria* (Multituberculata)

Suborder *Diprotodontia*

Suborder *Paucituberculata* (Cænolestoidea)

Suborder *Polyprotodontia*

OSBORN 1910.

Order MARSUPIALIA

Suborder *Polyprotodontia*

Superfamily DIDELPHOIDEA

Family 1. Didelphiidae

2. Myrmecobiidae

3. Dasyuridae

4. Thylacinidae

Superfamily PERAMELOIDEA

Family 1. Peramelidae

Superfamily NOTORYCTOIDEA

Family 1. Notoryctidae

Suborder *Diprotodontia*

Superfamily CÆNOLESTOIDEA

Family 1. Palæothentidae—†*Palæothentes*—†*Abderites*— *Cænolestes*

2. Garzoniidae

Superfamily PHALANGEROIDEA

Family 1. Phalangeridae

2. †Thylacoleonidae

3. Macropodidae

4. Phascolomyidae

5. †Diprotodontidae

Suborder *Allotheria*

Family 1. †Plagiaulacidae

SCOTT 1913¹

Order MARSUPIALIA

Suborder *Polyprotodontia*

Didelphiidae

Thylacynidae

Suborder *Diprotodontia*

Cænolestidae

†Garzoniidae

Suborder *Allotheria*

†Plagiaulacidae

†Polydolopidae

DISTRIBUTION.

The present known distribution of *Cænolestes* (and *Orolestes*) is shown by the accompanying map. Only a few localities are represented, all Andean, and extending from the Venezuelan border on the north nearly to the Bolivian boundary on the south. In the extensive regions between these localities, no specimens have been obtained but it is not improbable that the animal has nearly or quite continuous range throughout the central Andes so far as local conditions meet its needs.

¹ American forms only.

X Records of *Cænolestes* and *Orolestes*

The original type of *Hyracodon fuliginosus* was received from the well-known collector Louis Fraser in a shipment from Ecuador of which Tomes (1860, p. 211) says: "The greater portion of these are believed to have been collected at Pallatanga, on the western slope of the Cor-

dillera; but the exact locality is not certain, from the specimens having been unfortunately mixed together." The elevation of Pallatanga is scarcely more than 1500 meters and, since all exact records of *Cænolestes* are from greater altitudes, it is probable that the first specimen did not come from there but from the paramos of Mt. Chimborazo or Mt. Pichincha where Fraser also worked. The type of *Cænolestes obscurus* likewise came from an indefinite locality—the vicinity of Bogota, Colombia. In this case the probable actual locality is at the edge of the paramos north or east of the city of Bogota in the eastern Cordillera of the Andes. Exact records are only three in number: Paramo de Tama, Venezuelan—Colombian Boundary at 7000-9000 ft. (Osgood, 1912); Hacienda Garzon at 10500 ft., near the Paramo of Mount Pichincha, Ecuador (Stone, 1914); and Torontoy, Peru, at 14000 ft. between Lats. $10^{\circ} 30'$ and $13^{\circ} 30'$ S. and Longs. 72° and 73° W. (Thomas, 1917).¹

HABITS.

Cænolestes is of terrestrial and crepuscular or nocturnal habits. It feeds upon insects. These statements compass in a large measure what is known of its habits. It lives at high altitudes in cool forests not far from timberline and may be found also in grassy openings in mountain valleys. On the Paramo de Tama, near the head of the Tachira River, I obtained it at an altitude of about 7500 feet on both the Venezuelan and Colombian sides of the international boundary line which is here formed by the Tachira River. Most of the specimens were secured in the heart of a dense forest on the Colombian side of the river not far from a hacienda belonging to Don Mario Gonzales of San Jose de Cucuta, Colombia. Wishing to collect and observe in such a place under wholly natural and undisturbed conditions, I had employed men for some days to cut a new trail directly into the deepest part of this forest for a distance of several miles. Camp was established at the terminus of this main trail and short radiating trails were cut thence to various points of the compass, the forest and undergrowth being so dense that but little freedom of movement would have been possible otherwise. In the dense growth along these trails, *Cænolestes* were caught in small numbers, rarely more than one specimen to fifty "trap nights." The exact nature of the places in which individual specimens were taken varied widely, indicating that the animals move about freely. Some were caught in steel traps baited with small birds, others in rat traps baited with bacon rind,

¹ In a paper published since this was written, Thomas (1920, p. 246) gives three additional locality records: Machu Picchu, Peru, alt. 12000-13000 ft.; Ocabamba Valley, Peru, alt. 9100 ft.; Guala, Ecuador, alt. 6000 ft.

and still others in mouse traps with the conventional rolled oats as bait. In the case of the mouse traps, however, it was probable that the animals had not been attracted by the bait but had merely run across the lightly set traps. A preference for animal food was quite evident and it was only by continued effort with "meat baits" that a total of more than one or two specimens was obtained.

Beneath the heavy canopy of treetops and still further shaded by masses of low vegetation, the animals of the dark damp forests exercise their vision only to a limited degree and depend mainly upon the sense of smell. That this is the case with *Cænolestes* is indicated by its small eyes and its highly developed olfactory organs. In fact, although somewhat ratlike in appearance and from its rather long hind legs doubtless active and roving, *Cænolestes* is best described as to appearance by saying that it is a large forest shrew. Like a shrew, it is fond of fresh meat when such food offers itself, but its steady diet consists only of insects. The contents of three stomachs were submitted to the U. S. Biological Survey and as a result of examinations kindly made in their laboratories the following report was received:

- "No. 1. Contents: Parts of a weevil, three ants (Dorylidae), skin of caterpillar, lepidopterous pupa, adult lepidopteron, leg fragments of orthopteron, a Tipulid larva, centipede and spider. Dipterous and lepidopterous remains form the major portion.
- "No. 2. Contents: Fragments of two beetles and a weevil, a caterpillar, an orthopteron, a hemipteron (?), spider and one seed of composite plant. Caterpillar—at least 60 per cent.
- "No. 3. Contents: Unidentified insect fragments and parts of spider."

Other observations on the habits of *Cænolestes* so far as published are meager. Mr. Geo. D. Child, through whom the type of *C. obscurus* was obtained, contributed the following note: "The little animal you speak of is called 'Raton Runcho,' which means 'Opossum-Rat.'¹ It lives in the high brush-wood, and is supposed to feed on birds' eggs and small birds. It is very rare indeed, and is obtained with much difficulty." (Thomas, 1895, p. 877.)

Mr. Samuel N. Rhoads, who collected two females on Mt. Pichincha, Ecuador, writes that "The two specimens were secured in swampy ground, the edge of a large pasture on the Hacienda Garzon, within a few feet of a swiftly flowing stream of considerable size. They were caught in small cyclone mouse traps set in underground runways among the thick grass, these runways being about on the level with the waterline of the swamp. They were caught on the same day, soon after

¹ The book name *selva* was proposed by Lydekker in 1896 (Geog. Hist. Mamm.), but has seldom been used by others.

placing the traps in that locality, but although I continued to trap there for a week longer, having as many as 40 or 50 traps in that place, I secured no more specimens there, nor in any other similar localities where trapping was done. The stream alluded to runs over a bed strewn with volcanic rocks and boulders and is in an open cultivated valley-head, draining the south slopes of Mount Pichincha, about 8 miles south of Quito and at an elevation of about 10,500 feet, the valley at this point being about half a mile wide and extending to even greater widths as far as one can see, in a southerly direction." (Stone, 1914, p. 18.)

The type of *Cænolestes* (or *Orolestes*) *inca* was collected by an experienced naturalist, Mr. Edmund Heller, whose notes, when published, will doubtless furnish substantial additions to what is now known of the animal.¹

EXTERNAL CHARACTERS.

General form.—Although frequently referred to as "ratlike," the general form in *Cænolestes* might with equal propriety be called soricine. Insectivorous forms such as *Nesogale* approach it closely and, as elsewhere shown (Pl. XIX), have practically identical major proportions. Among marsupials, the species most closely resembling it are perhaps to be found in the genus *Phascologale*. Many species of *Marmosa* approximate it in size of body, but distinctions in the feet, ears, and tail are so obvious that the parallel is at once seen to be only of the most general nature. The legs are of moderate length, the hind pair 20 to 25 per cent longer than the front.

Head.—The head is elongate conical and somewhat shrewlike in general appearance. The eyes are very small, the distance between the anterior and posterior canthi measuring only 2.2 mm. in a female specimen in alcohol. The rounded ears project well above the pelage but are relatively small as compared to those of the didelphids. The thickness of the cartilaginous conch also is greater and it is set with fine short hairs within and without. The tragus is but little developed and the considerably larger antitragus is opposed mainly by the crus of the helix which forms a thin flap directed inward. Two parallel antihelical folds (metatragus) are prominent and separated from each other by a deep fossa (Pl. II, fig. 1). A similar condition is found in *Perameles* (cf. Thomas, 1888, Pl. XXI, figs. 5, 7). The nostrils are lateral and bounded by a rhinarium which is extended dorsally to form a small v-shaped plate. A shallow furrow traverses it anteriorly and dorsally.

¹I am informed by Mr. Heller that these and other notes on Peruvian mammals have been prepared for publication by the National Geographic Society, Washington, D. C.

The upper lips have a thin coriaceous edge that continues to the median sulcus which partially divides the rhinarium and forms an incipient cleft of the lips. The lower lips have a similar narrow edge which is only very slightly thickened in front. The lower lip is free in front from the base of the long incisors and has a high internal median ridge or frenulum which fits into the slight space between the incisors. Slightly in front of the angle of the mouth and just in advance of the molariform teeth is a pair of reciprocating labrets or outgrowths of the lips, one on the upper and the other on the lower lip (Pl. II, fig. 2). These appear to be an adaptation for the ejection of undesirable food particles during mastication. They are directed slightly outward and downward in such a way that with the lips wholly or partly closed in front they might in conjunction with the buccinator muscle serve to guide hard parts of insects or other matter to the exterior. A quite different function also is possible, that is, in closing the lips at this point and assisting in isolating the internal cheek pouch formed by the buccinator muscle. This pouch is fairly definite and includes a considerable space laterad of the molariform teeth.

Tail.—The tail is gently tapering and has no inordinate thickening at the base. The body fur extends upon it only a very slight distance. Thence to the tip it is thickly set with short stiff hairs which nearly or quite conceal the underlying scaly annulations. At the tip there is usually a slight hairy pencil. Nothing suggests prehensilism in the slightest degree.¹ The contour of the tail is rather distinctly quadri-lateral instead of cylindrical.

Feet.—The fore feet are pentadactyl, but the outer toes are decidedly reduced and furnished with small blunt nails instead of claws. The third or middle toe is slightly the longest and the second and fourth are subequal, all three bearing sharp curved claws. The reduction of the lateral toes and especially the absence of claws is in marked contrast to the condition general among Dasyuridae and Didelphiidae but is almost exactly paralleled in the Peramelidae in many of which the reduction of the first and fifth toes has proceeded much farther. In *Cænolestes*, however, although the elongated middle toes seem to indicate a digitigrade adaptation, the plantar surface is carried well backward on the outer side and supported by an unusually large pisiform bone. There are five plantar pads well distinguished from each other and having smooth non-striated surfaces.

The hind foot is relatively long and narrow. Its general proportions

¹Thomas (1895), in describing *C. obscurus*, states that the tail has "its terminal inch below wholly naked" and that "it is therefore presumably prehensile." No such condition has been observed in other specimens and it is probable the tail in the above-mentioned one was abnormal.

might be said to lie somewhere between those of *Phascologale* and *Antechinomys*. The hallux is reduced and bears only a weak nail. It is set at a slight angle but it scarcely reaches the end of the second metatarsal and its functional importance is slight. The remaining four toes are subequal, the third and fourth possibly a trifle longer than the second and fifth. All are furnished with well-developed curved claws. The third and fourth digits are connected at the base by an integumentary web which is slightly more extensive than that between the other digits, but this is scarcely to be regarded as a tendency toward syndactylism. The soles are naked and provided with six plantar pads in much the same relative positions as in *Phascologale*. Three of them are digital, one hallucal, and two metatarsal. All are smooth without transverse striations. See Plate II, figures 3 and 4.

Pelage.—The pelage is soft and thick over the entire body. Owing to the peculiar distribution of hairs of different quality the coat is not smooth or velvety but has an appearance of superficial looseness and unevenness not uncommon in mammals of humid regions. The color is practically uniform throughout, the underparts being only slightly lighter than the upper owing to the absence of the fine exserted blackish hairs which are generally distributed on the upperparts. The claws and a few hairs at their bases are whitish but the hairiness of the upper sides of the feet shows no contrast with the body. The tail also is colored the same above and below and from base to tip except that in four out of eleven specimens the tip is white. There are no facial markings and in general it would be difficult to find a more uniformly colored animal. The color is dull Mars brown, clearer on the underparts and more mixed with darker and subject to light reflections on the upper parts.

Marsupium.—No trace of an external pouch can be detected in adult specimens of either sex and no immature examples have been examined. Tomes (1863), in describing the original type of *Hyracodon fuliginosus*, which was probably immature, states that it had a "small and rudimentary pouch." Hence it is probable that the pouch in *Cænolestes*, as in *Phascologale* and some others, is present in the young but disappears in the adult.

MEASUREMENTS.

External measurements.—Fresh specimens measured in the field: Average of 5 males: Total length 240.6 (235-251); head and body 119.2 (113-135); tail vertebrae 121.4 (118-126); hind foot with claw 23.5 (23-24.5). Of 5 females: 223 (209-230); 107.6 (106-113); 115.4 (103-121)

22.5 (22-23). The ear in a female in alcohol measures 12.5 mm. from the intertragal notch to the apex. The tail at the base is 3.3 mm. wide and 3 mm. deep; at 10 mm. from the tip it is 1 mm. x 1 mm.

Measurements of skull.—Adult male and female, respectively: Greatest length 32.7, 29.2; basilar length 28.4, 25; zygomatic breadth 15, 13.8; mastoid breadth 11.4, 10.9; depth of braincase 8.2, 8.1; length of nasals 15.1, 14; greatest breadth of nasals 4.3, 4.3; least interorbital breadth 7.3, 7.3; length of palate from gnathion 18.3, 16.2; anterior palatal foramina 6.3 x 2.5, 5.4 x 2.2; palatal vacuities 6.7 x 3.2, 6 x 3.2; front of upper canine to back of M⁴ 12.9, 11.5; combined length of four upper molars 5.7, 5.5; combined length of three lateral incisors 3.1, 2.8; length of bone of lower jaw from condyle 20.5, 17.5; angle to coronoid 8, 7.2; depth of jaw at second molar 2.2, 1.7; combined length of four lower molars 6.6, 6.3; exposed length of median lower incisor 6, 5.

MYOLOGY.

Plates III-VI.

MUSCLES OF THE SKIN.

Panniculus.—Immediately beneath the skin and separable from it only by careful dissecting is a thin, partly membranous and partly muscular layer which may be divided into several rather definite parts but which taken as a whole forms a sac enveloping the entire body and having relatively few and weak ental attachments. On the fore and hind legs it becomes exceedingly thin and cannot be traced beyond the middle of the epipodials. Over the posterior part of the body it is entirely membranous, thicker over the back and sides and very thin over the pectoral muscles. It is attached posteriorly on the *biceps femoris* lengthwise of its caudal fibers; dorsally it becomes slightly muscular and dips beneath the "cruro-coccygeus" to a definite insertion on the dorso-ental surface of the tuberosity of the ischium.¹ It is almost discontinuous over the pectoral and interscapular regions where it is lost in a thin scarcely perceptible aponeurosis. Over the head it reappears as a thin sheet chiefly membranous except along the under side of the ear and forward on the side of the face to the anterior base of the zygoma where a few muscle fibers run. The principal muscular division proceeding directly from this outer plane, superficial fascia, or *panniculus* is that which enters the axilla forming the so-called *cutaneus maximus*. Be-

¹ Possibly this dorsal part corresponds to the "ischiotergal" slip described by Maculister in *Dasypus* and by Wilson (1894, p. 6) in *Notoryctes*.

ginning in the middorsal line near the origin of the *latissimus dorsi* (and immediately overlying and parallel to it), a thin sheet of converging fibers runs forward and downward to the axilla. Similar converging fibers from the mid-thoracic region and from the abdomen join those from above and unite to form a single muscle which enters the axillary region dorso-laterally just below the *latissimus dorsi* and ventrally parallels the outer edge of the *pectoralis major*. It makes a turn on itself and inserts as usual on the proximal two thirds of the inner edge of the deltoid ridge of the humerus. The abdominal part of this muscle is somewhat differentiated into a *humero-abdominalis* or *xiphi-humeralis* (*pectoralis quartus*). This begins to be fleshy on the middle of the abdomen and anteriorly divides to pass on either side of the *pectoralis* and parallel to it. On entering the axilla it becomes fused with the *cutaneus maximus* and has a common insertion with it. It has similar relations in most marsupials.

Cervico-auricularis.—This is a thin paired sheet arising from the occiput and the nuchal crest for a distance of about 10 mm. or to the fatty deposit in the interscapular space. It is inserted on the cartilage of the back of the ear and into the ental surface of the superficial fascia below the ear for some 8 mm. lengthwise of the side of the neck. Dorsally it leaves the base of the ear for a short distance and attaches to the ectal surface of the muscle which elsewhere underlies it and which is here called the *auriculo-occipitalis*.

Auriculo-occipitalis.—Arises from the nuchal crest entad of the *cervico-auricularis*, the anterior limit of its origin being even with that of the *cervico-auricularis* and the posterior slightly cephalad of it. Its two halves separate on the occiput and diverge over the parieto-squamosal region to overlie the caudal part of the temporal muscle. Directly above the ear, it passes into the ental surface of the superficial fascia or *panniculus* and thence around the anterior base of the ear the two are inseparable.

Dorso-cuticularis.—A long slender muscle arising on the side of the twelfth thoracic vertebra and running forward along the boundary between the *latissimus dorsi* and the *trapezius* to the side of the scapula where it widens slightly and inserts on the ental surface of the thin outer sheet of the *panniculus*. Its width at its insertion is 1.5 mm. I have been unable to find any trace of this muscle in *Didelphis* nor any record of its occurrence in any other marsupial. Its form and relations are almost exactly as in the Insectivora where it is of frequent occurrence. Somewhat similar muscles are found also in certain edentates. (See Plate III.)

MUSCLES OF THE BACK.

Spino-trapezius.—Origin from the spines of the thoracic vertebrae from the fifth to the twelfth. Insertion on the middle of the vertebral third of the scapular spine. A thin rather elongate muscle passing forward superficial to the dorso-cephalic part of the *latissimus dorsi*. Dorsally it is separated from the *acromio-trapezius* by a small fascia overlying the *rhomboideus*. A separation has been noted also in *Notoryctes* (Wilson, 1894, p. 6). It is the usual condition in the monotremes, insectivores and certain edentates. In *Didelphis*, the single muscle is weaker over the shoulder than elsewhere and in *Sarcophilus* (Macalister) also there is a tendency toward division. In *Petrogale*, Parsons (1896) found that it had a "continuous fleshy origin except opposite the first thoracic spine, where it is aponeurotic."

Acromio-trapezius.—Origin from the occipital crest slightly laterad of the median line and thence along the nuchal crest to the vicinity of the fifth thoracic spine. Insertion on the front of the scapular spine and metacromion and thence along a fascia crossing the acromio-deltoid to the clavicle. A broad thin sheet covering the dorso-caudal half of the side of the neck and part of the shoulder.

Latissimus dorsi.—Origin from the vicinity of the fifth thoracic spine along the vertebral column to the middle of the third lumbar vertebra and to the lumbar fascia. Insertion in common with that of the *teres major* entad of the *biceps* on the side of the bicipital groove. On its lower side near its insertion is the usual connection of the *epitrochlearis*. It is more definitely connected to the *teres* at its insertion than in *Didelphis*. It has no costal slips and in this agrees with *Didelphis*, *Dasyurus*, *Notoryctes*, *Phascolarctos*, etc.

Rhomboideus.—This is divisible into *Rhomboideus major (vertebralis)* and *R. minor (capitis)*. *R. major* has its origin on the nuchal crest from the occiput slightly caudad of the anterior border of the trapezius to the fifth thoracic spine. Its insertion is on the inner side of the vertebral border of the scapula cephalad of the insertion of the *serratus magnus*. *R. minor* has its origin fleshy from the junction of the mastoid, squamosal and supraoccipital bones and thence along the occipital crest a distance of about four millimeters. It is a thin flat ribbon free from the vertebral division until a point halfway between the occiput and the scapula where it unites with its fellow. Its insertion is on the anterior distal base of the scapular spine dorsad of the insertion of the "*atlanto-scapularis*." The *rhomboideus* in marsupials is usually undivided.

Levator scapulae.—Origin from the transverse processes of the last five cervical vertebrae just entad of the *scalenus*. Anteriorly a single slip

passes ectad over the *scalenus* from an origin on the superior transverse process of the third cervical. This arrangement differs from that of *Didelphis* in which the *scalenus* is entirely laterad of the *serratus*. Insertion on the posterior vertebral border of the scapula cranial of the insertion of the *serratus* and continuous with it.

Serratus posterior superior.—Insertion by digitations from the fourth to the ninth ribs inclusive. Dorsally it is mostly aponeurotic but a broad fleshy prolongation extends anteriorly to the lower border of the *splenius*, thence becoming aponeurotic as it crosses that muscle and continues between the *splenius* and the *rhomboideus* to the most anterior point of its origin on the nuchal crest above the sixth cervical spine. Thence caudad its origin follows the mid-dorsal line and its wide aponeurosis covers the *longissimus dorsi* and continues into the lumbar fascia.

Serratus posterior inferior.—Origin on the middorsal line with the lumbar fascia. Insertion from the tenth to the thirteenth ribs. Practically as in *Didelphis*.

Splenius.—Origin from the nuchal crest from the vicinity of the front of the axial spine to the second thoracic vertebra. Insertion on the occipital crest from the mastoid process of the squamosal to a point about 2 mm. from the median line. Its lower border is free as usual and near its caudal extremity it is somewhat attached to the aponeurosis of the *serratus posterior superior*.

Biventer cervicis.—Origin from the transverse processes of the fifth, sixth, and seventh thoracic vertebrae. Insertion fleshy or very slightly aponeurotic on the occiput between the median line and the *complexus* to which its lateral border is attached for some 4 mm. Anteriorly it reaches nearly to the parietal bones. It is a larger and thicker muscle than in *Didelphis* and lies with its counterpart for a great part of its length in the space between the elevated axial spine and the second thoracic spine.

Complexus.—Origin from the articular processes of and the aponeurotic arches between the last six cervical and the first five thoracic vertebrae. A large slip somewhat connected also with the *longissimus capitis* runs forward to the transverse process of the atlas. Insertion by a broad tendon on the occipital crest laterad of the *biventer* to within about 3 mm. of the mastoid bulla.

Longissimus capitis.—Origin from the articular processes of the last six cervical and the first four thoracic vertebrae in intimate relation with the slips of origin of the *complexus*. Insertion by broad tendon on the upper cranio-lateral border of the mastoid bulla and to a slight extent on the adjoining part of the squamosal.

The *biventer*, *complexus*, and *longissimus capitis* are well distinguished from each other and in their general relations are as in *Didelphis*. At least one distinct tendinous inscription crosses the anterior part of each some 4 mm. from the insertion.

Rectus capitis posterior superficialis.—Origin from the crest of the axial spine. Insertion entad of the *complexus* on the occipital crest. It runs downward and forward with free upper and lower borders, leaving a triangle on either side of the median line in which the much smaller halves of the *rectus minor* and part of the *rectus major* are exposed.

This muscle, often found in carnivores, is not generally reported for marsupials, although Parsons (1896, p. 694) reports the *rectus major* as bilaminar in *Petrogale*. It is thin and flat and covers a great part of the *rectus capitis posterior major*. In *Didelphis*, it is only partially separable from the *rectus major*, in *Marmosa* it is somewhat freer, and in *Cænolestes* it is quite independent.

Rectus capitis posterior major.—Origin from the craniad surface of the axial spine and intermuscular septa with the *obliquus inferior*. Insertion on the occiput from the base of the paroccipital process dorsad along the caudal border of the mastoid bulla to a point on the occipital crest midway to the median line. It meets its fellow of the opposite side in front of the axial spine and the two have tendinous connection there and to a slight extent also with the caudal border of the neural arch of the atlas. Its under surface has relation with the *rectus capitis posterior minor* and with the surface of the neural arch of the atlas. It is a rather thick muscle only slightly smaller than the *obliquus inferior* with which its lower-border is in apposition.

Rectus capitis posterior minor.—Origin from the straight anterior border of the neural arch of the atlas. In a female specimen this measures 3 mm. in length from side to side. Insertion on the occipital from a point near the median line nearly to the mastoid bulla. Its outer surface has relation with the *biventer*, *complexus*, and *rectus posterior major*. At the median line of the occiput it is separated from its counterpart by 1–2 mm. The two muscles approach each other and have tendinous union in their caudal halves.

Obliquus capitis superior.—Origin from the lateral surface of the transverse process of the atlas. Insertion tendinous on the occipital ridge immediately dorsad of the mastoid bulla. A short flat muscle crossing nearly at right angles to the large *obliquus inferior* and *rectus capitis posterior major* which has relation with its inner surface.

Obliquus capitis inferior.—Origin from practically the whole lateral surface of the axial spine. Insertion on the whole caudal surface of the transverse process of the atlas. It is a thick heavy muscle running

downward and forward. In *Didelphis*, this muscle extends to the third, fourth and fifth cervical spines, but in *Marmosa* it is much as in *Cænolestes*.

Rectus capitis anticus major.—Origin on the transverse processes of the last five cervical vertebrae. Insertion on the posterior third of the basisphenoid and the anterior two fifths of the basioccipital slightly laterad of the median line. Some of its superficial fibers extend caudad with the *longus colli* to the ventral side of the posterior thoracic vertebrae. Anteriorly it forms a good sized bundle and is not peculiar except for the extent of its attachment to the basisphenoid which is more than usual.

Rectus capitis anticus minor.—Origin on the middle third of the ventral surface of the neural arch of the atlas midway between the median tubercle and the transverse process. Insertion on the ventral surface of the basioccipital in its anterior lateral fourth. A thin flat muscle lying entad of the *rectus anticus major* and mediad of the *rectus capitis lateralis* but entirely free from both.

Rectus capitis lateralis.—Origin on the transverse process of the atlas. Insertion slightly tendinous on the outer edge of the basioccipital along the carotid canal and thence forward to the basisphenoid and by tendinous fascia to the presphenoid. Certain lateral fibers attach on the periotic. It is a well developed muscle and entirely distinct, extending farther cephalad than in *Didelphis*.

Longus colli.—The dorsal part of the *longus colli* is rather distinct, extending from the transverse process of the sixth cervical along the bases of the ribs and the sides of the vertebrae to the fourth dorsal vertebra. Between its two halves are fibers which extend into the cervical part but they are lateral in position and the median ventral surfaces of the dorsal vertebrae are mostly covered only with tendinous fascia. The cervical part consists of various converging slips from the transverse processes of the cervical vertebrae and caudally from the sides of the ventral surfaces of the vertebrae. Over the atlas it shows six rather distinct divisions, two medial ones inserting on the tubercle of the atlas and two on either side inserting on the body.

Spinalis dorsi.—The *spinalis dorsi* disengages from the medial side of the *longissimus* about opposite the eighth thoracic vertebra, the caudal part of the *biventer cervicis* forming a wedge between the two. Its inner fibers run to the spines of the anterior thoracic vertebrae but its lateral fibers form a practically distinct subvertical ribbon which runs forward and inserts on the side of the low spines and bodies of the fourth to the seventh cervicals and the first thoracic. It passes the high spine of the second thoracic without definite attachment, but between the

first thoracic and the tip of the spine of the second it is traversed by a strong tendinous inscription which may be partially attached to either or both of the vertebrae. Anteriorly it passes between the lateral division of the *semispinalis cervicis* and the long interspinal which connects the spines of the fifth cervical and the second thoracic. In the opossum the *spinalis* inserts as high up as the third cervical and the outer surface of its anterior part is wholly in relation with the *complexus*.

Semispinalis cervicis.—In the region between the axial spine and the second thoracic spine are two paired muscles which run ventrad toward each other and meet at an angle over the sixth and seventh cervicals, the posterior one passing inside the *spinalis* between the two halves of the anterior one. The anterior of these muscles has been regarded as the *semispinalis cervicis*. It arises on the bodies of the first thoracic and the fifth, sixth and seventh cervical vertebrae and converges to insert on the tip and side of the axial spine in opposition to the *obliquus capitis inferior*. Its outer surface has relation with the *complexus* and its inner with the *spinalis*. It is not apparent in *Didelphis*, but is present in *Marmosa*. The posterior paired muscle, above-mentioned, which is in the nature of an interspinal, may be described here also. Its origin is on the side of the spine of the fifth cervical and its insertion on the front of the tip of the spine of the second thoracic vertebra. It stretches directly between these spines without connection with the intervening vertebrae. Between its two halves the ligamentum nuchae passes to the vertebral spines. Beneath its caudal part, but distinct from it, a paired intervertebral connects the spines of the first and second thoracic vertebrae. Its outer surface has relation with the *biventer* and the *spinalis*, its inner with the ligamentum nuchae. A corresponding muscle is found in the opossum running to the sides of the thickened third cervical spine, its two sides therefore being widely apart and, although its actual relations are much as in *Cænolestes*, its appearance is quite different. In *Marmosa* the third, fourth and fifth cervical spines are not enlarged as in *Didelphis*. Consequently the arrangement of the neck muscles is more nearly as in *Cænolestes* and many other marsupials.

Longissimus.—Anteriorly this is divisible into *longissimus dorsi* and *longissimus cervicis*. Superficially the two divisions appear as one muscle but the outer part which continues caudad may be partially separated from an inner mostly concealed part which only extends caudad to the fifth thoracic vertebra and which may be regarded as the *longissimus cervicis*. It arises by slips from the laminae of the cervicals and from the transverse processes of the thoracic vertebrae from the first to the fifth. It inserts on the transverse processes of the cervical vertebrae from the second to the fourth or fifth, being inseparable at this

point from the *longissimus dorsi*. The *longissimus dorsi* separates from the *spinalis dorsi* about opposite the eighth thoracic vertebra and passes cranio-ventrad between the *iliocostalis* and the *biventer* and *longissimus capitis* from which it is entirely distinct. Its insertion is just caudad of that of the *longissimus cervicis* on the transverse process of the fifth and possibly also the fourth cervical vertebra. Its origin is from the lumbar tendinous sheet from which it becomes fleshy in the vicinity of the second lumbar vertebra. Passing forward it is conjoined with the *spinalis* and consists of slips from the transverse processes of the vertebrae.

Iliocostalis.—An anterior division or *iliocostalis dorsi* and a posterior or *iliocostalis lumborum* are fairly distinct. The *iliocostalis dorsi* arises by slips from the sides of the ribs from the first to the eighth. It passes cranio-ventrad on the side of the neck laterad of the *longissimus* and inserts on the superior transverse processes of the sixth and seventh cervical vertebrae. For a short distance just before its insertion it is merged with the *longissimus*.

The *iliocostalis lumborum* arises from the complicated lumbar tendons of the *sacrospinalis* and becomes fleshy opposite the fourth lumbar vertebra. It inserts by thin flat slips on the sides of the ribs from the sixth to the thirteenth. Between the ilium and the ribs it forms the anterior lateral boundary of the mass of muscle filling the space between the articular and transverse processes of the lumbar vertebrae. Over the ribs it is thin and the slips are fairly distinct from each other. Between the sixth and the ninth ribs it overlaps the *iliocostalis dorsi* which only reaches the eighth rib.

Sacrospinalis.—Origin from the crest and the ventro-medial surface of the cranial fourth of the ilium, meeting the origin of the *iliacus*, and from the tendinous lumbo-dorsal fascia. Insertion on the dorsal surfaces and caudal edges of the transverse processes of the lumbar vertebrae and the twelfth and thirteenth dorsals. A thick rounded bundle extending from the ilium to the ribs. Easily separable from the multifidae which lie mediad of it but covered by the tendinous fascia. It fills the space on the side of the vertebral column between the articular and the transverse processes of the lumbar vertebrae.

Quadratus lumborum.—Origin from the bodies of the 12th and 13th dorsal vertebrae and the lumbar vertebrae from the first to the fourth. Insertion on the tips and ventral surfaces of the transverse processes of all the lumbar vertebrae except the last. It lies somewhat flattened between the *psaos parvus* and the large rounded *sacrospinalis* and connects the series of transverse processes with each other by series of overlying tendons, the fibers from the under side of one running to the

upper side of the one succeeding. Anteriorly it is closely associated with the *psaos parvus*.

Psoas parvus.—Arises fleshy and also by tendinous heads from the first, second, third, fourth and fifth lumbar vertebrae. Thence it continues caudad as a broad flat tendon crossing the surface of the medial part of the *psaos magnus* and inserting dorso-cranial of the marsupial bone on the iliopectineal tubercle. In its cranial part it is closely connected with the *quadratus lumborum* which lies laterad of it. The caudal part of its fleshy origin on the fifth lumbar vertebra is overlapped by the *psaos magnus* which intervenes between it and the *quadratus lumborum* at this point.

Psoas magnus.—Origin from the sides of the bodies and the ventral bases of the transverse processes of the fourth, fifth, and sixth lumbar vertebrae and the cranial half of the first sacral. Anteriorly it reaches to the posterior epiphysis of the third lumbar. Insertion with the *iliacus* on the lesser trochanter and the adjacent inner surface of the femur nearly as far distad as the insertion of the *pectineus*.

MUSCLES OF THE HEAD.

Plates III-IV.

Masseter.—The superficial masseter has its origin on the lower side of the zygoma by a short stout tendon nearly opposite and slightly caudad to that of the inferior zygomaticus and in the vertical plane of the middle of the eye. It spreads over the lower part of the mandibular insertion of the deep masseter and covers the ventro-caudal part of the mandible with its insertion from the base of the coronoid process just caudad of the *buccinator* and along the lower side of the mandible to the angle and the *pterygoideus*. It does not reach the inner surface of the mandible and especially in its ventro-caudal portion is relatively weaker than in *Didelphis*. The deep masseter lies below it and is closely connected, but a division is evident. It has its origin on the inner and lower sides of the zygoma and the fascia over it and is intimately associated with the *temporalis*. It is broadly attached to the whole outer face of the coronoid fossa.

Temporalis.—This is in two layers. The superficial or outer temporal arises from the fascia along the middle of the parietal bone and across the caudal part of the underlying layer. Thence it spreads as a rather thin sheet over the deep temporal and narrows to pass into the orbital fossa and insert on the mandible at the anterior base of the ascending ramus. It is also attached superficially to the membrane between it and the eye. The deep temporal is more extensive; its origin runs from the

glenoid fossa across the mastoid bulla along the supraoccipital crest halfway to the median line and thence directly forward along the rather indistinct parietal and supraorbital crests to the plane of the posterior canthus of the eye. Its insertions are extensive on the coronoid process, mostly on its ental surface and its anterior edge with a few of the fibers from the mastoid origin going to its ectal surface.

Buccinator.—Relatively extensive and forming a pouch behind the angle of the mouth. Its origin is along the alveolar border of the maxillary from the vicinity of the infraorbital foramen to the anterior base of the zygoma and likewise along the ramus of the mandible from the front and base of the coronoid process forward to the vicinity of the mental foramen. It is attached to or merged with the *orbicularis oris* around the lips.

Pterygoideus internus.—Origin from the whole of the external pterygoid fossa from the carotid foramen cephalad, passing the tips of the pterygoids but attached to their bases and thence along the ridge to the dorsal side of the lateral tuberosity of the palatal ridge. Insertion broad on the inner surface of the angle of the mandible and by certain medial fibers running to the fascia over the tympanic. Medial of the internal pterygoid the small but distinct *tensor veli palatini* runs from the periotic and the edge of the basisphenoid to the hamular process of the pterygoid bone.

Pterygoideus externus.—Origin by two heads. The larger ventral one arises fleshy from the whole of the lateral base of the pterygoid bone laterad of the *pterygoideus internus*. The smaller dorsal head arises from the alisphenoid caudad and laterad of the sphenoidal fissure. The two heads spread over the convex surface of the alisphenoid between the sphenoidal fissure and the glenoid fossa and, after uniting as they pass the front of the tympanic, insert at the base of the mandibular condyle principally on the stylomandibular ligament.

The muscle is relatively large and broad, bulking fully two-fifths as much as the internal pterygoid. It is thus quite different from that of *Didelphis* in which it is single-headed and very weak. It is described (Macalister) in *Phascolarctos* as a "small rudiment crossing and inseparable from the internal" and in *Sarcophilus* as "exceedingly feeble"; in *Myrmecobius* also, it is weak and single (Leche); in *Charopus* (Parsons, 1903, p. 67), it is unusual and inserts on the inside of the mandibular ramus; in *Petrogale* (Parsons, 1896) and other kangaroos, it is small; for most other marsupials, data are lacking.

Mandibulo-auricularis.—The only prominent ear muscle seems to correspond to the one for which this name has been used by various authors. It is a small subcylindrical muscle arising in the fascia between

the superficial and deep masseters near the mandibular condyle and issuing between the masseters to an insertion on the anterior base of the ear.

Zygomatikus.—This is in three divisions all arising on the zygoma. Two of them originate from the lateral surface of the posterior half of the zygoma behind the eye. They are small well-formed fasciculi closely similar to each other and somewhat distinguished from the third division which lies below (entad) them. They run forward to the level of the angle of the mouth where they become tendinous and proceed thus for some distance to insertions in the tissue on the side of the upper jaw at the base of the whiskers.

The third division, to which some other name may be applicable, is similar in appearance but arises from the side of the lower anterior angle of the zygoma and its slender tendon is inserted into the skin of the inside of the upper lip in the vicinity of the base of the labret.

Retractor naris.—A slender muscle arising from the fascia between the eye and the zygoma. It crosses the outer surface of the anterior root of the zygoma and becomes a very slender threadlike tendon which runs forward to an insertion on the nasal cartilages.

Digastricus.—Arises on the paroccipital process between the condyle and the mastoid bulla and runs down around the ear and thence forward to spread over the interramal space and attach along the sides of the mandible from the base of the angular process to the symphysis. Anteriorly it is largely aponeurotic. A tendinous slip diverges at a point ventrad of the masseter and passes dorsad to the caudal median border of the mylohyoid. It is essentially as in *Didelphis*.

MUSCLES OF THE NECK.

Plates IV, VI.

Sterno-hyoideus.—This is the principal muscle exposed on removing the skin from the lower side of the neck. It arises on the ental surface of the second sternebra and runs directly forward opposite its mate and inserts on the posterior side of the hyoid bone caudad to the insertion of the genio-hyoid. Some of its superficial fibers inosculate with those of the genio-hyoid. Like the other muscles connected with the hyoid and the tongue, it is not peculiar.

Genio-hyoideus.—A flat paired muscle immediately cephalad of the *sterno-hyoideus* and entad of the *mylo-hyoideus*. It arises on the body of the hyoid bone and inserts on the mandibular symphysis.

Sterno-thyroideus.—A thin, paired muscle lying between the trachea and the *sterno-hyoideus*. Laterally it is thin and transparent and toward the median line it is somewhat thicker, especially in its caudal

portion where its two sides become fused. It arises on the sides and ental surface of the second sternebra and inserts on the thyroid cartilage.

Mylo-hyoideus.—A very thin, paired, transverse muscle between the mandibles and immediately entad of the *digastricus*. It arises from the inside of the mandible near the alveolar border and has no bony insertion, merely meeting its fellow in a raphe in the median line and becoming aponeurotic anteriorly and posteriorly. Along the dorsal side of this raphe it is attached to the *genio-hyoideus*. Its posterior fleshy border runs from the angle of the mandible forward to the median line opposite the base of the angular process.

Hyoglossus.—Arises laterad and dorsad of the *genio-hyoideus* on the ventrocaudal border of the thyrohyals and basihyal cephalad of the insertion of the *thyrohyoideus*. It passes ventrad of the ceratohyals and free from them, diverges to enclose the genioglossus, and passes dorsad and cephalad into the body of the tongue.

Genioglossus.—Well-developed and having the usual relations, arising on the sides of the mandibles near the symphysis and forming a wedge between the two halves of the *hyoglossus*.

Styloglossus.—Runs laterad of the *hyoglossus* and merges with it about midway of the mandible. Its origin was not determined, having been destroyed inadvertently.

Thyrohyoideus.—Origin on the thyrohyal bone entad of the sternohyoid and the omohyoid and opposite the *hyoglossus*. Insertion on the posterior corner of the bony wing of the thyroid cartilage. A short stout triangular muscle, relatively well-developed.

Omo-hyoideus.—Origin on the lateral fourth of the posterior side of the hyoid bone. Insertion on the upper posterior angle of the scapula just craniad of the insertion of the *levator scapulae*. On the ental side near its origin it is joined to the sternohyoid by thin membrane.

Sterno-mastoideus.—Origin on the manubrium of the sternum slightly overlapped by the edge of the *pectoralis*. Insertion divided, one rather thin sheet inserting by broad aponeurosis on the occipital crest from a point midway between the median line and the auditory meatus to the middle of the anterior border of the mastoid bulla. The other division is a thickened bundle and inserts by a short tendon on the mastoid process slightly ectad and cephalad of that of the cleido-mastoid. The two divisions become fused about midway between origin and insertion.

Cleido-mastoideus.—Origin on the front of the distal end of the clavicle. Insertion fleshy or only slightly tendinous on the mastoid process. Its outer surface has relation with the *splenius*, the *sterno-mastoideus* and the *cleido-occipitalis*; its inner with the *digastricus*, *atlanto-acromialis* and *omo-hyoideus*.

Cleido-occipitalis.—Origin on the clavicle just laterad of the sternomastoid. Insertion by aponeurosis on the side of the occiput just laterad of and in the same plane with the *acromio-trapezius*. On its outer surface its relations are with the *cervico-auricularis*; on its inner with the *rhomboideus*, the *atlanto-acromialis*, and the *cleido-mastoideus*.

Omo-cleido-transversarius.—This includes two distinct muscles as in *Didelphis*, a dorsal part or "*atlanto-scapularis*" and a ventral part or "*atlanto-acromialis*." The dorsal part has its origin on the side of the body of the atlas immediately laterad of the origin of the ventral part. The insertion is on the front of the scapular spine in its vertebral fourth just ventrad of the insertion of the *rhomboideus minor*. It is a weaker more slender muscle than the ventral part and does not widen so at its insertion. The ventral part has origin on the side of the hypapophysial tubercle of the atlas. Its insertion is broad from the caudal side of the metacromion process of the scapula and for a short distance ventrad along the deltoid fascia over the acromio-deltoid muscle where it sends a few fibers ventrad.

MUSCLES OF THE THORAX.

Pectoralis minor.—Origin from the sides of the fourth and fifth sternebrae. Insertion on the lesser tuberosity of the humerus and the deltoid ridge by a broad slightly tendinous aponeurosis much wider than in *Didelphis*. Its outer surface has relation with the *pectoralis major*, its inner with the *rectus abdominis*, *scalenus*, etc.

Pectoralis major.—Origin from the median line of the entire sternum. Insertion on the distal part of the inner edge of the deltoid ridge. Its outer edge is curled under and near the insertion it has a slight aponeurotic connection with the *xiphihumeralis* as in *Didelphis*.

Supracostalis.—Origin by broad aponeurosis from the second, third, fourth and fifth sternebrae. Insertion fleshy on the side of the first rib. This muscle, which Coues called "*sterno-costalis*" and others designate *transversus costarum*, has practically the same relations as in *Didelphis*. Its broad aponeurosis for its entire length covers the *rectus abdominis*.

Subclavius.—Arises from the first rib and inserts on the outer side of the distal end of the clavicle, not extending to the scapular spine as in *Didelphis*. In this it agrees with *Phascologale*, and some other Australian forms (*Macropus* sp., *Phalanger* sp.).

Serratus magnus.—Origin posteriorly by six very distinct digitations on the sides of the ribs from the third to the eighth inclusive. Anteriorly there are no distinct digitations and the origin is from the joints of costal cartilages with the first, second and third ribs. Insertion on the pos-

terior border of the scapula where it is continuous with the *levator scapulae*.

Scalenus.—This has two fairly distinct divisions, at least posteriorly. The *scalenus medius*, or larger division, takes origin from the superior transverse processes of the third, fourth, and fifth cervical vertebrae over which it is somewhat narrow and thickened. Thence passing caudad it becomes thin and flattened and inserts by broad distinct slips on the middle of the third and fourth ribs, one slip on the third and two on the fourth, passing beneath and separating digitations of the *serratus magnus* in so doing. The *scalenus anticus* has its origin in coalescence with the *medius*, but about midway between the third cervical vertebra and the first rib it separates and inserts on the front of the first rib directly opposite the insertion of the *supracostalis*.

MUSCLES OF THE ABDOMEN.

Rectus abdominis.—Origin from the pubis and marsupial bone. Insertion on the sternal part of the first rib and on the adjoining side of the manubrium of the sternum. It is strong and well developed anteriorly, broader but thinner abdominally.

Obliquus externus abdominis.—Origin from the middle of the sides of the ribs from the fourth to the twelfth, interdigitating anteriorly with the *serratus*. Thence it follows the lumbar fascia to Poupart's ligament which is inserted at the inner anterior base of the marsupial bone. Ventrally it spreads over the abdomen covering the *rectus* and inserting by aponeurosis on the *linea alba*.

Obliquus internus abdominis.—Origin from Poupart's ligament and the ilium and thence dorsad to the lumbar fascia and craniad to the posterior borders of the 10th to 13th ribs. It is fleshy in its dorsal portion down to a line from the ventral third of the costal cartilage of the tenth rib diagonally across the side of the abdomen to the pubic region. Ventrad of this line the *transversalis* appears and no aponeurosis of the *obliquus* is demonstrable in the material available.

Transversus abdominis.—Origin from the inner sides of the 11th, 12th, and 13th ribs and from a coalescence with the diaphragm and thence along the lumbar fascia to the ilium. Ventrally it is inserted into the *linea alba*. Posteriorly it is broadly aponeurotic and anteriorly mostly fleshy.

Cremaster.—Arises in the male beneath the internal oblique near the crest of the ilium, and passes ventro-caudad across the *iliacus* and *psaos major* in the plane of the *transversalis* but separated from it by a slight space. It passes out of the body cavity through the ring in front of the

marsupial bone and proceeds to the mid-ventral line and the testes. It is a well developed strand of muscle about one mm. in width but simply parallels or covers but does not enclose the spermatic cord until near the base of the testes where it nearly or quite surrounds the cord before spreading over the base of the testes and becoming aponeurotic.

MUSCLES OF THE SACRUM.

Plates III, V.

Gluteus maximus.—Origin from the crest and the anterior part of the acetabular border of the ilium and from the articular processes of the sacral and anterior caudal vertebrae as well as from the tendinous fascia covering the *erector spinae* as far caudad as the origin of the *femorococcygeus*. Insertion somewhat divided, an anterior part on the cranio-lateral and a posterior one on the caudo-lateral surface of the distal end of the great trochanter of the femur. It is a broad fan-shaped muscle, relatively thicker than in *Didelphis*, being fleshy from the *erector spinae*. Its cranial border is thickened and somewhat curled around the anterior fifth of the ilium and distad lies in the groove between the *gluteus medius* and the *iliacus*, continuing ectad of the proximal end of the *vastus externus* to the cranial border of the *biceps femoris* to which it has a short semitendinous attachment distad and ectad of its insertion on the great trochanter. Its caudal border parallels the *femorococcygeus* but except in the dorsal fascia is distinct from it. No separate *tensor fasciae femoris* is evident and it is doubtless fused with the *gluteus* as in various other marsupials.

Gluteus medius.—Origin extensive from the dorsal fascia, the laminar tendons of the caudal muscles, the transverse processes of the sacral vertebrae, and the crest and ischial border of the ilium. Insertion on the caudo-lateral proximal part of the great trochanter of the femur ectad of that of the *gluteus minimus* which is distinct. It lies entad of the *gluteus maximus* and is almost equal to it in superficial extent. It is imperfectly divisible into two layers, the posterior one overlapping the anterior. A similar bilaminar arrangement is reported in *Phascolarctos*, *Thylacinus*, and *Dasyurus*, but not in *Didelphis*¹ and *Cuscus* (see Leche, Bronn's Thierr., p. 853). It is likewise in *Notoryctes* (Thompson and Hillier, 1905, p. 310). On its ental side it is somewhat connected with the *pyriformis* not only near the insertion but also elsewhere except along the anterior tendinous border of the *pyriformis*. Anteriorly it is curled under from the crest of the ilium to the femur but in the distal half of this extent its edge is free and merely overlies the *gluteus minimus*.

¹ Sidebotham (1885, p. 13) reports three layers in *Chironectes*.

Gluteus minimus.—Origin from the dorsal surface of the ilium from a point on the acetabular border about midway between the crest and the acetabulum to a point opposite the middle of the acetabulum. Insertion broadly covering the antero-internal surface of the great trochanter of the femur. It is relatively large and completely distinct not only anteriorly from the *gluteus maximus* and *gluteus minimus* but posteriorly from the *gemelli*. In the didelphids and various other marsupials it is partially joined to the *maximus*.

Pyriiformis.—Origin fleshy from the dorso-lateral surface and the caudal edge of the crest of the ilium and from the tips of the transverse processes of the two sacral and the first caudal vertebrae.

Insertion on the apex of the great trochanter of the femur just caudad of the insertion of the *gluteus minimus* and entad of the *gluteus medius*. A very large muscle lying entad of the *gluteus medius* and ecto-caudad of the *gluteus minimus*. Its anterior border which is full and rounded runs direct from the trochanter to the iliac crest and at least in its distal half is tendinous. Fleshy fibers pass dorsad from this broad tendinous edge to the sacral border. Its outer surface, at least caudally, is somewhat connected with the *gluteus medius*.

Quadratus femoris.—Origin on the inner side and slightly ventrad of the tuberosity of the ischium and on the inner edge of the ischium just craniad of the tuberosity. The fibers curve over the lesser sciatic notch to pass outward to their insertion on the middle of the ventral edge of the great trochanter of the femur and on the intertrochanteric ridge. It is relatively large and well developed and fairly distinguishable from the *gemelli* anteriorly.

Gemelli.—No distinction is evident between a *gemellus superior* and *inferior*. The origin is on the body of the ischium dorsad of the origin of the *ischiococcygeus*. The insertion is with the tendon of the *obturator internus* on the caudal edge of the great trochanter of the femur and in the digital fossa proximad of the insertion of the *quadratus femoris*. Its cranial border is distinct from the caudal border of the *gluteus minimus*.

Iliacus.—Origin from the ventro-lateral or iliac surface and the acetabular border of the cranial two-thirds of the ilium from the crest caudad. In its caudal portions its insertion is opposite that of the *iliococcygeus* on one side and the *gluteus minimus* on the other with a part of the *psoas magnus* lying loosely between. Its general relations are closely similar to those in *Didelphis*.

Obturator internus.—Origin from all of the caudal border of the ramus of the ischium which is ventrad of the *quadratus femoris*, from the inner side of the pubic symphysis, and from the caudal border of the obturator foramen. Insertion by a flat tendon in the digital fossa of the femur.

Obturator externus.—Origin from the side of the ascending ramus of the ischium from the lower edge of the tuberosity to the symphysis and thence on the side of the pubic ramus to a point nearly opposite the anterior base of the marsupial bone. Insertion by a thick tendon in the digital fossa and on the intertrochanteric ridge of the femur. It is a broad thickened muscle lying at its origin entad of the adductors. Dorsally it meets the edge of the *quadratus femoris*.

The *obturator externus* is said by Leche (p. 858) to be "nichts bemerkenswerthes" in the Marsupialia, but Macalister (1870, p. 167) states that it is "large and normal in *Phascolomys* and *Sarcophilus*, as well as in *Macropus giganteus*, the Wallaby, Phalanger, and Opossum." Thompson and Hillier (1905, p. 312) found it well developed in *Notoryctes*. The so-called differentiation of the *obturator*, which Leche names *intermedius* and reports only for *Didelphis* and *Philander* is doubtless not peculiar to these forms. It was not specially sought in *Cenolestes* and may have been overlooked and included as an integral part of the *externus*. Whether it be called *externus* or *intermedius*, it is apparent that it is not peculiar in *Cenolestes*.

MUSCLES OF THE TAIL.

All the usual tail muscles of the pubic cavity are present and distinct but their muscular parts do not extend far beyond the base of the tail. On removal of the skin and the underlying fascia, the tail appears almost entirely encased in shining tendons from its base to its tip. On the dorsal aspect the continuations of the back muscles proceed somewhat farther caudad but these also soon become entirely tendinous. After all the tendons have been removed, however, a series of thin muscles connecting the vertebrae are exposed. These are short muscles, those next the median line extending from the chevron bone of one vertebra to that of the one succeeding; those laterad extend from the caudal transverse process of one vertebra to the cephalic articular process of the second succeeding vertebra, each muscle thus spanning a vertebra.

Dorsally the tendons from the dorsal extensors cover the angles of the tail, passing over the articular processes of the vertebrae, but in the median line the short intervertebral tail muscles are exposed.

Abductor caudae externus.—Origin on the medial side of the iliac crest and by tendinous connection with the lumbo dorsal fascia. Insertion on the dorsal surfaces and tips of transverse processes of the caudal vertebrae. This is a spindle-shaped muscle filling the great part of the space between the transverse and the articular processes of the anterior caudal vertebrae and passing out on the side of the tail where it narrows and soon becomes wholly tendinous.

Extensor caudae lateralis.—Origin by slips from the articular processes of the sacral and caudal vertebrae. Insertion on the sides and transverse processes of the caudal vertebrae. Lies between the *abductor caudae externus* and the *multifidae* which are continued to the base of the tail as an *extensor caudae medialis*.

Pubococcygeus.—Origin from the whole inner caudal border of the pubic ramus. It forms a thin sheet over the obturator foramen and continues fleshy to the front of the fifth caudal vertebra whence it finds attachment in the aponeurosis on the surface of the tail. In some cases it is slightly tendinous on its inner side and attached to the chevron bone of the fifth caudal. Laterally it has slight attachment to the colon.

Sacrococcygeus.—Origin from the ventral surfaces of the transverse processes of the sacral and the first four caudal vertebrae. Thence by numerous slender tendons it proceeds along the side of the tail to the distal caudal vertebrae. These tendons lie slightly laterad of those of the *infracoccygeus* and fill the space between the cephalic articular and transverse processes of the caudal vertebrae. Its muscular heads are thick and fleshy and lie between the caudal part of the *infracoccygeus* and the *pubococcygeus* and ventrad of the *ischiococcygeus*.

Ischiococcygeus.—Origin from the inner side of the ilium from a point opposite and slightly craniad of the caudal border of the *gluteus minimus* caudad about halfway to the tuberosity of the ischium. Insertion on the tips of the transverse processes of the first four caudal vertebrae.

Iliococcygeus.—Origin from the ischial border of the ilium from a point just caudad of the sacrum (posterior inferior spine of ilium) and thence caudad along the whole inner surface of the ilium (great sacro-sciatic notch) nearly to the anterior border of the obturator foramen. A few fibers rise from the tendons of the *psaos minor* and at this point are closely connected with those of the *pubococcygeus*. Insertion by four distinct tendons which converge to the middle of the ventral surface of the tail. The outer tendon runs just laterad of the mid-ventral line of the tail to the chevron bone of the eighth caudal vertebra. The next one runs similarly to the seventh caudal. The inner pair of tendons run first mediad, then entad, and finally laterad of the outer pair. Of this inner pair the outer one inserts on the chevron bone of the ninth and the other on the tenth caudal. Near their attachments each of the tendons gives off a slight muscular fasciculus.

Infracoccygeus.—Origin from the medio-ventral surfaces of the bodies of the fifth and sixth lumbar and the two sacral vertebrae. Insertion by long slender tendons running on the ventral side of the tail next to the median line. The innermost tendon runs to the middle of

the front of the first caudal. The next tendon laterad runs to the eleventh caudal, the next to the twelfth and so they continue to the twentieth caudal. The muscle is broadly spindle shaped and occupies the sacral basin, its two halves forming an elongate ellipse between the two halves of the *psaos magnus* anteriorly and of the *iliococcygeus* posteriorly. It consists of superposed laminae each becoming tendinous before issuing from the pubis.

MUSCLES OF THE ANTERIOR LIMB.

Plate V.

Spino-deltoides.—Origin from the middle third of the scapular spine. Insertion on the distal end and outer edge of the deltoid ridge of the humerus. At its insertion it passes beneath and nearly at right angles to the *acromio-deltoides*.

Acromio-deltoides.—Origin on the acromion and the proximal half of the clavicle. Insertion on the inner end of the flattened surface of the deltoid ridge opposite the insertion of the *pectoralis major*. It covers the rather broad surface of the deltoid ridge and is a thicker muscle than in *Didelphis* and more distinct from the *spino-deltoides* and the *pectoralis*. It is fan-shaped, the fibers from the clavicle and those from the acromion converging toward a median raphe and coming to a point ventrally.

Subscapularis.—Origin from practically the whole of the expanded ental surface of the scapula. Insertion by a short broad tendon on the lesser tuberosity of the humerus. It is slightly thicker dorsally than ventrally but in general is thin and flat.

Supraspinatus.—Origin from the surface of the supraspinous fossa. It passes entad of the clavicle between the coracoid and the acromion and inserts on the greater tuberosity of the humerus. It is about twice as large as the *infraspinatus* and relatively larger than in *Didelphis*, agreeing more nearly with *Phalanger*, *Perameles*, *Phascologale*, etc.

Infraspinatus.—Origin from the infraspinous fossa, including the ental surface of the metacromion and acromion, the axillary border of the scapula, and from a tendinous raphe between it and the proximal part of the *teres*. Insertion partly fleshy and partly tendinous on the greater tuberosity of the humerus. A weak aponeurosis from the infraspinous fossa near the base of the acromion may represent the *teres minor*, but it is very indefinite. A *teres minor* is reported for *Chironectes* (Sidebotham) but not for *Didelphis*.

Epitrochlearis.—Origin from the lower border of the *latissimus* slightly (2 mm.) before it passes beneath the *biceps*. Insertion on the olecranon process of the ulna.

Teres major.—Origin from the posterior ventral angle of the scapula in close relation with the *subscapularis*. Insertion on the bicipital groove slightly proximad of and in conjunction with that of the *latissimus*.

Coracobrachialis brevis.—Origin on the coracoid process entad of the *biceps*. Insertion on the humerus just proximad of the insertion of the *teres*. It is very slightly developed, almost rudimentary, and in its fleshy part consists of a very thin layer of fibers instead of a thick belly as in *Didelphis*. In one specimen it appeared to insert in the tissue of the medial head of the *triceps* rather than on the humerus.

Biceps brachii.—Origin by a single tendon from the coracoid process of the scapula and adjacent fascia. Soon after becoming fleshy it is readily divisible into two distinct muscles which run in close apposition to pass into the forearm between the *pronator teres* and the *extensor carpi radialis brevis*. The outer muscle inserts by a narrow flat tendon on the tubercle of the radius. The inner muscle is partially attached to the *brachialis* near its insertion and then its tendons divide and pass on either side of the tendon of the *brachialis* to insertions on the side of the ulna distad of the lesser sigmoid cavity. In some of the Macropodidae the *biceps* is completely divided but in some other diprotodonts and in polyprotodonts generally its relations are reported essentially as above.

Brachialis anticus.—Origin from the lesser tuberosity and the border of the head of the humerus passing around the middle of the front of the humerus and thence distad. At about the middle of the front of the humerus it becomes attached to the bone; between this and its origin it is free or nearly so. Thence it continues along the humerus to the elbow joint where it twists further and has final insertion just distad of the lesser sigmoid cavity on the inner surface of the ulna by a flattened tendon which passes between the divided tendons of the *biceps*.

Triceps brachii.—The long head has its origin as usual on the axillary border of the scapula just dorsad of the glenoid cavity. Its insertion by common tendon with the lateral head is on the outer angle of the olecranon process of the ulna.

The lateral head arises beneath the posterior border of the head of the humerus and thence is free from the humerus to its union with the long head. Its posterior aspect is hollowed to receive the long head for most of its length and the two are fairly distinct. The median head arises from the posterior aspect of the humerus for practically its entire length, reaching dorsally within 2 mm. of the lesser tuberosity and thence to the olecranon fossa. Its insertion is on the inner angle of the olecranon process of the ulna quite distinct from that of the other two heads.

The *triceps* of *Cænolestes* is characterized by the distinctness of the

median head which in *Didelphis* and many other marsupials is united with the lateral head near its origin.

Anconeus.—Origin from the inner condyle of the humerus. Insertion on the inner angle of the olecranon process entad of the *epitrochlearis*. A small muscle connected on its lower side by a tendinous raphe with the *extensor carpi ulnaris* and on its upper side slightly connected with the median head of the *triceps*.

Pronator teres.—Has origin as a thick fleshy bundle from the posterior side of the inner condyle of the humerus and the epicondylar ridge. Insertion by broad aponeurosis on the middle third of the front of the radius.

Flexor carpi radialis.—Origin from the inner humeral condyle and the septa connecting with the *pronator teres* between which and the *palmaris longus* it lies. Its long slender tendon passes over the extremity of the radius and thence over the scaphoid to insertion at the bases of the second and third metacarpals.¹

Palmaris longus.—Origin from the septa between it and the *flexor carpi ulnaris*, the *flexor carpi radialis*, and the digital flexors and by a few fibers from the inner condyle of the humerus. It covers the digital flexors almost completely and lies between the *flexor carpi ulnaris* and the *flexor carpi radialis*.

In the single specimen dissected it was divided into two parts. The superficial one (*palmaris accessorius*?) sends its tendon beside that of the *flexor carpi ulnaris* to a slight attachment on the inner side of the fascia on the pisiform bone and thence superficial to or partly attached to the flat transverse ligament of the wrist which extends across the wrist from the pisiform to the base of the first metacarpal. Above this transverse ligament it extends into the palmar fascia. The deeper division sends its tendon bound with its fellow to the transverse ligament where the two separate and the second dips beneath the ligament to reappear in the fascia and send aponeurotic branches to the extremities of the second, third, and fourth digits. The fleshy parts of the two divisions are distinct for about one-half their length.

Flexor carpi ulnaris.—Origin from the inner condyle of the humerus, from the septa between it and the *palmaris longus*, the *extensor digitorum communis*, and the *anconeus*, and from the outer angle of the olecranon opposite the insertion of the *epitrochlearis*. Insertion on the end of the pisiform bone. A broad fan-shaped muscle with two sets of fibers converging toward its tendon which extends a short distance into its fleshy

¹ This is also the case in *Didelphis*, although Coues states that the insertion is on the first metacarpal; Cunningham (1882, p. 17) found a similar disposition in *Phalanger maculata*.

part. The fleshy divisions end abruptly and equally about one-third of the distance from the olecranon to the carpus. The posterior division, therefore is not longer than the anterior as it is in the opossum and the tendon is shorter.

Flexor digitorum sublimis.—Origin by three heads in the septa of the ulnar head of the *profundus*. These give rise to three distinct slips which send their tendons loosely bound together until they reach the palm where they pass beneath the annular ligament and insert ectad of the large *profundus* tendons at the bases of the first phalanx of the second, third, and fourth digits, respectively. Their tendons are slender and practically of the same size throughout. From their point of origin a short thick muscle with fibers running diagonally opposite those of the ulnar head of the *profundus* extends to an origin from the caudad surface of the inner humeral condyle and from the proximal edge of the glenoid cavity of the ulna. Whether it may be one head of the *sublimis* is doubtful. It is very distinct.

Flexor digitorum profundus.—Origin by four heads each of which sends a distinct tendon to the palm where all unite in a thick flat tendinous mass which occupies most of the palmar space. From this three strong tendons radiate to the middle digits. Two others going to the pollex and the fifth finger are much smaller and arise from the ectal surface of the common tendon. The heads are as follows:

1. Ulnar head. Origin on the flattened posterior surface of the proximal two-thirds of the ulna from the olecranon at least to a point opposite the neck of the radius. Its tendon is very large and strong, rounded in its proximal part, slightly tapering toward its middle, and flattening as it joins the outer side of the palmar tendinous mass.

This is the largest fasciculus of the forearm, lying beneath the *flexor ulnaris* and giving shape to the limb.

2. The second head lies superficially next to the *flexor carpi radialis* and connected with it in its proximal part by intermuscular septa. It is similarly connected with the fourth head for a distance of nearly half its fleshy extent. Its origin is from the inner condyle of the humerus entad of that of the *palmaris longus*. Its tendon joins the common tendon ectad of that of the third head, the two being approximately of equal size and lying midway between the tendons of the first and fourth heads which are much larger.

3. The third head is very distinct and wholly free but lies deeper than the second and is not exposed until that is reflected. It lies between the *flexor ulnaris* and the second head of the *profundus* together with the conjoined part of the *flexor radialis*. Its origin is from the inner side of the lip of the lesser sigmoid cavity of the ulna. Its small round tendon

runs direct to the common palmar tendon which it joins next to that of the ulnar head and entad of that of the second head.

4. The fourth head, which is almost as large as the first or ulnar head, has origin from a distinct slip from the front of the inner condyle of the humerus and from a broad fleshy expansion arising from the inner side of the proximal fourth of the radius. The two fleshy parts join to form a single tendon which runs along the radius to join the common tendon in the palm.

While obviously very different from that of *Didelphis*, in which there is no division above the palm, this muscle is so variable that detailed comparisons have not been attempted. In most marsupials it sends tendons only to the four ulnar digits.

Abductor pollicis longus (*Extensor ossis metacarpi pollicis*).—Lies in the deep groove in the proximal part of the ulna and takes origin from the adjacent sides of the ulna and the radius, the fibers converging from the two sides to the central tendon which extends some distance into the fleshy part. Its origin (male specimen) is thus some 5 mm. in length. Its rather broad flat tendon passes beneath the ulnar and lateral digital extensors crossing the forearm and becoming superficial between the *extensor digitorum communis* and the *extensor carpi brevis*. It then curves around the radius and inserts on the lateral surface of the first metacarpal.

Pronator quadratus.—This muscle, which is present in didelphids and so far as known in all other marsupials except *Notoryctes*, is not found in *Cænolestes*, or at most is represented by no more than an indefinite non-muscular fascia. It is reported as poorly developed in *Thylacinus*, *Sarcophilus*, *Phalanger* and *Phascogale*. It is lacking in monotremes, insectivores, bats, and certain edentates (*Dasypodidae*, *Manidae*).

Brachioradialis (*Supinator longus*).—Origin from the outer epicondylar ridge of the humerus proximad of the *extensor carpi radialis longus*. Its tendon passes under that of the *abductor pollicis longus* and over a groove in the extremity of the radius to insertion in the external lateral ligament which overlies the extremity of the radius.

Extensor carpi radialis longus.—Origin from the outer epicondylar ridge of the humerus distad of the *brachioradialis*. Insertion on the base of the second phalanx of the second digit.

Extensor carpi radialis brevis.—Origin from the outer epicondylar ridge and septa of the *extensor digitorum communis*, between which and the *extensor carpi radialis longus* it lies. Insertion on the outer lateral surface of the proximal third of the first phalanx of the third digit. It is somewhat larger than the *longus* and separate from it as in *Thylacinus* and *Phalanger*. In most other marsupials, the long and short radial

extensors are reported to be fused at least proximally, but Parsons (1896, p. 697) expresses the belief that they are normally separate in the kangaroos.

Extensor digitorum communis.—Origin from the distal part of the outer epicondylar ridge, the outer condyle and from the septa of the muscles lying on either side, namely the *extensor carpi radialis brevis* and the *extensor digitorum lateralis*. The fleshy part divides into two slips, the more superficial of which sends tendons to the second and third digits and the deeper one to the fourth and fifth. The tendons are separate but closely parallel until they pass under the transverse ligament and separate to insertions on the terminal phalanges of the second to the fifth digits. The tendons become somewhat flattened or aponeurotic distally.

Extensor digitorum lateralis.—Origin on the front of the outer condyle of the humerus next to the *extensor communis* but quite distinct from it. Connected by septa with the *extensor carpi ulnaris*. The head of the radius lies immediately entad of the line of separation between the two muscles. In the distal two thirds of its fleshy extent it is divided into three distinct slips from which separate parallel tendons run one to the fourth and two to the fifth digit. On the fourth digit the insertion is at the base of the ungual phalanx. On the fifth the insertion in one case is at the base of the ungual phalanx and in the other in the fascia covering the second and third phalanges. A double tendon from this muscle to the fifth digit has been noted in *Sarcophilus* (Macalister, 1870, p. 164), but not in other marsupials.

Extensor carpi ulnaris.—Origin from the front of the outer condyle of the humerus and the adjacent side of the ulna; also connected by septa with the *extensor digitorum lateralis*. Its tendon passes over a groove on the antero-internal surface of the extremity of the radius and inserts at the base of the fifth metacarpal. According to Macalister (l. c.) and Wilson (1894, p. 451), this muscle has no ulnar origin in *Sarcophilus*, *Didelphis*, *Trichosurus*, and *Dasyurus*, but like *Cænolestes*, has one in *Macropus*, *Phascolomys*, *Phascolarctos*, and *Perameles*.

Supinator (brevis).—Origin tendinous on the sesamoid laterad of the head of the radius. Insertion on the arched anterior surface of the proximal third of the radius. A broad flat muscle. It occupies the upper two-thirds of the radius in *Phascolomys*, two-fifths in *Phascolarctos*, one-third in *Macropus* and *Sarcophilus* and only one-fourth in *Didelphis* and *Dasyurus* (Macalister l. c.). The presence of a sesamoid in its tendon of origin appears to be unique among marsupials. It is reported in certain edentates and bats and an origin from the orbicular ligament of the radius was found in *Dasyurus* by MacCormick (fide Wilson, l. c., p. 48).

Extensor indicis.—Origin from the antero-external edge of the ulna opposite the greater sigmoid cavity. Its tendon divides into two about midway of its length. These pass under the digital extensors and across the carpus to insertions one at the base of the ungual phalanx of the first digit and one at the same point on the second digit.

MUSCLES OF THE FOREFOOT.

The delicate intrinsic muscles of the forefoot have been worked out as carefully as possible, but in all the specimens examined some of the metacarpal bones were found to be broken, making it very difficult to ascertain exact relations. In general the muscles of the hand differ from those of *Didelphis* more than they do from some of the Australian forms. The adductors are much less highly developed than in *Didelphis* and this in connection with the absence of a *pronator quadratus* indicates a relatively limited power of action of hand and fingers.

Lumbricals are present but not greatly developed.

Adductors.—Three adductors are readily distinguishable, the *indicus*, *annularis*, and *minimus*. They arise in the palmar fascia at the base of the metacarpals and radiate to their insertions as practically separate muscles without any development of a central raphe. The *indicus* arises over the bases of the second and third metacarpals and inserts on the ulnar side of the distal extremity of the second metacarpal. It is separable with some difficulty from the ulnar head of the *flexor brevis indicis*. The *annularis* and *minimus* arise over the bases of the third and fourth metacarpals and insert on the radial sides of the distal ends of the fourth and fifth metacarpals respectively. They are separated from the flexors by a large palmar nerve. An *adductor pollicis* could not be distinguished. In general the adductors seem to be very similar to those of *Thylacinus* as figured by Cunningham.

Flexors.—Short flexors are well developed on all the fingers. The *flexor brevis pollicis* could not be divided into two parts. It arises on the ulnar side of the base of the first metacarpal and inserts medially with a sesamoid at the first joint of the pollex. It is a relatively thick muscle and possibly may include a fused adductor. A rather definite aponeurotic strand running from the carpus to the ulnar side of the first joint of the pollex may represent a second head of this flexor. The remaining flexors are divided except proximally and lie in the usual position on the metacarpals with their tendons of insertion embracing the metacarpo-phalangeal joints of each of the digits.

Abductors and interossei.—Two marginal muscles regarded respectively as *abductor pollicis* and *abductor digiti minimi* are present.

There is also an abductor of the index digit which is separately described owing to its more palmar position than the other interossei.

The *abductor pollicis* is a short thick muscle with origin from the trapezium, the transverse ligament and the base of the first metacarpal. Its insertion is on the radial side of the distal extremity of the first metacarpal.

The *abductor digiti minimi* also is a relatively large muscle. It originates on the pisiform bone opposite the insertion of the *flexor carpi ulnaris* and inserts by a flattened tendon running under the annular ligament to the outer side of the metacarpo-phalangeal joint of the fifth finger.

The *abductor digiti indicis* arises on the trapezium and the base of the first metacarpal and inserts near the distal end and on the radial side of the second phalanx in close apposition with the short flexor. It is in nearly the same plane as the flexors and is visible alongside the *abductor pollicis* before removal of the large tendons. It is also visible from the dorsal aspect of the hand.

Excluding the above described *abductor indicis*, there are three dorsal interosseus muscles. These lie over the spaces between the four outer metacarpals, are almost wholly dorsal in position, and resemble those described and figured for *Phalanger* (*Cuscus*) by Cunningham (loc. cit., p. 23, pl. II, fig. 2). The one between the index and the medius has origin principally on the ulnar base and side of the second metacarpal and insertion with the tendon of the short flexor on the radial side of the third metacarpal. That between the medius and the annularis arises from the bases and the sides of the third and fourth metacarpals and inserts on the tendon of the short flexor on the ulnar side of the third metacarpal. That between the annularis and minimus arises from the bases and sides of the fourth and fifth metacarpals and inserts on the tendon of the short flexor of the ulnar side of the fourth metacarpal, also sending a short tendinous slip to the radial side of the fifth metacarpal.

MUSCLES OF THE HIND LIMB.

Plates V-VI.

Adductor longus.—Origin from the side of the ramus of the ischium from the symphysis pubis to the insertion of the *semimembranosus*. Its origin is ectad of the *adductor magnus* and entad of the *gracilis*. Its caudal border parallels the *semimembranosus*. Insertion on the inner condyle of the femur and the inner side of the head of the tibia.

Adductor magnus.—Origin from the side of the pubic symphysis and thence dorso-cranial along the edge of the pubis to the middle of the

base of the marsupial bone and dorsad along the ramus of the ischium to the *semimembranosus*. Insertion on most of the length of the caudal surface of the femur from the insertion of the *adductor brevis* near the lesser trochanter to the insertion of the *longus* at the base of the inner condyle. This muscle lies between the *brevis* and the *longus* and its exposed surface lies next to the *pectineus*. At its origin it lies entad of the *longus* which parallels it distad.

Adductor brevis.—Origin from the side of the dorsal half of the ascending ramus of the ischium to the side and front of the tuberosity of the ischium and for a slight distance craniad. Insertion on the distal caudal edge of the great trochanter of the femur caudo-entad of the insertion of the *gluteus maximus* and thence transversely of the femur to a point slightly distad of the lesser trochanter. The outer surface of this short broad muscle is wholly covered by the *gracilis*, the *semimembranosus*, and the other adductors.

Caudofemoralis.—Origin by a broad tendinous aponeurosis from the fascia covering the *multifidae* and somewhat attached to the articular processes of the third caudal vertebra. The insertion is double, one slender slip inserting by a short tendon on the inner edge of the inner condyle of the femur between the *adductor longus* and the *gastrocnemius*. The other division of the muscle passes on the opposite side of the *adductor longus* and inserts by several fasciculi on the caudal surface of the distal fourth of the shaft of the femur. Near its insertion it is slightly connected with the adductor. This muscle, which is not present in the didelphids, is well developed, but shows some variation even in the small number of specimens dissected. At its origin it lies immediately entad of the *femorococcygeus* and slightly distad is separated from it by the great sciatic nerve. It passes transversely to the bases of the adductors, but is free from the femur to its insertion. Its ental surface near its origin has relation with the caudal muscles, the *quadratus femoris*, and the *obturator internus* as it passes distad just craniad of the *biceps femoris*. Its presence is reported in *Notoryctes*, *Phalanger*, *Thylacinus*, *Myrmecobius*, and *Dasyurus*.

Pectineus.—Origin from the side of the pubis at the anterior base of the marsupial bone. Insertion on the inner surface of the femur slightly proximad of the middle. A short thick muscle lying ectad of the insertion of the *psoas major*.

Femoro-coccygeus.—Origin aponeurotic from tendinous fascia between the cephalic and caudal articular processes of the third caudal vertebra. Insertion by several fasciculi on the distal third of the caudolateral surface of the femur. It is separate from the *gluteus maximus* except near the insertion where the fascia join. It has a slight twist

in passing between the *biceps* and the *gluteus* and lies "edgewise" for a short distance, its distal part being covered by the *biceps*. It is separate from the *gluteus* in *Sarcophilus* (Macalister, 1870, p. 166) and *Notoryctes* (Thompson and Hillier, 1905, p. 310), but in other marsupials is in the same plane and more or less fused with it.

Vastus internus.—Origin from the proximal three-fourths of the anterior and dorsal surface of the femur, extending from the anterior edge of the great trochanter to the lesser trochanter and thence distad. Insertion with the *vastus externus* on the inner side of the patella and thence by aponeurosis to the head of the tibia. A very large muscle greatly exceeding the *vastus externus*.

Subcrureus.—Origin on the dorsal surface of the femur just distad of the *vastus internus*. Insertion in the tendinous fascia surrounding the patella. A short flat muscle.

Vastus externus.—Origin from the anterior border of the great trochanter of the femur. Insertion conjoined with the *rectus femoris* on the patella. About equal in size and form to the *rectus*, with which it forms the anterior border of the thigh. Decidedly smaller than the *vastus internus* to which it is closely appressed but from which it is distinct except for a short distance near its origin.

Biceps femoris.—Origin by a short tendon from the lateral surface of the tuberosity of the ischium craniad of the inner division of the *semitendinosus* and entad of the outer division or *crurococcygeus*. Insertion by broad aponeurosis from the patella across the proximal third of the leg to the outer edge of the tibia. This aponeurosis covers a large part of the outer side of the leg. A large triangular muscle of great width but moderate thickness. Its anterior border meets the *vastus externus* and covers the insertion of the *femorococcygeus*.

Rectus femoris.—Origin from the dorso-lateral ridge of the ilium immediately craniad of the acetabulum. Insertion with the vasti on the patella. It passes between the *gluteus minimus* and the *ileopsoas* and forms the inner anterior boundary of the thigh.

Semitendinosus.—Origin by two heads, an outer one ("*crurococcygeus*") from the fascia over the fourth caudal vertebra and an inner one from the tuberosity of the ischium just caudad of the *biceps*. Midway between the tuberosity and the flexed femur these two slips unite and then divide into three parts, two passing inside the *gastrocnemius* and one outside. The inner slips insert by aponeurosis, one with a twist on the front of the tibia one-third of the distance from the tibial head and the other (entad) on the same surface of the tibia halfway between its fellow and the head of the tibia. The outer slip inserts on the outer surface of the tibia and in the fascia of the *biceps femoris*. The relations

of this muscle are essentially as in *Didelphis* and different from those of most other marsupials, in which the so-called *crurococcygeus* is not so well developed, although tendinous intersections have been found in *Thylacinus*, *Dasyurus*, and *Trichosurus* (fide Thompson and Hillier, 1905, p. 316) and in *Chæropus* (Parsons, 1903, p. 73).

Gracilis.—Origin from the pubic symphysis and the ascending ramus of the ischium nearly half way to the tuberosity. Anteriorly a few fibers reach to the base of the marsupial bone. Insertion aponeurotic on the inner surface of the distal half of the proximal third of the tibia.

Semimembranosus.—Origin fleshy from the outer side of the ascending ramus of the ischium, occupying the middle two-thirds of the space between the symphysis and the tuberosity. Insertion by a broad tendon on the inner side of the head of the tibia entad of the internal lateral ligament of the knee joint which is strongly developed. A large thick muscle elliptical in cross section. No *presemimembranosus* was distinguished, but it may be represented in the somewhat anomalous muscle here called *caudofemoralis* although that has no ischial origin.

Sartorius.—Origin on the dorsal border of Poupart's ligament. Insertion in the fascia extending from the knee joint and the patella to the proximal third of the front edge of the tibia. It is a thin band consisting of one layer of muscle fibers only and can scarcely be functional. This muscular part is clearly differentiated from the fascia on either side of it which cover a great part of the inside of the leg immediately beneath (entad) the mass of fat. Between the *sartorius* and the next layer of muscle is the crural nerve and its branches, of which the saphenus innervates the *sartorius* and runs down the side of the leg (Pl. VI, fig. 2). It is practically imbedded in the *sartorius* at this point and is visible through the thin muscle layer. The insertion of the muscle is ectad of the *gracilis*.

This much-reduced, almost rudimentary muscle, is easily overlooked and it was at first thought that a *sartorius* was entirely absent or fused with the *gluteus maximus*. It was possible also that it might be a second *gracilis*. It has been demonstrated in two specimens, however, and its relations with the saphenus nerve seem to justify the conclusion that it is in reality the *sartorius*. Its origin from Poupart's ligament is paralleled in certain edentates, as sloths and armadillos. A partial origin from Poupart's ligament has been noted in *Sarcophilus* by Macalister (1870, p. 170). In the Insectivora, although usually absent, the *sartorius* is found in *Gymnura* partly attached to the *gracilis*.

Gastrocnemius.—The medial head arises by a short thick tendon from the inner transverse surface of the inner condyle of the femur and the intercondyloid notch just distad of the *caudofemoralis* and the *adductor*

longus. It lies against the lateral head with various nerves intervening and is joined with it in the distal fourth of its fleshy extent. It thus agrees with most marsupials and differs from *Didelphis* in which it is wholly free from the lateral head. The lateral head has two divisions, one of which probably represents the *soleus*. These are:

1. The principal part arises from the extremity of the fibular sesamoid opposite the end of the collateral tibial ligament and joins the second part about midway of its fleshy extent. Except for a small triangular area along its outer proximal edge where the second part is visible beneath the fascia, it overlies the second part.

2. The second part arises from the caudal surface of the fabella and from the tendinous edge of the *plantaris* for about two-fifths of the fleshy extent of the *plantaris*. It joins the outer part about midway of its fleshy extent. This second and more deep-seated part of the muscle may be the *soleus* which is not otherwise evident. The conjoined parts form a rounded tendon which runs to an insertion on the calcaneum. As a whole the *gastrocnemius* forms a deep belly on the leg curving rather abruptly inward to its tendon and occupying in its fleshy part only forty per cent of the length of the leg, the remainder being tendinous. Its outer or anterior edge is continuous with the fascia lata which extends to the patellar region and down the leg entad of and intimately associated with the fascia of the *biceps femoris*. In one specimen a slight cleft in this outer edge was observed separating a thin slip which may represent a third or patellar division of the *gastrocnemius*.

Plantaris.—Origin from the side of the fabella, mediad of the second division of the lateral head of the *gastrocnemius* and more or less united with it. It continues joined with the *gastrocnemius* for about two-fifths of its fleshy extent and then becomes free as a slender bundle which proceeds distad covered by the *gastrocnemius*. Its slender tendon, which at first is entad of that of the *gastrocnemius*, curves around it and passes over the calcaneum ectad of the *gastrocnemius* tendon and widens to merge into the plantar fascia where its prolongations could not be traced.

Flexor digitorum fibularis.—This has two divisions scarcely separable in their fleshy part. The smaller one, which is on the tibial side, sends a small tendon beneath the larger one but separate from it until just before it passes the os calcis when the two merge. The fleshy part of the muscle is relatively large, lying next below the *gastrocnemius* and taking origin from the greater part of the inner and posterior aspects of the fibula from the inner side of the outer tuberosity distad not only in the grooved inner expanded part but also on the proximal part of the rounded posterior aspect. Proximally it is connected by septa with the *tibialis anticus* and distally some of its fibers arise from the interosseous mem-

brane. Its large tendon passes inside the calcaneum where it becomes broad and heavy and soon divides into four strong tendons bearing the lumbricals and running to the ends of the outer toes. Just before the common tendon divides it receives on its ental surface the two flattened tendons of the *flexor digitorum tibialis* between which passes the *flexor accessorius*.

Flexor accessorius.—Arises on the tuberosity of the distal outer extremity of the os calcis, spreads over the ventral surface of the distal part of the calcaneum, passes beneath the plantar ligament and then becoming tendinous, continues diagonally entad of the common tendon of the large *flexor fibularis*. Thence it perforates the tendon of the *flexor tibialis*, or at least is somewhat attached to it, and proceeds to insertion on the distal phalanx of the hallux. Just before passing through the annular ligament at the first joint of the hallux its tendon unites with that of the *flexor tibialis*.

This muscle is present in *Didelphis* and has been described by Coues (1872, p. 134) under the name *flexor brevis pollicis obliquus*. In *Didelphis*, however, it passes ectad of the common tendon of the *flexor fibularis* and its tendon joins the tendon of the *flexor tibialis* before it reaches the hallux. An *accessorius* is recorded also for *Chironectes* (Sidebotham, 1885, p. 16).

The *accessorius* is stated by Leche (Bronn's Thierr., p. 904) to be generally lacking in marsupials and perhaps the above described muscle should be regarded as a part of the *flexor digitorum communis brevis* which is otherwise rather weak and slightly represented (see postea, p. 57). Whatever its name, it is evident that it is more independent in *Cænolestes* than in other marsupials or most other mammals, especially in the distinctness of its tendon for a considerable distance on the hallux. A somewhat similar *accessorius* found in an insectivore, *Gymnura*, is regarded as abnormal by Parsons (1898, p. 324). It is of regular occurrence in edentates.

Flexor digitorum tibialis.—Origin from the greater part of the proximal half of the interosseous membrane, from the *popliteus*, and by a few fibers from the inner side of the inner tuberosity of the fibula. Its tendon, closely associated with that of the *tibialis posticus*, passes the inner malleolus and soon divides into two, one of which runs directly to the ental surface of the large tendon of the *flexor fibularis* and the other joins the small tendon of the *accessorius* and after a lateral connection with the main tendon of the fibular flexor it continues to an insertion on the terminal phalanx of the first digit.

Peroneus brevis.—Origin tendinous from the inner tuberosity of the fibula and proximally by septa with the muscles lying on either side and

beneath it. Its large tendon passes beneath the annular ligament outside the external malleolus and after passing over the front of the calcaneum turns downward and inserts on the outer side of the tuberosity of the base of the fifth metacarpal. Just above its insertion it is distinctly increased in size and is perforated by the tendon of the *peroneus tertius*. The same arrangement has been noted in *Trichosurus* (Thompson and Hillier, 1905, p. 322).

Tibialis anticus.—Origin from the deeply excavated outer surface of the proximal part of the tibia. Its muscular part lies mostly in this broad groove but is directly attached to the bone only in its proximal third. Its principal tendon passes along the fibular edge of the tibia and after passing over the astragalus turns down entad of the tendon of the *extensor hallucis longus* and widening slightly inserts on the under side of the entocuneiform bone. A second and smaller tendon runs from an indistinct division of the fleshy part of the muscle and, paralleling the larger tendon, inserts in front of it on the side of the base of the first metatarsal.

In *Myrmecobius* (Leche, Bronn's *Thierr.*, VI, p. 893) it is joined by the tendon of the *extensor hallucis longus*, but in this case that muscle and its tendon are independently developed. An accessory tendon to the first metatarsal similar to that in *Cænolestes* was found in *Notoryctes* by Thompson and Hillier.

Peroneus tertius.—Origin from the external condyle of the femur and the under surface of the collateral ligament jointly with the *peroneus longus*, from the outer surface of the inner condyle of the fibula and also by a distinct slip or second head from the surface of the fibula just distad of the outer tuberosity. It lies entad of the *peroneus longus* and although not so completely bicipital there is a distinct correspondence between the divisions of each. It has only slight connection with surrounding muscles, being distinct throughout its fleshy extent. Its tendon passes through the malleolar ligament, perforates the tendon of the *peroneus brevis* and continues directly to its insertion in the fascia on the outer side of the base of the first phalanx of the fifth digit.

This muscle is regarded by Leche as a division of the *extensor brevis digitorum*. Its relations in *Didelphis* are similar to those in *Cænolestes* and also in *Thylacinus*, *Myrmecobius*, *Cuscus*, and in *Phascolarctos*, but in these latter it has no femoral origin (Leche).

Peroneus longus.—Arises by two heads which are separate proximally but which join midway of their fleshy extent to form a single tendon. The inner head lying next to the *peroneus brevis* takes origin from the inner tuberosity of the fibula and from the outer edge of a collateral ligament running from the external condyle of the femur to the inner tuber-

osity of the fibula. A few fibers also arise from the outer base of the *peroneus brevis*. The outer head arises from the outer tuberosity of the fibula and the inner surface of the fibular sesamoid. Its tendon lies deep to that of the *peroneus brevis* and passes with it through the annular ligament over the external malleolus. Distad it passes over the peroneal tubercle of the calcaneum and down the side of the cuboid deeply imbedded in tissue. Turning to the under side of the tarsus, it passes over the cuboid and after being somewhat enlarged and attached near the base of the fifth metacarpal, it passes the bases of the third and fourth metacarpals and goes to insertion on the inner base of the hallux. Its course and insertion are thus much the same as in *Phascolarctos*, *Phalanger* and *Didelphis*.

Tibialis posticus.—A distinct muscle with its fleshy part lying between the *flexor digitorum fibularis* and the *flexor digitorum tibialis*. Its origin is from the inner side of the outer tuberosity of the fibula, from the expanded inner surface of the fibula and from septa of the fibular flexor. Its tendon runs beneath that of the *flexor digitorum tibialis* and crosses it midway to the ankle finally inserting on the palmar surface of the scaphoid.

Extensor brevis digitorum.—This is in three distinct divisions two of which are of fibular origin while the third is on the dorsum of the foot. The first is a thin flat muscle arising from a small space on the outer expanded surface of the fibula slightly mediad and distad of the outer condyle. Its tendon runs through the outer malleolar ligament and inserts in the fascia on the outer side of the base of the first phalanx of the third digit. The second division is slightly larger in its muscular part which arises subjacent to the first and between it and the *extensor hallucis longus*. Its origin occupies the distal two-thirds of the proximal half of the outer expanded surface of the fibula. The proximal part of the fibula between the tuberosities is without muscular attachments and directly in relation with the inner surface of the *peroneus tertius* which overlies the above-described divisions of the short extensor. The tendon of the second division passes the outer malleolus with that of the first and proceeds over the dorsum of the pes to insertion in the fascia on the outer side of the base of the first phalanx of the fourth digit. These two divisions obviously belong to the peroneal group of muscles and are quite as distinct as the others. They are united in *Didelphis* and *Dasyurus* and send tendons to the second, third and fourth toes; in *Thylacinus* and *Myrmecobius*, they are in three divisions (Leche).

A very slender tendon imbedded in fascia crosses the third metatarsal and connects the tendon of the second division with that of the third or pedal division. The third division or *extensor hallucis brevis*

consists of a short flat muscle much involved in fascia and arising chiefly from the dorsal surface of the fibular side of the distal annular ligament and the fascia over the os calcis. Its fleshy fibers scarcely reach beyond the base of the metatarsals and its tendon runs to the fascia on the fibular side of the base of the second metatarsal. Fascial connection with the hallux may include a minute tendon but this was not distinguished. A similar *extensor hallucis brevis* is found in *Didelphis*, but it is lacking in *Dasyurus*, *Thylacinus* and *Myrmecobius*. Its presence in *Cænolestes* appears to be an exclusive resemblance to *Didelphis*. It is common, however, in other mammals, as for example, the *Dasypodidae* among edentates.

Extensor longus digitorum.—The digital extensors lie next to the *tibialis anticus* and take origin from the outer tuberosity of the tibia, the inner tuberosity of the fibula, and septa joining them to the *peroneus brevis*. Their origin is more largely from the fibula than the tibia. They are united for no more than the proximal third of their fleshy extent and in the remainder of their length are distinct both as to the distal fleshy parts and the tendons. Passing over the ankle they are bound down in common with the *tibialis anticus* and the *extensor hallucis* by a broad annular ligament (*ligamentum cruciatum*) and also by a short stout annular ligament (*ligamentum fundiforme*) which serves for the passage only of their four ligaments. The most deep-seated of these extensors is that lying next to the *tibialis anticus* and its tendon goes to the terminal phalanx of the second digit, the next to the third digit and the others to the fourth and fifth, the tendons passing on the inner side of the second phalangeal joint in each case and thence to the dorsal surfaces of the third phalanges.

Extensor hallucis longus.—Origin from the inner edge of the fibula just distad of the inner tuberosity and thence about one fourth of the way to the junction of the fibula and the tibia. A few fibers also spring from the popliteal fascia. Its thin flat muscular part passes beneath the common digital extensor and the peroneus muscles in a diagonal direction from the head of the fibula to the base of the tibial flexure and thence its tendon passes down the tibia beneath the large annular ligament and inserts on the base of the first phalanx of the hallux.

Popliteus.—Origin from the greater part of the inner surface of the proximal half of the tibia. Proximally it reaches nearly to the inner condyle of the tibia while distally it extends almost to the union of tibia and fibula. Its fibers run diagonally across the interosseous space but do not reach the fibula except proximally. Distally the fibers reach not more than half way across the interosseous membrane to which they are attached as a thin nearly transparent sheet. Proximally the muscle has

tendinous attachment to the posterior side of the inner tuberosity of the fibula and the tendon continues broad through the muscle, which is pinnatifid. An indistinct layer lies superficial to this part of the muscle and is also pinnatifid, but in reverse direction, the tendon attaching to the shaft of the tibia and the fibers radiating toward the fibula. This may represent the so-called *pronator tibiae*, indications of which are found also in *Myrmecobius* and *Didelphis* (Leche, p. 899), forms in which there is some possibility of rotatory movements between tibia and fibula, whereas in *Cænolestes* such movements are quite precluded by the closely appressed distal halves of the bones. The retention of these muscles in *Cænolestes*, therefore, is possible evidence that the arrangement of the bones is a recent specialization. This is partly negated, however, by the entire absence of the *pronator quadratus* in the foreleg. There is no trace of a distal *pronator tibiae* such as is described by Macalister (1870, p. 173) for *Phascolumys*, *Sarcophilus*, *Phalangista*, and *Perameles*.

MUSCLES OF THE HIND FOOT.

The number and relations of the muscles of the hind foot, the so-called intrinsic muscles, are broadly similar to those of various other pentadactylous marsupials. The size and independence of the hallucal adductor and the divided *abductor indicis* are noteworthy, as well as the sesamoid origin of certain of the short flexors. Although the other adductors and abductors are present it seems doubtful whether many of them can be more than very slightly functional on account of the closely appressed metatarsals. These are almost in contact proximally for nearly half their length in such a way as to permit of only very slight lateral motion.

Lumbricales.—These are fairly well developed, but of uneven size, arising from the sides of the tendons of the *flexor digitorum fibularis* and inserting with them at the ends of the toes. Each tendon has two of these small muscles. Those on the tendons of the second, third and fourth digits are progressively shorter and rise from the sides of the tendons a considerable distance distad of their separation from the common tendon. Those of the tendon of the fifth digit are very unequal, the outer being short and rising far distad while the inner is long and rises as high as the point of separation of the tendon from the common tendon.

A single lumbrical rises on the medial side of the *flexor digitorum tibialis*. Another thin muscle rises on the ventral surface of the common tendon of the fibular flexor just above its digital divisions and sends weak aponeurotic connection to the fourth digit and possibly also to

others but this could not be determined. It represents, at least in part, the *flexor digitorum communis brevis*, which is similarly but more highly developed in *Didelphis*.

Adductors.—All the adductors except the *medius* are present. Those of the outer digits arise in the membranous fascia just proximad of the metatarsals. The *indicis* goes to the medial or fibular side of the first joint of the index digit, the *annularis* goes to the tibial side of the same joint of the fourth digit, and the *minimi digiti* to the tibial side of the fifth digit. In one specimen the *indicis* and *annularis* were joined proximally for at least half their length, the *minimi digiti* being distinct from the base of the metatarsus. They are rather flat, wide, and quite delicate, but have small central tendons.

The *adductor hallucis* lies along the lateral surface of the first metatarsal between the *abductor hallucis* and the short flexors and is the largest of the intrinsic hallucal muscles. It is not connected with the other adductors but has independent origin from the scaphoid by a relatively stout tendon which passes along the grooved surface of the ectocuneiform becoming fleshy at the base of the metatarsal. It inserts with the abductor on the tibial side of the distal extremity of the first metatarsal. The *adductor hallucis* is thus quite different from that of *Didelphis* and somewhat similar to that of *Phascologale* as described by Cunningham (1882, p. 55), but is even more independent, its origin being completely separate from that of the other adductors.

Abductors.—These consist of an *abductor hallucis*, *abductor minimi digiti* in two divisions, *abductor ossis metatarsi minimi digiti*, and four interosseus muscles.

The *abductor hallucis* is rather poorly developed and muscular only in its short fan-shaped proximal part. It arises in the fascia surrounding the small sesamoid laterad of the base of the first metatarsal and sends a thin tendon to insert on the radial side of the distal extremity of the same phalanx.

The *abductor ossis metatarsi minimi digiti* is well developed arising on the os calcis and running laterad of the plantar tendon to insert on the tuberosity of the fifth metatarsal.

The *abductor minimi digiti* is in two divisions. The first arises as a thin fan-shaped fleshy muscle from the os calcis ectad of the foregoing and soon becoming a slender tendon continues by a marginal course to its insertion on the radial side of the distal extremity of the fifth metacarpal. The second division, which is muscular throughout, arises from the sesamoid at the inner (medial) base of the fifth metacarpal and inserts on the outer (radial) sesamoid at the distal extremity of the fifth metacarpal, passing between the divided short flexors. The external

plantar nerve passes entad of it near its origin and its general relations are much as described by Cunningham for *Dasyurus*, *Phascologale* and *Thylacinus* but it is a relatively larger muscle than in these, rather thin and flat, and it seems not impossible that it may be an *opponens minimi digiti*. The usual four dorsal interossei are present but have a tendency to fusion with the short flexors. With the exception of the first, they are entirely palmar in position and cannot be seen from the dorsal aspect of the foot owing to the nearly united basal metatarsal bones.

The *abductor indicis* is in two divisions. The larger has tendinous origin as high up as the scaphoid near the origin of the *abductor hallucis* and continues tendinous to the medial base of the first metacarpal where it has slight attachment and becomes a well-formed fleshy belly running diagonally to insertion on the side of the first joint of the index digit. The second and smaller division of this abductor arises on the medial side of the base of the hallux and runs close to the *flexor indicis* slightly dorso-mediad of the first division and joins the tendon of the first division on its medial side slightly before its insertion. This two-headed arrangement of the *abductor indicis* is reported for *Thylacinus*, a tetradactylous animal, and a somewhat similar condition is found in *Phalanger* which is syndactylous. In *Didelphis* and *Dasyurus* the *abductor indicis* is single-headed.

The third metacarpal has the usual two abductors inserting on each side of the first joint on the sesamoids. They are dorsad of the flexors at their extremities but mostly laterad in the middle where they are visible without disturbance of the flexors. The outer one is rather closely associated in its proximal half with the *flexor brevis* of the fourth metacarpal. The fourth interosseous is less dorsal in position than the others and runs nearly laterad of the flexor from origin in the palmar cartilage to insertion on the outer side of the end of the fourth metacarpal. The interossei insert very close to the extensor tendons but do not appear to be definitely connected with them. Near the insertions of the second, third and fourth interossei there is a tendinous connection with the joint opposite the one on which they are directly inserted and thus the ends of the metacarpals are bound together and necessarily have but little freedom of movement.

Short flexors.—There are five pairs of short flexors, one for each metatarsal. They are only slightly connected at their origin in the palmar fascia just above the bases of the metatarsals and thence to their insertions on the distal sesamoids the divisions are clearly marked. Those of the hallux are weaker than the others and the lateral division is partly aponeurotic and smaller than the median division. Those of the second and third digits enclose a small flattened sesamoid in the fascia

where they arise. This sesamoid may not be homologous with that of *Notoryctes*, but it is otherwise unique among marsupials and, as noted by Thompson and Hillier,¹ is suggestive of certain edentates. The flexors of the fifth digit are separated at their insertion by the tendon of the shorter division of the *abductor minimi digiti*. It is somewhat doubtful as to whether the outer or radial division is properly to be regarded as a flexor, especially since it inserts by a slender tendon whereas its fellow has a broad fleshy insertion extending to both sesamoids. Possibly this supposed outer division of the fifth flexor should be regarded as a third division of the *abductor minimi digiti*.

SUMMARY OF MYOLOGICAL CHARACTERS.

The trunk muscles of *Cænolestes* are, as was to be expected, of the prevailing marsupial type. It is in the muscles of the limbs and head that the most significant deviations are found and although these seem quite numerous when comparison is made only with American forms, there is such variation among Australian forms that similarity is found with first one and then another. It is quite evident that some Australian forms, especially in the Dasyuridae, approach *Didelphis* more closely than does *Cænolestes*. It is also clear that many of the points in which *Cænolestes* differs from *Didelphis* are to be found in Australian forms. Although some of these resemblances are plainly cases of convergence, others do not seem so. Therefore, aside from the question as to the nearest relationships of *Cænolestes*, it is evident that it contributes in a positive way to the broad general conclusion, already well grounded, that American and Australian marsupials are of common derivation. That is, many of the points in which *Didelphis* fails to show resemblance to Australian forms are supplied by *Cænolestes*.

From the standpoint of myology there seem to be few if any distinctions that can be definitely drawn between Polyprotodontia and Diprotodontia. Nevertheless, in viewing the possibility that *Cænolestes* is to some extent transitional between the two large groups, the musculature can scarcely be adduced as contrary evidence, since it shares many features with the diprotodonts that are not possessed by all polyprotodonts. That it shares more than any polyprotodont cannot be said,

¹ Regarding the sesamoid in *Notoryctes*, these authors say: "In connection with the plantar sesamoid bone, which as far as we know has not been recorded in any other marsupial, it is interesting to find that in some edentates there is a very similar arrangement. Windle and Parsons state that in the Dasyopodidae the tibial and fibular flexors unite in the lower part of the leg, and are inserted into a very large sesamoid bone in the sole of the foot, which is held in place by a fibrous band from the calcaneum, the equivalent of the accessorius. From the front of the sesamoid bone five tendons pass to the terminal phalanges of the five digits." Jour. Anat. & Phys., XXXIX, p. 326, 1905.

although better comparisons than are here possible might prove this to be the case. It is worthy of particular note that it has a considerable number of points in common with the extraordinary form *Notoryctes* and that, as in *Notoryctes*, some of its most exclusive features as a marsupial are suggestive of edentates and insectivores. Among such are the divided *trapezius*, the *dorso-cuticularis*, the absence of a *pronator quadratus*, the sesamoid origins of the *supinator* and the short flexors of the hind foot, and the peculiar *sartorius*. It is also to be noted that some of the resemblances between *Cænolestes* and *Didelphis* not shared by most other marsupials are however to be found in other mammals, especially edentates, and therefore cannot be taken as certain indications of close relationship between *Cænolestes* and *Didelphis*. The well-developed pedal *extensor hallucis brevis* and the *flexor accessorius pedis* are perhaps examples of this kind.

The muscles of the legs and feet in *Cænolestes* are adapted to a terrestrial, almost cursorial, life. The leg muscles have short thick fleshy parts and very long tendinous extensions, relatively longer even than in such terrestrial forms as *Phascogale*, and in this respect are perhaps most similar to those of the saltatorial but otherwise generalized Australian polyprotodont *Antechinomys*, the myology of which has not been thoroughly described. The proportion of tendinous to fleshy parts is about as 60 to 40 and the outline of the leg thus resembles that of a digitigrade animal. It is markedly different from that of *Didelphis* in which fleshy extensions reach nearly or quite to the carpus and tarsus. Specialization for terrestrial life has proceeded far beyond that in *Peromys* whose habits are presumably not greatly different from those of *Cænolestes* but whose musculature is only slightly different from that of *Didelphis*.¹ The absence of the *pronator quadratus* is not accompanied by such reduction of other muscles as might be expected in view of the facts that otherwise a large number of muscular elements are present in the limbs and that their distinctness from each other is marked. There is some reduction of the intrinsic muscles of the feet, especially in the adductor sets, and the grasping power is distinctly limited.

All the important muscles have been worked out with considerable care, but the nature of the material, the small size of the animal, and the lack of comparative material, have made it impossible to be certain of relations in all cases. Such comparative notes as are ventured are based on examination of the scattered literature and the comprehensive work of Leche in Bronn's *Thierreich*. Comparison of actual specimens has only been possible with *Didelphis*, *Marmosa*, and to some extent with

¹ No careful study of *Peromys* has been possible, but cursory examination of its feet and legs indicates close similarity to *Didelphis*.

Peramys and *Macropus*. That this is not an ideal method of comparative work can be apparent to no one more fully than the author; but, it is the best that circumstances permit and if small errors have crept in it is hoped that they will not seriously affect general conclusions. The nomenclature adopted has been that of Bronn's Thierreich with occasional changes where it seemed feasible to conform to the BNA terms.

Features in the myology of *Cænolestes* which seem especially noteworthy are as follows:

Trapezius divided into *spinotrapezius* and *acromiotrapezius*.

Rhomboideus divided.

Dorso-cuticularis present.

Distinct *rectus capitis posterior superficialis*.

Pterygoideus externus relatively large and bicipital.

Subclavius not extending to scapular spine.

Median head of *triceps brachii* distinct.

Origin of *supinator brevis* including a sesamoid.

Pronator quadratus absent.

Distinct *caudo-femoralis* present.

Femoro-coccygeus practically distinct from *gluteus maximus*.

Gluteus medius partially divided into two layers.

Sartorius on inner side of thigh, much reduced, and originating from Poupart's ligament.

Extensor carpi ulnaris with ulnar as well as humeral origin.

Radial extensors of carpus separate from each other.

Median and lateral heads of *gastrocnemius* partly united.

Fibular divisions of *extensor brevis digitorum* distinct from each other.

Pedal *extensor hallucis brevis* well-developed.

Flexor accessorius present.

Pronator tibiae partly separate from *popliteus*.

Short flexors of second and third toes arising from a sesamoid.

URINOGENITAL SYSTEM.

The interesting urinogenital system has been studied only by gross dissection and the use of the binocular microscope. With the scanty material available and the means at hand, no other course seemed possible. It is to be regretted that serial sections could not be obtained of the embryo. One of the imperfect females examined was pregnant, having three embryos, two in the right uterus, and one in the left. The uteri were greatly expanded and the foetuses lay surrounded by their

thin membranes and the thickened walls of the uteri. The largest was approximately 4 mm. in length and well advanced in development. By a most unfortunate accident, these embryos, after being removed and set aside for special study, were totally destroyed.

Mere gross dissection, however, is sufficient to show that the reproductive system of *Cænolestes* is markedly different from that of the primitive polyprotodonts.

MALE GENERATIVE ORGANS.

Plates VIII-X.

The prostate gland is exceedingly large, relatively larger than in any other marsupial known to me and therefore probably larger than in any other mammal. It is a prominent ovate body occupying practically the entire space in the posterior part of the body cavity and extending from the kidneys to the pubis. In a normal adult with a body length of 113 mm., it measures 16 mm. in length and 11 mm. in greatest diameter. The bladder, which is small and inconspicuous in comparison, is situated on the dorsal surface of the prostate somewhat caudad of its anterior expansion and thus scarcely visible from the ventral aspect. The gland is not tapering and radiform as usual but is only slightly smaller posteriorly than anteriorly and the narrow urethral canal makes exit from its walls abruptly on its dorso-caudad surface. The prostate is enclosed in a thin membranous or slightly muscular covering of transverse fibers. The thickened portion consists of numerous slender cylinders of glandular tissue radiating from the urethral canal to the periphery. The urethra in its course through the prostate is quite small and is longitudinally plicated on its inner surface (Pl. X, Fig. 5).

The ureters and vasa deferentia enter a groove between the base of the bladder and the prostate, the former going to their orifices on the dorsal side of the bladder near its neck and the latter turning sharply caudad into a groove in the prostate to a common opening into the urethra a short distance beyond its enclosure by the prostate (Pl. X, Fig. 1). The vasa deferentia have the usual course, leaving the base of the testes and diverging to enter the body cavity and pass dorsally on each side of the colon to their terminations. Before entering the raphe in the prostate they are somewhat enlarged and slightly convoluted. There are no obvious seminal vesicles.

From its exit from the prostate the urethra continues as a slender tube for about 5 mm. to its bulbous portion where it enlarges and receives the ducts of the Cowper's glands. This bulbous urethra is short and wide and has a slight forward projection or cul de sac on its ventral

side. There are three pairs of Cowper's glands. The anterior pair are elongate pear shape and open into the anterior part of the bulbous urethra ventro-laterally. These have been designated in the figures as Cowper's gland No. 1. The two other pairs are situated dorso-laterally just anterior to the base of the stalk of the corpus spongiosum and have a common opening (Pl. IX, Fig. 1). In fact the smaller one (No. 2) appears to be scarcely more than a division of the larger (No. 3) since they unite at an appreciable distance from the urethra. This gland No. 3 is very large, several times larger than the others, even larger than the paired testicles, and more than half the size of the enormous prostate. It measures 12 mm. x 8 mm. Although not examined histologically, its superficial appearance indicates a structure somewhat different from that of gland No. 1. Gland No. 2 appears to have the same structure as No. 3, of which, as noted above, it is scarcely more than a division. In *Marmosa* and *Didelphis* there are only two pairs of Cowper's glands, but in *Peromyscus* I find three as in *Cænolestes*, the posterior pairs being on a common stalk but showing no such discrepancy in size as in *Cænolestes*. The substance of gland No. 1, except near the base, appears as nearly solid, slightly granular tissue of a whitish color, whereas No. 2 and No. 3 are highly fenestrated and spongy and even in gross examination are seen to consist of numerous thin-walled tubules which converge toward the duct and divide into branches toward the periphery of the gland. In the glands of *Marmosa* and *Didelphis*, the same difference of structure obtains. In a pouch specimen of *Macropus* there are three glands, as in *Cænolestes*, and the posterior ones have a common stalk, but their structure seems identical with that of the anterior pair. Histological study of fresh material is necessary before the significance of the differences in these glands can be understood.¹ That some of them may function as seminal vesicles is perhaps not impossible.

The corpus cavernosum has its origin in a bulb invested with muscles and situated slightly caudad of the first pair of Cowper's glands. The stalk of this bulb, which is slightly bilobate, expands into a layer around the ventral wall of the urethra and continues to the base of the glans penis. The muscle covering the bulb of the corpus cavernosum turns on itself and attaches on the stalk of the bulb instead of continuing to form a levator penis as in the opossum. However, there are three pairs of small thin muscles lying flat on the base of the corpus cavernosum and the adjoining part of the bulbous urethra, some one of which may be the homologue of the levator penis, although this seems

¹ Such work may have been done already, but the scope of the present study has not extended to the literature in which it might be recorded.

doubtful (Pl. IX, Fig. 2). In the opossum the levator penis arises as a direct continuation of the muscles covering the postero-ventral part of the lateral bulb of the corpus cavernosum and proceeding inward turns abruptly caudad and continues to the base of the glans penis. There is no muscle occupying this position in *Cenolestes* and none that could function as a levator penis. There is another paired muscle in the opossum, of which the function is not apparent, which corresponds closely in position to one of the small paired muscles in *Cenolestes*, the most anterior and superficial one. These small muscles have been designated in the figure as Nos. 1, 2 and 3. No. 1 arises along the median line on the anterior part of the bulbous urethra and continues as a slender, free, and slightly thickened fascia to the anterior surface of the muscle on the bulb of the corpus cavernosum. No. 2 lies partly beneath No. 1 and overlaps No. 3, which covers the anterior side of the stalk of the corpus cavernosum. There is no trace of any such muscles in *Macropus*. From the ventral side of the corpus cavernosum near the extremity of the stalk a large flattened tendon extends to an attachment on the postero-internal surface of the ischium opposite the origin of the *gracilis* muscle (Pl. IX, fig. 2, *i. t.*). No such tendon is found in *Didelphis*, but a similar attachment is normal in *Macropus*. A plexus of nerves, blood vessels, and connective tissue emerges between the *pubococcygeus* muscle and the ischium and reaches the corpus cavernosum and adjacent glandular surfaces, assisting in attaching the very large postpubic mass to the body.

Behind the bulb of the corpus cavernosum but more dorsal in position is the bulb of the corpus spongiosum, its stalk meeting the bulbous urethra slightly below (ventrad) the common opening of the posterior Cowper's glands. It is also enclosed in a thick layer of muscles but these do not extend beyond its stalk which expands within the dorsal wall of the urethra in a very thin layer. This continues to the base of the glans penis where it rapidly thickens and extends to the end of the glans. The corpus cavernosum continues, forming the central part of the glans, practically to the pointed tip.

The glans penis is so deeply cleft as to appear like two separate organs. Each half is of acicular shape, narrowing at the tip to a sharp point. The extreme tip is separated by a circular fold from the main body of the glans as a sort of secondary glans. In front of the base of the glans the urethra forks to empty into a deep groove which runs down the dorsal surface of each division of the glans, terminating at the circular fold which marks off the tip of the organ. This groove is expanded in its ventral part and occupies the center of the glans. In cross section in a preserved specimen it appears as a transverse opening connected with

the dorsal surface of the glans by a vertical slit. In life it would doubtless function almost as efficiently as a closed tube. The surface of the glans is slightly rugose in its proximal two-thirds and distally it is covered with small circular papillae. The total length of the cleft part of the glans is approximately 12 mm. The penis when retracted lies in a pouch formed by the expanded distal part of the urethra and opening into the cloaca (See Pl. IX, Fig. 3).

The retractor muscles of the penis are attached one on each side near the anterior end of the penis pouch. They originate from the narrow tendons of the *infracoccygeus* on the ventral surface of the fifth lumbar vertebra and run for a short distance caudad above the colon and then turn slightly ventrad, inclosing the colon and continuing caudad along its latero-ventrad surface, and pass dorsad of the Cowper's glands to their insertion on the sides of the penis pouch.

The cloaca, the penis pouch, and at least certain of the Cowper's glands are enveloped in a thin layer of muscle, the sphincter cloacae and its accessory folds. About the cloaca it is thicker and the fibers have a circular course, but more anteriorly it becomes thinner and follows the interstices between the organs where it is lost in the connective tissue. Anteriorly it is attached by the flat tendons to the ventral surface of the pubic symphysis. These, in crossing the ventral surface of the urethra, have partial attachment near the base of the corpus cavernosum.

The testes are of relatively large size, broadly elliptical in shape and slightly flattened on one side where the two are closely appressed in the scrotum. A single one in a preserved specimen measures 6.5 x 4.5 mm. The testis is clasped by the epididymis in the usual manner and loosely attached. The globus major is somewhat thickened and has considerable convexity. It is slightly larger than the globus minor which lies folded upon itself and in its course toward the exit of the vas deferens has several convolutions which lie against the testis in the inferior curvature of the epididymis. Part of this is evidently the vas deferens itself and its position here is evidently somewhat peculiar since the arrangement in *Didelphis* and *Macropus* is not quite the same.

The most noteworthy features of the male generative organs of *Cænolestes* are the extraordinary size of the prostate and Cowper's glands, the very deeply cleft glans penis, and the absence or great reduction of the levator penis muscle. It is difficult to see indications of relationship in any of these. The bifid glans penis occurs in both polyprotodonts and diprotodonts. It is most pronounced in polyprotodonts and, according to Owen (Anat. Vert., III, p. 648), is associated with the presence of a levator penis muscle. But in *Cænolestes* there is the unusual condition of a deeply bifid glans penis and a rudimentary

levator muscle. The divided glans and the well-developed levator muscle are doubtless both to be regarded as generalized conditions and it may be concluded that if *Cænolestes* has retained one it is at least in a secondary stage as regards the other. It is by no means certain that any of the small paired muscles on the cephalo-ventral surface of the bulbous urethra are homologous with the levator penis, but if not, then that muscle, which is so well developed in *Didelphis* and some other polyprotodonts, is entirely absent in *Cænolestes* as it is in the Macropodidae.

FEMALE ORGANS.

Plate VIII.

The female specimen dissected was a virgin. The ovaries were found lying entirely exposed on the latero-ventral surface of the colon slightly clasped by the fimbriated funnel of the oviduct. They are smooth and disklike, about 2 mm. in diameter. The Fallopian tubes are rather short and heavy as compared to those of the opossum. The uterine bodies also are relatively short and less flattened than in *Didelphis*. They appear simply as gradual expansions of the Fallopian tubes with only a slight line of division. They are shorter than the outstretched vaginal canals and measure approximately 6 mm. in length from the end of the oviduct to the base of the median vaginae. The general relations of the peritoneal folds forming the uterine fossa and the broad ligament are as in other marsupials.

The round ligament is well developed. In gross dissection its relations to the ovary and the uterus could not be determined positively. Apparently it does not extend directly to the uterus but is lost in the thickened part of the broad ligament to which the ovary is attached and from which a well-defined branch runs to the anterior part of the uterus. Possibly this branch is a continuation of the round ligament, but this could not be determined. This round ligament is attached by a membranous fold which extends in directly transverse relation to the broad ligament and thence to the dorso-parietal wall of the body cavity. Its antero-ventral border is rounded and about equal in diameter to the oviduct. Seen from the ventral side it passes at right angles beneath (dorsad) the folds of the oviduct and the broad ligament and makes a sharp turn cranial to join the thickened mass upon which the ovary rests. It is so distinct and well-developed that it does not seem likely that it can be anything other than the homologue of the round ligament of higher mammals. Its occurrence is one of the most noteworthy features of the female urinogenital system of *Cænolestes*. I am unable to detect any trace of a similar ligament in *Marmosa*. A round ligament,

apparently of similar character, is recorded by Hill (Proc. Linn. Soc. N. S. W., XXIV, p. 46, 1899) for *Perameles* and *Macropus*.

The bladder in its posterior half and the adjoining part of the urogenital sinus have their ventral surfaces connected with the abdominal wall at the linea alba by a vertical membrane, the "Ligamentum vesicae medium" of Brass and Hill. The bladder is also attached at the base laterally by a fold on each side running forward into the broad ligament. The ureters pass dorsad of the uteri and dorso-laterad of the uterine neck, enveloped in the broad ligament, and enter the base of the bladder dorso-laterally, passing between the vaginal cul de sac and the lateral vaginal canals and parallel to them.

The uterine necks are very closely attached to each other along the median line but may be separated without destruction of their walls to the beginning of the median vaginae. The os tincae, however, are completely coalesced and in the virgin female examined they are of extraordinary size, extending the full length of the vaginal cul de sac and having thick, slightly rugose walls. They are attached dorsally and ventrally by strong septa, forming two distinct lateral vaginal chambers. In an older female they were found to be relatively much smaller.

The thin-walled vaginal cul de sacs extend nearly to the base of the bladder on its dorsal surface and are attached to it and to the urogenital sinus by connective tissue, but without direct communication in the virgin female specimen examined. In a pregnant and probably multiparous female a possible connection was observed between the median vaginae and the urogenital sinus. An irregular opening was found but some destruction of tissue had occurred before the specimen was examined, so the point needs verification. If this should prove to be of regular occurrence it would be an important indication of relationship to the diprotodonts, for it is essentially characteristic of most of them.

The lateral vaginal canals leave the median vaginae in a dorsal direction and turn sharply caudad and then craniad in several convolutions, nearly to their origin, where they again turn and run directly caudad to their endings on the lateral walls of the urogenital sinus posterior to the ureters. The length of one of the canals fully extended is 10 mm. or two-thirds longer than the uteri. In this respect, therefore, *Cænolestes* shows some approach to the condition in *Macropus*, where the vaginal canals are always longer than the uteri. The urogenital sinus from the base of the ureters to the cloaca is 9 mm. long in a young female.

The rectum and the urogenital sinus open into the common cloaca at about the same level, 3 mm. from the lips of the external orifice. The clitoris is situated in a pocket of membrane on the dorsal wall of the

sinus and is a highly developed organ, deeply bifid, and attached only at its base. It consists of two long, smooth, and sharply pointed rods the apices of which reach nearly or quite to the cloacal opening of the urogenital sinus. Each of these divisions is sharply and deeply grooved on the dorsal surface of its basal half. The entire organ is 3.5 mm. in length and of this each fork is 3 mm. The form of the clitoris, therefore, is not very unlike that of the divided glans penis of the male. In an old female, the clitoris appears to be a single organ, hanging free, and having several longitudinal plications and a transverse cleft behind its rather blunt apex.

The cloaca is inclosed in an extensive muscle, the *sphincter cloacae*. It surrounds the cloacal orifice and spreads over the large rectal glands, terminating anteriorly in a transverse plexus of nerves and muscle fibers which is attached to the ventral surface of the urogenital sinus and which sends fibers and fascia into the pelvic cavity. The relations of the rectal glands have been described elsewhere (p. 76).

From the comparative standpoint, the female reproductive organs of *Cænolestes* are more significant than those of the male. They are distinctly of a diprotodont rather than a polyprotodont type. This is evidenced at once by the deep median vaginae and the long lateral vaginal canals. Many polyprotodonts give birth to their young by way of the lateral vaginae. In fact this seems the only method possible in such forms as *Didelphis* which have extremely small median vaginae. Certain others do not conform to this rule, as *Perameles*, which Hill (1899) has shown to have a parturition somewhat less perfected but similar to that of the Macropodidae, in which the foetus has direct passage from the median vaginae to the urogenital sinus and thence to the exterior. In certain other respects the reproductive system of *Perameles* is regarded by Hill as primitive. Data are not available as to the phenomena of parturition in all diprotodonts, but the direct median passage has been observed, according to Fletcher (fide Hill, 1899, p. 74), in no less than twelve species of Macropodidae. Hill found it also in *Tarsipes*, and, although it is evident that much study of well prepared serial sections is necessary for a thorough understanding of the subject, the inference is very strong that those forms having large median vaginae and relatively long convoluted lateral canals give birth to the young by median rather than lateral passage, some by an opening from the median vaginae to the urogenital sinus which becomes permanent after the first parturition, and others by a rupture at this point and subsequent closing for each parturition, the connection being made through a "pseudo-vaginal canal" in the connective tissue. This method of parturition is at least functionally an approach to the method in the higher mammals having

a single vagina, but is regarded as a special development not necessarily leading to nor derived directly from the single eutherian vagina. That it should have been independently developed in different phyla seems highly improbable. Hill's statement (p. 74) in regard to this is as follows: "That this median passage has not been twice independently acquired within the Marsupial Class I am convinced, and its existence in *Perameles* in a condition so obviously unspecialized and in association with such a persistently embryonic condition of the genital organs, tends to suggest that its acquisition is of ancient date." Elsewhere he says: "Finally, as regards parturition, it seems to me that the foregoing discussion sufficiently upholds the conclusion that *Perameles*, in respect to the phenomena connected with that process, in no way stands alone amongst Marsupials as an aberrant and specialized type, but quite on the contrary, exhibits more primitive features in the mode of birth of the young than are shown by any other Marsupial hitherto described as possessing a direct median passage. That the direct passage in *Perameles* is in a much more primitive condition than that of Macropods, will, I think, be admitted without question. Indeed, the condition of the passage in *Perameles* can only, in my opinion, be regarded as the precursor of the Macropine one and as showing us in use today the earliest stage in the evolution of that direct median passage which reaches its highest development in the specialized Macropodidae."

So far as gross examination indicates, *Cænolestes* is more advanced than *Perameles* and, although final proof must await microscopic study of serial sections, all the present evidence points to the conclusion that the direct median passage is nearly or quite as well-developed in *Cænolestes* as it is in various Australian diprotodonts. In a virgin female, the median cul de sac extends almost to contact with the urogenital sinus and in an old female, openings were found in the vaginae and the adjacent surface of the sinus, but the connection between them was imperfect and possibly due to adventitious circumstances in the history of this particular specimen.

ALIMENTARY CANAL.

Mouth.—The hard palate has numerous transverse ridges. Including the one at the posterior edge of the palate, there are nine of these ridges that extend completely across the palate and five which do not but merely reach to varying points on either side of the median line (Pl. XVIII, Fig. 3). The ridge between the last molars has 8 to 10 forwardly directed protuberances giving it a somewhat serrate appearance. The ridge between or just behind the canines projects forward in

a sharp angle. Of the incomplete ridges, the hindmost is scarcely more than a rounded papule, the next one is posteriorly angled, two others are straight and another has its ends directed backward. This is the condition found in a male specimen. A female shows a similar but not identical arrangement, two of the short ridges being absent and one of the long ones, that between the third molars, being interrupted in the middle.

The soft palate and the naso-pharyngeal cavity are relatively long. The naso-pharyngeal isthmus is guarded on the dorsal side by several slender filaments directed anteriorly.

Tongue.—The tongue (Pl. X, Figs. 3 and 4) is rather slender and somewhat pointed at the tip. In a male specimen, it measures 22.5 mm. from the tip to the epiglottis and 4.5 mm. in greatest width. The under surface is sharply keeled and from the median ridge a longitudinal flap extends from the basal part of the tongue to the tip. A slight median raphe is indicated on its upper surface from near the middle almost or quite to the tip.

The upper surface is densely covered with coronate papillæ. No microscopic preparations were made, but under a strong binocular lens these coronate papillæ appear simpler than those of *Didelphis*, not being strongly hooked and including relatively few secondary hairlike papillæ. No differentiation of the anterior from the posterior coronate papillæ was detected (see Poulton, 1883, p. 618).

Fungiform papillæ are scattered somewhat irregularly among the coronate papillæ and are rather small and disklike. There is nothing comparable to the highly developed fringe of fungiform papillæ on the end of the tongue which is found in *Didelphis* and *Marmosa*.¹

The circumvallate papillæ are as usual three in number. They are about equidistant from each other and the angle formed by the posterior one is somewhat obtuse. Each has a peculiar retrorse process developed on its dorso-caudal surface, giving it a form resembling a helmet. Similar backwardly directed circumvallate papillæ were found by Poulton (l. c., p. 610) in *Dasyurus* and said by him to be unique in his observations of the tongues of various marsupials.

Filiform papillæ are present in small numbers in the usual position. No lateral organ could be discerned.

On the whole, the tongue of *Cænolestes* seems much more similar to

¹ This characteristic of these American forms appears not to have been noted previously, since the specimen (*Metachirus*) used by Poulton (l. c.) lacked the extreme tip. The tongues of *Didelphis* and *Marmosa* are broad at the tip and thickly studded with somewhat elongate and pedunculate fungiform papillæ, giving the end of the tongue a fringed or dentate appearance not reported for any other marsupials.

that of Australian than American forms. It agrees quite closely with Poulton's description of the tongue of *Dasyurus* and that in turn is said to be of the same type as *Halmaturus*.

Oesophagus.—The oesophagus from the larynx to the stomach is 30 mm. long. It has membranous attachment to the vertebrae and lies just dorsad of the aorta until it passes through the diaphragm and almost immediately joins the stomach in the middle of the lesser curvature. Its lining has four to six longitudinal plications which continue into the lumen of the stomach and hang free there. This condition of the lining of the oesophagus evidently has general similarity to that reported for most other marsupials except *Didelphis*. The transverse folds of the lower oesophagus, which are characteristic of *Didelphis*, are not represented in *Cænolestes*, but they are absent also in *Marmosa* and *Peramys*.¹ The lower oesophagus of *Marmosa* is smooth, however, and markedly different from that of *Cænolestes*.

Stomach.—The stomach (Pl. VII, Figs. 4 and 5) of a male specimen in somewhat distended condition measured 24 mm. in longitudinal diameter and 16.5 mm. in transverse diameter. It is regular in form, roughly elliptical, with the cardiac expansion only slightly exceeding the pyloric. In gross examination, the interior shows three marked divisions which may be called oesophageal, cardiac and pyloric, without regard to their histological structure. The first or oesophageal division is the smallest and consists of thin plications or lamellae surrounding the orifice of the oesophagus and extending along the lesser curvature to a point nearly midway between the oesophagus and the pylorus. The cardiac division consists of a strongly differentiated glandular area, forming a compound gastric gland comparable, at least in certain respects, to the gastric glands of *Phascolarctos*, *Phascolomys*, and *Manis*. It is developed principally on the sides of the stomach, being separate on the two sides toward the pylorus but uniting around the cardiac side of the oesophagus. In an undistended stomach, it is plainly evident from the exterior through the rather thin outer muscular walls. From within it appears as a thick, elevated, and well circumscribed area of glandular tissue inclosing 40-60 slitlike openings (Pl. VII, Fig. 5). Especially toward the cardiac end of the stomach, these openings have a distinctly linear arrangement and seem to be the result of an amalgamation of the usual rugae of the stomach. In an undistended specimen there is one prominent fold or ruga which extends three-fourths of the way to the pylorus and runs about midway between the compound gland and the greater curvature. The pyloric division of the stomach

¹ This is a rather important distinction between *Didelphis* and *Marmosa* which, so far as I am aware, has not been noted previously.

is relatively smooth and undifferentiated and is continuous with the area along the greater curvature to the fundus.

The differentiation of areas of the stomach in *Cænolestes* is of course a specialization and as such is referred to in discussion elsewhere, but in view of the wide range of the character of the stomach in marsupials, it is difficult to draw any direct inference of relationship. In fact, the stomach of *Cænolestes* is unique and serves to strengthen the general conclusion that the animal stands by itself quite as independently as any of the highly specialized Australian forms.

Intestines.—The intestines lie in the abdominal cavity with the main mass of the small intestines, or Meckel's tract, in ventral position. The duodenal loop is well differentiated. On leaving the stomach (from which it is separated by the pyloric sphincter, showing externally as a distinct band 2 mm. in width), it passes slightly dorsad, then caudad and turning sharply on itself runs cephalad a short distance ventrad of and almost in contact with the colon, and thence it passes ventrad to the center of the mass of Meckel's tract. Throughout this course it appears to have slightly thicker walls than the distal part of the small intestine and in an alcoholic specimen is lighter colored. The diameter of the collapsed duodenum just beyond the stomach is nearly 9 mm. The duodenum is weakly attached to the dorsal mesentery near the end of its loop and very close to the colon. The folds of the intestines are simple and on a continuous mesentery but cannot be straightened in a preserved specimen without cutting. They seem to be in the condition figured as stage II by Beddard (1908, p. 592, Fig. 121).

The large intestine is relatively very short and straight and has no indication of loops. It is scarcely more than 30 mm. in length as compared to 340 mm. for the small intestine from the pylorus to the caecum. The hindgut is imperfectly divisible into colic and rectal parts. The colon has the usual longitudinal folds on its inner surface but at its termination, that is at the beginning of the rectal part for a distance of about 4 mm., both the large primary folds and the interspaces are crossed by numerous delicate transverse ridges which in relaxed condition appear as undulating transverse lamellae. Externally the rectum is covered with longitudinal muscle fibers, somewhat more numerous dorsally than ventrally.

The caecum is very small, scarcely more than a vermiform appendix (Pl. VII, Fig. 3). In one specimen it is only 4.5 mm. in length; in another it is 5.5. It opens into the intestine by a simple aperture and its inner surface does not differ from that of the ileum. Immediately distad of its opening the colon begins with a muscular, plicate inner surface in contrast to the villous surface of the ileum.

The short colon combined with the small caecum in *Cænolestes* is contrary to the usual condition in marsupials and is most nearly met in certain of the dasyures. Although the short colon is perhaps primitive, the small caecum is obviously a secondary condition.

GLANDS.

Liver.—The liver is relatively large in size. It has three primary divisions, the left lateral, the right central and the right lateral, the last being subdivided into right lateral proper, caudate and spigheian (Pl. VII, Fig. 2). The largest lobe is the left lateral, although the right central or cystic lobe almost equals it and the combined bulk of the three subdivisions of the right lateral is nearly or quite as great as either of the other primary divisions. In a male specimen hardened by preservative, the right central lobe is 18 mm. wide; the left lateral has practically the same width but is slightly more bulky.

The large left lateral is entire without trace of a division, although a slight projection of its dorsal part is somewhat separated to curve around the oesophagus. In *Didelphis* and *Marmosa* there is at least one cleft in its lateral boundary partially dividing it into left lateral and left central.

The right central is entire except for one fissure extending about halfway to its base and inclosing the gall bladder in a wholly concealed position similar to that in some other marsupials but unlike that of *Didelphis* in which it is exposed and in contact with the diaphragm.

The right lateral proper lies next to the right central and is regular in shape although its dorsal border has a slight cleft. It is about half as large as the right central and its dorso-caudal border is in contact with the kidney. Although for the most part separate from it by a deep fissure, it is united at its base to the caudate.

The caudate is in ventral position and incloses a considerable part of the right kidney. It has an anterior projection and two posterior ones through the inner of which the vena cava passes.

The spigheian lobe arises from the base of the caudate and divides into two lobes almost at right angles to each other, one appressed to the left lateral and the other hanging free caudad of it and ventrad of the caudate lobe. Its peduncle attaches it to the caudate but the portal vein and the ductus choledochus pass between the free surfaces of the caudate and the spigheian. The pancreatic duct and the ductus choledochus unite just before entering the duodenum.

The liver of *Cænolestes* differs from that of *Didelphis* and at least some other polyprotodonts in its large bipartite spigheian lobe, in its

small concealed gall bladder, and in its complete fusion of left lateral and left central lobes. In *Didelphis*, the spigelian lobe is quite small and undivided. In *Antechinomys* (Beddard, 1908, p. 563) the spigelian is entirely absent and left lateral and left central are completely divided. A spigelian lobe is present in most of the Macropodidae and in one (*Dendrolagus*), at least, it is bilobate (Beddard, 1895, p. 134).

Spleen.—The spleen is suspended from the omentum and the dorsal mesentery and lies closely appressed to the dorso-caudad surface of the stomach. It is an elongate organ with a single triangular lobe directed dorsad from the right hand third of the dorsal border. The spleen thus has a general shape resembling a figure four or a bisected head of a blunt-pointed arrow (Pl. VII, Fig. 1).

Pancreas.—The pancreas is an extensive, much divided gland lying ventrad of the spleen and suspended between it and the intestines over the surface of which some of its smaller digitations run. In common with the spleen it has some attachment to the omentum and dorsally it is connected with the main dorsal mesentery. One long branch of it lies between the principal longitudinal folds of the duodenum. The main pancreatic duct, joined with the common bile duct, enters the duodenum some 10 mm. below the pylorus (Pl. VII, Fig. 2). The two ducts meet nearly at right angles a very short distance from or practically at the surface of the duodenum. Thence through the membranes of the duodenum there is an appreciable common duct carrying both biliary and pancreatic products to the common opening on the inner surface of the duodenum. At a point nearly opposite the union of the main ducts, the pancreatic duct receives an important branch or accessory duct derived from that lobe of the pancreas which lies along the duodenal fold.

Salivary glands.—The parotid gland (Pl. IV, Fig. 2) is extensive and invests the base of the external ear except in its dorsal third. It covers a part of the trapezius muscle, the sternomastoid and all the superficial muscles of the side of the neck. Three lobules are distinguishable, one anterior and two posterior, the dorsal of these latter being the larger. In a male specimen, the length of the gland is 16 mm. and its greatest dorso-ventral depth 10 mm. Its duct leads forward crossing the masseter superficially and then, sinking slightly, opens into the mouth near the first upper molar tooth.

The submaxillary gland also is large and is intimately associated with the parotid. It lies in ventro-lateral position relative to the parotid and has a detached lobe lying between its principal longitudinal lobe and the parotid. Its duct was not traced with certainty.

A large sublingual gland lies along the inside of the mandibular

ramus. Posteriorly it lies between the mylohyoid and the styloglossus muscles and anteriorly it is between the mandible and the genioglossus. It extends from the vicinity of the base of the angular process to the symphysis. Its ducts were not sought but it is evidently independent of the other large salivary glands.

Thyroid.—In a male specimen, the thyroid was found in two elongate pisiform divisions (Pl. VI, Fig. 1) lying on the lateral surface of the trachea. The larger anterior one extended from the vicinity of the thyroid cartilage caudad about 4 mm. The smaller lay just caudad in the same longitudinal plane and was about one-third the size of its fellow. In a female specimen, only one thyroid was detected.

Thymus.—All the specimens examined were fully adult and show only traces of the thoracic thymus. No indication could be found of the presence of the superficial cervical thymus regarded as characteristic of the Diprotodontia by Symington (1898) and Johnstone (1901). Four distinct fatty lobes lying ventrad of the carotids and cephalo-ventrad of the base of the heart were noted in one specimen as probably representing the thymus.

Mammæ.—There are four mammæ, regularly arranged, two on each side of the abdomen inside the thighs. This is the number present in the majority of diprotodonts and less than the number usually found in the polyprotodonts. In comparison with the *Didelphiidae*, however, the number is perhaps less important than the regularity of arrangement, for members of that family are distinguished from all other marsupials by having the mammæ present only in odd numbers and more or less asymmetrically arranged. The number in diprotodonts never exceeds four; in polyprotodonts it is usually more than six, although *Thylacinus*, *Myrmecobius* and a few species of *Phascologale* have only four, while the greatly specialized *Notoryctes* has but two. The following are the number found in the principal marsupial genera:

POLYPROTODONTIA:

Peragale	8
Perameles	6-8
Chaeropus	8
Thylacinus	4
Sarcophilus	4
Dasyurus	6-8
Phascologale	4-10
Sminthopsis	8-10
Myrmecobius	4
Notoryctes	2
Didelphis	5-13
Metachirus	7-9
Lutreolina	9
Philander	7
Marmosa	9-15
Peramys	9-27

DIPROTODONTIA:

Macropus	4
Acrobates	2
Distechurus	4
Dromicia	4
Petaurus	4
Pseudochirus	4
Phalanger	4
Phascolarctos	2
Phascalomys	2
Cænolestes	4

Miscellaneous glands.—The smaller sebaceous and lymphatic glands were not systematically sought but a few have been noted in the course of dissecting other parts.

Three rather conspicuous axillary glands are found imbedded in a fatty mass lying laterad of the *cutaneus maximus* and partially enclosing the elbow. The largest of these is long ovate pyriform and lies dorso-laterad of the cutaneous muscle, projecting also over the *latissimus*. It is about 4 mm. long and 2 mm. wide. Two others, slightly smaller, lie together ventro-laterad of the *cutaneus maximus*.

Two large superficial inguinal lymphatic glands in the male are situated in the deposit of fat ectad of the cremaster muscle. In one specimen, the larger of these glands measured 7 mm. in length and the other 5 mm.

A rather prominent precardial lymphatic gland is situated in the acute angle between the base of the oesophagus and the fundus of the stomach (Pl. VII, Fig. 4). In one male, this gland measured 4 mm. in length, being approximately equal to the diameter of the oesophagus.

An ovoid lymphatic gland about 3 mm. in greatest diameter is situated in the angle between the sternomastoid, omohyoid, and digastric muscles (Pl. IV, Fig. 1). It lies close to the trachea and is separated from it only by the carotid blood vessels.

In the male, a large paired lymphatic gland (Pl. VIII, Fig. 2) lies laterad of the large Cowper's gland (No. 3). It is about 6 mm. long by 3.5 mm. in width. Its wider end rests in the angle at the base of the penis pouch between the base of the penis and the bulb of the corpus spongiosum. Its smaller end lies in connective tissue and fat on the side of the large Cowper's gland. It is not directly connected with the rectum and has no duct that is apparent.

In the female, there is a large ovate pyriform rectal gland lying on each side of the cloaca with its small end directed dorsally and a duct leading from near its large end to the ventro-lateral wall of the cloaca slightly caudad of the opening of the rectum. The gland is 4 mm. long and has a tough thickened wall somewhat differentiated from the glandular mass within.

RESPIRATORY SYSTEM.

Plate X, Fig. 2.

Lungs.—The left lung is entire. In two specimens, the right lung has two lobes besides the azygos and the anterior of these is slightly notched at a point corresponding to that where a complete division is found in the didelphids. In a third specimen, the right lung was found to be trilobate although the anterior division was relatively small. The

evidence is therefore inconclusive as to whether the normal arrangement in *Cænolestes* is bilobate or trilobate. In most marsupials, except the wombat and some of the Macropodidae, it is said to be trilobate. In several specimens which I have examined of *Didelphis*, *Marmosa*, and *Peramys* it is trilobate. In *Cænolestes*, there is at least a tendency for it to be bilobate.

Trachea.—The tracheal rings are about nineteen in number and are not united dorsally but separated by a broad interspace.

Larynx.—The epiglottis is large and elevated, and has a slight anterior emargination which is most evident in the cartilage after removal of the soft external membranes. It is attached to the thyroid cartilage ventrally. The thyroid is continuous medio-ventrally with the cricoid and its bony wings partially inclose the cricoid laterally. Its anterior or superior horn has ligamentous attachment to the thyrohyal; its posterior or inferior horn attaches similarly to the outer extremity of the cricoid which is here very slightly ossified. The arytenoids are large, subtriangular and partially united dorsally. The cartilage of Santorini is developed into a prominent curved protuberance which is continuous with the high muscular crest of the arytenoid which, meeting its fellow of the opposite side, nearly closes the cavity of the larynx with the exception of a very small passage on the ventral side.

CIRCULATORY SYSTEM.

Owing to the condition of the material, no attempt was made to study the circulatory system in detail. The heart is as usual in marsupials. In comparison with that of *Didelphis* it shows no very important characters. In ventral view the auricles meet over the base of the pulmonary artery, entirely concealing it, whereas in *Didelphis* it is broadly exposed. The bases of the right subclavian and right carotid are very closely approximated and both join the left carotid scarcely higher than its junction with the aorta. In *Didelphis* there is an appreciable space between the origins of the right subclavian and right carotid.

SKELETON.

Plates XI-XVI.

CERVICAL VERTEBRAE.

Plate XI.

The seven cervical vertebrae are freely movable and distinct. As in the majority of marsupials, the only one having a prominent spinous process is the axis. The neural arches have relatively thin, flattened

laminae which are weak as compared to the bodies. They are well separated from each other and have no tendency to imbrication. All except the first and seventh have vertebrarterial foramina. Only the first and second have hypapophyses.

The atlas is practically entire, having a body which evidently ossifies in normal manner and having no large foramina for the passage of nerves and blood vessels. A very minute perforation (Pl. XI, fig. 1, *n. f.*) at the posterior dorsal base of the transverse process is of doubtful significance, but its position does not seem to indicate homology with either of the usual foramina of the atlas. A perforation with exactly similar relations is found in the extinct *Borhyaena* (Sinclair, 1906, p. 350, Pl. LIII). The cephalic border of the neural arch of the atlas is nearly straight for about the width of the neural canal and thence on either side it is angled latero-caudad to meet the base of the cephalic articular process. A groove between this angle of the border of the arch and the articular process serves for the passage of the first cranial nerve and supplies the office of the foramen found in most other marsupials. That this foramen is formed by the closing of the points bounding the mouth of this notch is shown by immature examples of *Didelphis* in which the union is slightly cartilaginous. In the adult of *Didelphis*, the foramen is well isolated as it is in *Philander*, *Dasyurus*, *Myrmecobius*, *Thylacinus*, *Sarcophilus*, *Phascologale*, *Trichosurus*, *Petaurus*, *Petrogale*, *Macropus*, *Phascolomys*, and *Phascolarctos*. The only living marsupial genera agreeing with *Cænolestes* in this open "intervertebral notch" are *Marmosa*, *Peramys*, and *Perameles*. The atlas of *Perameles* is otherwise similar to that of *Cænolestes*, but that in *Marmosa* and *Peramys* resembles it still more closely, differing chiefly in the pedunculation of the transverse process and its slight obliquity and in the tendency to the formation of a vertebrarterial canal with a partly cartilaginous border.

The transverse process of the atlas has decided obliquity to the spinal axis and is relatively short and broad with no constriction at its base and with only a shallow groove between it and the posterior zygapophysis in the position corresponding to the usual course of the vertebrarterial canal. The only genera examined in which the arterial canal has complete bony boundaries are *Philander*, *Dasyurus*, *Sarcophilus*, *Myrmecobius*, and *Petaurus*. In all others except *Cænolestes*, however, the transverse process is pedunculate and in some, as *Phascologale* and *Peramys* (*Marmosa* and *Didelphis* to a lesser extent), there is a slight bony elevation on the pedicel of the arch cephalad or laterad of the postzygapophysis from which a cartilage extends to the posterior angle of the transverse process and forms the arterial canal. So far as material at hand shows, the gradual formation of the vertebrarterial canal of the

atlas is illustrated in various stages from the wide open channel to the heavy bony ring in a series of forms the sequence of which would be about as follows: *Cænolestes*, *Perameles*, *Didelphis*, *Phascologale*, *Peramys*, *Dasyurus*, *Philander*, *Petaurus*, *Myrmecobius*. In this series, *Didelphis* is about midway between the extremes and various other forms, including the macropods and phalangers, would occupy a similar position.

The atlas in *Cænolestes* has no trace of a spinous process. The body of the atlas is directed downward and slightly forward, being almost vertical and nearly at right angles to the laminae of the arch. Its lower part is thus a thin triangular plate with a hypapophysis or median ridge on its posterior aspect and forming a pointed tubercle at its apex. The odontoid process, therefore, does not have its usual flattened seat but rests on the sharp edge of this nearly vertical body of the atlas. This part of the atlas differs from that of practically all other marsupials, although it is closely approached in *Phascologale*, *Marmosa*, and *Peramys*. The neural canal of the atlas is slightly wider than deep and measures 4 mm. x 3.4 mm.

The imperforate condition of the atlas in *Cænolestes*, both anteriorly and posteriorly, is doubtless primitive. It is found in the monotremes and among living marsupials examined only in *Marmosa*, *Peramys*, and *Perameles*.¹ It is characteristic also of the extinct *Borhyaena* but, so far as known, of none of the other forms of the Santa Cruz beds. The atlas, however, is unknown in all the fossil forms placed by Sinclair (1906) in the family *Cænolestidae*. In other mammals, including all the higher groups, no instances have been found in which both the atlanteal and vertebrarterial foramina are absent.

The axis has a high laminate spinous process similar in general to that of the majority of marsupials. The anterior wing of this process is not equal to the posterior and reaches only to the posterior part of the dorsal surface of the atlas. The posterior wing overhangs the third cervical. The front border of the neural arch seen from the side is a continuous curve, and the prominent dentation found in *Didelphis*, *Peramys*, *Marmosa*, *Philander*, *Trichosurus*, *Petaurus*, and some others is absent. At the base of this curve is the prezygapophysis which is an oval facet only very slightly elevated from the pedicel of the arch. The odontoid process is completely ankylosed in the adult and forms a simple projecting knob of the usual character. The body of the vertebra is a fairly thick shield-shaped piece with two relatively large foramina in the floor of the neural canal. The hypapophysis is peculiar. It is

¹In *Notoryctes* the atlas is imperforate, but this is obviously due to extreme specialization and has no significance in this connection. (See Stirling, 1891.)

divided into two parts one of which probably appertains to the odontoid and the other is the usual hypapophysis of the axis. The anterior of these is a sharp hooked process rising from the base of the odontoid and projecting backward along the median line. The posterior process is a ridge rising beneath this hook and gradually becoming more elevated to terminate in a tubercle which has slight projection caudad beyond the centrum and thus to some extent effects imbrication with the front edge of the body of the third vertebra. In *Marmosa* there is a prominent axial hypapophysis and in the skeleton of a half-grown *Didelphis* with the odontoid still distinct there is a small median knob but in adults I find nothing in any marsupial comparable to the hooked process described above. It very strongly suggests the large recurved hypapophyses of *Ornithorhynchus*. The posterior border of the neural arch in lateral view presents an even curve from the spine to a prominent angle above the postzygapophysis. The neural canal has the outline of a horseshoe and its width and depth are nearly equal.

The transverse process of the axis has a superior and an inferior division, each a small backwardly directed projection from the caudal aspect of the pedicel of the arch, ventrad of the postzygapophysis and caudad of the prezygapophysis. The extremities of these processes are articulated to the slightly produced basal angles of a small, distinct, subtriangular pleurapophysis or cervical rib which serves to complete the boundary of the arterial canal and which is obviously the homologue of the fully ankylosed processes serving the same purpose in the succeeding vertebrae. In a fully adult animal this cervical rib has only cartilaginous union with the vertebra and in the undesiccated specimen it is freely movable (Pl. XI, fig. 7). It is essentially similar to the persistently distinct cervical ribs in *Ornithorhynchus* and in reptiles. It has not been reported among marsupials or higher groups except in *Perameles* and *Phascologale*. In *Perameles lagotis*, it is noted by Owen (Trans. Zool. Soc. Lond., II, p. 394, 1841) in a single instance. The only skeleton of *Perameles* (*P. nasuta*) I have been able to examine, has the axial pleurapophysis fully ankylosed. Whether it is normally so in this genus, therefore, is uncertain; but its persistence as a distinct part in *Cænolestes* is none the less significant. Beddard (1902, p. 119) mentions a separate axial rib for *Phascologale*.

The third, fourth and fifth cervicals are not peculiar and resemble each other in general characters. In the third the neural canal is relatively deeper, its vertical and transverse diameters being nearly equal, whereas in the fourth and fifth the width considerably exceeds the depth. In the third the spinous process is very slight; in the fourth and fifth it is a low but distinct mammillate process. In all three the inferior

branch of the transverse process is only slightly developed. The superior branch of the transverse process is directed obliquely backward and upward and at its base is pierced by the arterial canal. In the fifth it is slightly thicker than in the third and fourth, has less backward deflection and is definitely faceted at the tip.

The sixth cervical is similar in general features to that in the majority of marsupials. The arterial canal perforates the base of the transverse process as usual but it is to be noted that the dorsal wall of the canal is very much thicker than the ventral. The inferior branch of the transverse process is expanded into a dolabriform plate practically parallel with the spinal axis. This has only very slight forward projection but is extended caudad nearly or quite to the plane of the centrum of the first thoracic vertebra.

The seventh cervical has a slightly higher spinous process than the three preceding vertebrae, but the distinction is not marked. It has the superior branch (diapophysis) of the transverse process with its tubercle directed almost at right angles to the spinal axis in the horizontal plane of the body of the vertebra. The inferior branch (parapophysis) and consequently the arterial canal are obsolescent, that is, they are variable, being present in some specimens and reduced or absent in others. In one specimen, the arterial canal on the left side of the vertebra is enclosed by the merest bony thread representing the parapophysis and having no tubercle. On the right side of the same vertebra, the condition is similar but the canal is exceedingly small, scarcely one-tenth as large as that on the left side. In several other specimens, the seventh cervical is quite imperforate, the vertebral side of the base of the transverse process being merely grooved (Pl. XI, figs. 4-6). In a single specimen of *Phascologale*, the lower boundary of the arterial canal is incomplete, indicating a condition very similar to that in *Cænolestes*; in a single specimen of *Sarcophilus*, one side is perforate and the other imperforate. The seventh cervical is imperforate in *Trichosurus*, *Dasyurus*, *Perameles*, *Sarcophilus*, *Didelphis*, *Marmosa*, and *Peramys*. The arterial canal has complete bony boundaries in *Thylacinus*, *Petaurus*, *Philander*, *Phascolomys*, *Phascolarctos*, and the Macropodidae. It is complete also in the extinct genera *Borhyaena* and *Cladosictis*. Thus in the majority of diprotodonts the seventh cervical is perforated and in the majority of polyprotodonts it is imperforate, although the perforate condition is more primitive. In this as in other respects, therefore, *Cænolestes* combines the characters of the two groups.

None of the cervicals have foramina piercing the lateral walls of the neural arches like those in some other marsupials (*Thylacinus*, *Dasyurus*, *Didelphis*, etc.).

Considered as a series, the cervical vertebrae of *Cænolestes* seem more similar to those of *Perameles* than of any other marsupial. The imperforate atlas, the persistent cervical rib attached to the axis, the evenly curved front edge of the neural canal of the axis, and the imperforate seventh vertebra all are found in *Perameles*. The inferior lamellae of the transverse processes of the third, fourth, and fifth cervicals are much larger in *Perameles* than in *Cænolestes* and this constitutes the principal distinction between them. In comparison with those of other American forms, the cervicals of *Cænolestes* show a general resemblance to those of *Marmosa* and *Peramys* and, among Australian small polyprotodonts, to those of *Phascologale*, but the characters of the atlas and axis are distinctive. With the highly modified cervicals of *Didelphis*, comparison is scarcely necessary.

THORACIC VERTEBRAE.

There are thirteen thoracic vertebrae as in all other marsupials with the exception of *Phascolarctos* (11), certain species of *Phascolumys* (15), and *Notoryctes* (15).¹ The spine of the first thoracic is scarcely higher than that of the last cervical (Pl. XII, fig. 4). The second spine is slender and much the highest of the series, measuring about 3.5 mm. The third to the eighth also are slender and progressively shorter and more recumbent. The ninth is still shorter, and although directed backward, it is longer basally and less acicular in form. The first thoracic differs from the last cervical mainly in the elevation of the transverse process and in its differentiation for articulation with the rib. In most marsupials there is an abrupt transition from the last cervical to the first thoracic, especially in the length of the spinous process, but in *Cænolestes* this break is between the first and second thoracics. The only form examined which agrees with it in this respect is *Perameles nasuta*. In *Phascolarctos*, the spine of the first thoracic only slightly exceeds that of the last cervical but it is nevertheless the longest of the thoracic spines and there is nowhere any marked break in the series. Owen (1841) in his paper on the osteology of marsupials does not mention a relatively short spine for the first thoracic of *Perameles* and it is therefore possible that it is not found in all species of the genus.

The tenth is the anticlinal vertebra and its spinous process is a low upright plate with a straight crest occupying practically the entire length

¹ In a single specimen of *Didelphis paraguayensis* from Ceará, Brazil, there are fourteen ribs instead of the thirteen normal in *Didelphis*. Moreover, there are twenty vertebrae in the thoracic-lumbar series, the extra vertebra apparently being the fourteenth, corresponding to the extra rib. This is probably an abnormality, but the examination of other specimens of this species is desirable, especially since, as noted elsewhere (p. 86), it has other skeletal peculiarities. A pouch specimen of *D. p. audinus* has the normal thirteen ribs.

of the dorsal surface of the arch. The eleventh, twelfth and thirteenth spines are similar, but although only slightly elevated, they are definitely directed forward instead of backward. The thirteenth is slightly lower than the eleventh and twelfth and closely resembles the spines of the succeeding lumbar.

The position of the so-called center of motion sharply distinguishes *Cænolestes* from all living American polyprotodonts and aligns it with the majority of Australian forms, as well as with most higher mammals. In *Didelphis*, *Marmosa*, *Philander*, and *Peramys* there is practically no anticline since the spines of all the thoracic vertebrae and of all the lumbar, except the last or at most the last two, are inclined backward. The only other marsupials having this characteristic are the koala, the wombats, and certain of the phalangers. In all the Australian polyprotodonts and at least in the macropods of the diprotodonts the center of motion is in the posterior thoracic vertebrae and in most of these forms the anticline is very marked.¹ This is also the case in *Cænolestes* and in the Patagonian fossil form *Cladosictis*. Whether the condition in the didelphids is palaeotelic or not is problematical. Essentially the same condition is found in the monotremes and edentates, and it is usual among reptiles, whereas among higher mammals it is rare and generally correlated, perhaps secondarily, with high specialization, as in seals and anthropoids. In practically all insectivores, rodents, bats, carnivores, ungulates, and in Primates, except the higher apes and man, there is a marked convergence of vertebral spines to a center of motion. In agreeing with the majority of mammals in this respect, therefore, *Cænolestes* at least shows a lack of affinity with the didelphids.

The first nine thoracic vertebrae are essentially similar in form. The arches and pedicels, although progressively increasing in length, are relatively narrow and each vertebra is separated from the succeeding one dorso-medially by a considerable space between the projecting zygapophyses. Each has a transverse process bearing a facet for the tubercle of a rib. The tenth begins to assume the character of the lumbar. It has only a trace of a facet for the tubercle of a rib, its spinous process is low and ridgelike instead of spinous, its laminae and pedicel are broad and its body is less compressed. The eleventh has these characters further developed. There is no trace of a lateral rib attachment and posteriorly there is a tiny pointed accessory process below the postzygapophysis. In the twelfth and thirteenth this accessory process is slightly larger and in these the prezygapophyses begin to turn upward, but there is only a slight rudimentary mammillary process.

¹ In the single skeleton of *Sarcophilus* examined there is a double anticline, one at the eleventh thoracic and another at the third lumbar.

In the character of the first thoracic and essentially in other respects, the thoracic series of vertebrae, like the cervical, resembles that of *Perameles* more closely than that of any other marsupial. It differs chiefly in the last three thoracics which in *Perameles* have relatively high spines and prominent mammillary processes.

LUMBAR VERTEBRAE.

Plate XIII, Figs. 1-2.

The six lumbar have no important peculiarities. The first three increase slightly in size caudad but aside from the appearance of small inferior anterior transverse processes they scarcely differ from the last thoracic. There are still small accessory processes (anapophyses), the spinous processes are low and ridgelike and the mammillary processes (metapophyses) scarcely protrude beyond the articular surfaces of the zygapophyses. The fourth lumbar retains the ridgelike spinous process, but the transverse process, directed downward and forward, is considerably increased in size. The fifth and sixth lumbar are somewhat differentiated from the others. The transverse processes are longer, the spinous processes, which are directed forward, are higher and more hooklike, and the postzygapophyses are nearer together and turned slightly upward. There are no accessory processes but the mammillary processes are more prominent than in preceding vertebrae and the laminae of the arch are compressed posteriorly for a narrowing neural canal. The ventral surfaces of the bodies of the lumbar are slightly differentiated posteriorly into two slight ridges or tubercles for the psoas muscles. A single median ridge or a long hypapophysis is found in various forms, in *Phascologale*, *Antechinomys*, *Perameles* and to some degree in *Marmosa*. Double ridges on the posterior half of the body of the vertebra similar to those in *Cænolestes* have been observed in *Didelphis*, *Dasyurus*, and *Petaurus*.

The lumbar series as a whole may perhaps be said to be characterized by the low and inconspicuous condition of the vertebral processes which in most other forms are more highly developed. In this respect there is some resemblance to immature specimens of *Didelphis*, but even in these the anterior and posterior zygapophyses are connected by a sharp ridge and the accessory processes are broad, heavy, and blunt instead of small and pointed.

SACRAL AND CAUDAL VERTEBRAE.

Plate XIV.

Two vertebrae have been regarded as sacral, although the second touches the ileum only at the anterior extremity of its transverse process.

The first has a much broader and heavier transverse process, the whole lateral surface of which is firmly attached to the ileum. The condition is much like that in *Phascologale*, *Sminthopsis*, *Philander*, and *Myrmecobius*. A slightly greater proportion of the second sacral touches the ileum in *Marmosa*, *Peramys*, *Dasyurus* and *Perameles*; a still greater in *Didelphis*, *Phascolarctos* and *Phascalomys*; and in *Trichosurus*, *Petaurus*, *Petrogale* and *Macropus* the second vertebra is almost or quite as fully attached as the first; while in a single specimen of *Thylacinus*, the second sacral, although heavily ankylosed to the first, does not reach the ileum at all. The two sacrals are ankylosed to each other laterally, but a large perforation persists between their transverse processes. Their general character corresponds closely with that of the last lumbar, the chief differences being in their widened and more horizontal transverse processes and in the greatly compressed zygapophyses, both anterior and posterior in the case of the second, and mainly the postzygapophysis in the case of the first. The first has a well-developed and forwardly directed spinous process similar to that of the last lumbar. The second has a smaller, more pointed spinous process directed slightly backward.

The first caudal and sometimes the second is ankylosed to its predecessor by a narrow bony strip stretching between the extremities of the transverse processes. The first and second are essentially similar to the sacrals except that they have no traces of spinous processes and that the zygapophyses are expanded slightly in the first and greatly in the second, with short rounded metapophyses. The third and fourth caudals are similar to the second but their transverse processes are inclined forward and the first chevron bone is found over their intervertebral space. The fifth caudal is the first to lack the neural canal and postzygapophyses, but the transverse process is still undivided, although somewhat reduced in lateral extent and lengthened antero-posteriorly. In some specimens this transverse process is distinctly emarginate laterally. The sixth caudal is elongate and subcylindrical with rather prominent metapophyses and short crescentic anterior and posterior transverse processes. The succeeding vertebrae are almost exactly similar up to the sixteenth which has only the merest traces of processes. The remainder taper gradually to a very slender tip, the total number of caudals being twenty-seven. The ninth and tenth are the longest of the caudals each measuring about 7.7 mm. Nineteen chevron bones are present from the third to the twenty-second caudals. These have one to four longitudinal ridges on their under surfaces which give them a fluted appearance. They are nearly as wide as the bodies of the vertebrae.

The characters of the caudals are of course of little or no significance in the indication of relationships, but it is of some interest to note that

the absence of spinous processes on the proximal caudals is a more frequent condition among Australian than American forms. With the exception of *Perameles*, which has six caudal spines, and *Phascolarctos* and *Phascolomys*, with practically complete sets, all the Australian forms examined either have no caudal spines or have them few and extremely rudimentary. In no form are they so completely absent as in *Cænolestes*. On the other hand, there are no less than four in *Marmosa*, three in *Peramys*, and three to six in *Didelphis*.¹ Slight caudal spines are reported in *Notoryctes* (Stirling 1891).

The angle at which the sacrals are set between the ilia is less in *Cænolestes* than in most of the polyprotodonts examined and, as in *Trichosurus* and *Macropus*, the spinal axis is nearly parallel with the longitudinal axis of the superior outline of the pelvis. Thus, if the caudals are extended in the axis of the sacrals the transverse processes of the fourth and fifth caudals lie below the ischial tuberosity instead of well above it as in *Didelphis*, *Dasyurus*, *Perameles*, *Myrmecobius*, etc.

RIBS AND STERNUM.

Plate XII.

There are thirteen pairs of ribs as usual, seven pairs articulating directly with the sternum and six pairs not reaching it. Nine pairs are doubly articulated with the vertebrae by tubercles and capitulae. The succeeding four pairs lack tubercles and arise from the bodies of the vertebrae. The first rib is much heavier than the others and its costal element is longer than the vertebral which is true of no other marsupials examined except *Trichosurus* and *Petaurus*. Its costal element is fully ossified, basally flattened and slightly bicapital near its union with the lateral extension of the manubrium. The vertebral element is similarly expanded at the head and narrowed toward its union with its fellow. A strong first thoracic rib, it may be noted, is an edentate character. The remaining ribs from the second to the thirteenth are very slender and subcylindrical, especially the vertebral series. The sternal series are slightly flattened, the seventh being rather noticeably wider than the others. The sternal ribs from the first to the seventh show a greater degree of ossification than usual in marsupials. The seventh pair of costals articulate directly with the caudal end of the last segment of the mesosternum and therefore lie ventrad of the xiphisternum and nearly or quite touch each other at this point (Pl. XII, fig. 2). In *Didelphis* and most other polyprotodonts these ribs only reach the sides of the meso-

¹Although six caudal spines are distinguishable in *Didelphis virginiana*, a specimen of *D. paraguayensis* from Ceará, Brazil, has only traces of three and the spine of the second sacral also is undeveloped.

sternum and are separated at their extremities by the xiphisternum, whereas in *Perameles* and most diprotodonts the condition is as in *Cænolestes*.

The manubrium of the sternum is like that of most of the smaller clavicate marsupials. It consists of an anterior prong which is somewhat deflected ventrad and slightly faceted dorso-cephalad for the clavicular union. A small triangular lateral process projects from each side of the base of this prong and the body of the bone extends thence caudad as a subcylindrical rod with a slight keel on its ventral surface. In a male specimen the manubrium is 7 mm. in length.

The four segments of the mesosternum are of nearly uniform width throughout without pronounced median constriction or lateral expansion even in the fourth which is scarcely wider posteriorly than anteriorly. The first is somewhat more rounded than the others which are slightly flattened but with their ventral surfaces rather more convex than otherwise.

The xiphisternum is a long slender bony rod with slightly greater width than thickness, but, in general, rounded in form especially proximally where it is somewhat expanded. Distally it has a more abrupt and more flattened slight expansion where it joins the xiphoid cartilage, which is relatively large and scutate. The xiphisternum is separated from the last segment of the mesosternum dorsally by a broad epiphysial cartilage and ventrally by the extremities of the seventh pair of ribs. Hence, although the plane of the dorsal surface of the xiphisternum is continuous with that of the mesosternum, its ventral plane is distinctly higher. This epiphysial cartilage and this relation of the proximal end of the xiphisternum, although apparently trivial, are nevertheless characteristic of the diprotodonts as opposed to the polyprotodonts. In all polyprotodonts examined, except *Perameles*, both surfaces of the xiphisternum are practically continuous with the corresponding surfaces of the mesosternum and the sixth or seventh ribs do not meet ventrad of the base of the xiphisternum. In all diprotodonts, including *Cænolestes*, the reverse is the case. *Perameles*, as judged by one specimen, seems to occupy an intermediate position in this respect, for although ribs meet on the end of the mesosternum, the epiphysial cartilage is so reduced as to be scarcely noticeable in a dried specimen.

PECTORAL GIRDLE.

The clavicle (Pl. XII, fig. 5) is well developed and stretches from the sternum to the acromion in the manner usual among marsupials. Its lateral or acromial end is compressed and curved upward. In the adult, the union with the acromion is almost direct, the intervening cartilage being scarcely evident. It is attached also to the coracoid process by a

broad flat coraco-clavicular ligament. The medial or sternal end of the clavicle is somewhat expanded and deeply notched for the short stout omosternal cartilage¹ by which it is attached to the front of the manubrium.

The scapula (Pl. XIII, fig. 3) is less angular than in the majority of marsupials, having a slightly elongate, gently curving outline without sharp prominences. The suprascapular notch, so-called, is not a notch but a long sweeping curve which does not reach its highest point until about two-thirds of the distance from the coracoid process to the vertebral border of the scapula. The axillary border has a slight convexity, instead of being practically straight, and it meets the vertebral border in a modified curve which in some specimens may be a trifle angular. The slightly thickened vertebral border is covered by a small, elongated, cartilaginous, or sometimes partly osseous, suprascapular epiphysis.²

The spine is relatively low and continuously deflected so that it is nowhere perpendicular to the fossae on either side of it. Thus the antero-ventral triangular half of the infraspinous fossa is overhung by the flat deflected surface of the spine, which continues to the well-developed acromion, turning sharply forward beyond the triangular metacromion. The coracoid process is a prominent prong with a slight notch in its glenoid surface. The superior fossae are nearly equal in extent, the supraspinous usually having slightly the greater surface. The inferior surface or subscapular fossa presents a deep longitudinal channel opposite the base of the spine and on either side of this the surface is somewhat bulging.

Taken as a whole the scapula of *Cenolestes* is more similar to that of *Sminthopsis* than to that of any other form which is available for comparison. It agrees with this in its general outline and in its much deflected spine, whereas practically all other forms have a more angular shape, a less deflected and usually higher spine, and a shorter more abrupt suprascapular notch.

PELVIC GIRDLE.

Plate XIII, Fig. 4.

The pelvis is completely ossified and shows no traces of division in adult specimens. It is relatively long and deep but light in structure. Although it is not especially peculiar in other respects, it is strikingly distinguished from that of all other marsupials by its extremely short

¹ In a skeleton of *Philander*, this cartilage is partially ossified.

² A distinct bony suprascapular epiphysis is especially evident in *Perameles* and *Thylacinus*.

pubic symphysis. This scarcely equals the diameter of the acetabulum. The general proportions of the pubes and ischia are thus essentially similar to those in various edentates and insectivores. The rami of the pubis and ischium together with the body of the ischium form a nearly equilateral triangle having its angles respectively at the ischial tuberosity, the pubic symphysis, and the cephalic border of the acetabulum. The pubic ramus is very slender and is slightly ridged on its lateral surface. The ischial ramus is slightly wider and more compressed and the body of the ischium is a trifle more so, but all these bones are relatively slender and the obturator foramen which they enclose is large and subtriangular. The ischial rami are not greatly divergent but more nearly approach the perpendicular than those of any other marsupial and in this respect show resemblance to certain insectivores. The ischial tuberosity is fairly well developed and in some specimens forms a decided protuberance.

The ilium is slender and of the ordinary prismatic shape, being somewhat flattened and everted at the tip. Its ischial border forms a high ridge continuous with the inner edge of the ischium and to some extent overhanging the anterior part of the obturator foramen in shelf-like manner. A slight iliopectineal tubercle is present. The acetabular border of the ilium forms a fairly well marked ridge, especially anteriorly where it separates the gluteal and iliac surfaces of the ilium. The acetabulum and cotyloid notch are of the usual character.

The marsupial bones are well developed, their unusually long straight bases being attached to almost the entire anterior edge of the long ascending pubic ramus, that is, from the symphysis to a point directly below the acetabulum and caudad of the iliopectineal tubercle. The base of the marsupial bone is thus almost as long as the anterior projecting spine and proportionally longer than in any other marsupial. In a male specimen, the marsupial bone measures 5.6 mm. in length as against a base of 4.8 mm., the ratio of base to length being 85.7. This ratio is approached only in a few forms in which the marsupial bones are notably rudimentary. The following ratios indicate how widely *Cænolestes* differs from the majority of marsupials in this respect: *Cænolestes* 85.7; *Didelphis*, 40; *Dasyurus*, 34; *Myrmecobius*, 15; *Sminthopsis*, 52; *Perameles*, 14.2; *Trichosurus*, 41; *Phascolarctos*, 37.3; *Petrogale*, 9.

ARM AND FOREARM.

Plate XV.

In proportion to the size of the animal, the humerus is large and heavy, being much the largest single bone in the body. It differs conspicuously from that of most marsupials in its great degree of curvature

and in its prominent flat-faced deltoid ridge. This occupies about three-eighths of the inner proximal part of the front of the humerus, and instead of forming a single trenchant ridge it presents an elevated flat rhomboidal surface extending downward from the greater tuberosity and ending bluntly and abruptly. Its surface is slightly roughened and lies at a considerable angle to the long axis of the humerus. From its distal extremity the front of the shaft abruptly recedes and forms a long sweeping curve to the condyles. The deltoid ridge or tuberosity has a flattened surface in some other forms, as *Perameles*, *Dasyurus*, and *Peramys*, but in these it is triangular, coming to a point distally. In the fossil form *Garzonia patagonica*, of which a humerus is figured by Sinclair (1906, Pl. XLIII, fig. 11), the deltoid surface is very broad, and although it is less blunt distally, it seems to approach the condition in *Cænolestes* more nearly than does that of any recent form.

The head of the humerus is rather large and has considerable backward extension beyond the axis of the shaft. The external or so-called greater tuberosity is much smaller and less marked than the internal. It is greatly compressed and its outer surface is continuous with the deltoid ridge. Its upper surface is continuous medially with the surface of the head of the humerus. The internal tuberosity is prominent and distinctly separated from the head by a deep groove as well as from the external tuberosity by the bicipital groove. A well-marked ridge extends distad from it parallel with the internal border of the deltoid ridge and terminating in a roughened area for the insertion of the latissimus muscle.

The middle of the shaft of the humerus for about one-third the length of the bone is relatively smooth, slightly compressed, subcylindrical, with only traces distally of the upper continuation of the external epicondylar or supinator ridge. The caudal outline from the head of the humerus to the supinator ridge is a deep sigmoid curve. The distal end of the humerus is broadly and rather abruptly expanded. The internal condyle is prominent and the sulcus between it and the ulnar trochlea is developed as an articular surface for the inner upper lip of the sigmoid cavity of the very obliquely situated ulna. The trochlea is somewhat smaller than the capitellum with which it is continuous. Thus there are three articular surfaces at the distal end of the humerus, a very unusual condition. The third or internal articular surface is fully developed only on the caudal aspect of the bone and does not continue to the front as the trochlea and capitellum do. Although partly confluent, these three articular surfaces are sufficiently distinct from each other, but of course the third or inner one could be regarded as an extension of the trochlea which might be developed without difficulty.

Nevertheless, I have been unable to find a case of a similar extension in any other marsupial or in other mammals. Its effect is a much wider articulation with the ulna than usual and, as described below, there is a corresponding increase of the articular surfaces of the ulna. Functionally it may have some correlation with the distinctness of the median head of the triceps muscle, and although it might not be remarkable in an animal of fossorial habits, it seems quite incompatible with the distal elements of the arm and leg which are adapted to cursorial life.

A small oval epicondylar foramen is present and the ridge bounding it has a smooth rounded surface directed with a spiral curvature toward the distal end of the deltoid crest but widely separated from it by the smooth shaft. The outer epicondylar ridge is similarly twisted and gradually merges with the surface of the shaft without any abrupt angle or hooked process. Its edge distally is broadly curled so that it presents a relatively broad flattened lateral aspect instead of the sharp edge of most other marsupials. The external epicondyle is little more than the extremity of this ridge. It has a decidedly concave surface for the seat of the sesamoid of the tendon of the *supinator brevis* muscle. This sesamoid, as mentioned elsewhere, is not present in other marsupials. It is sufficiently distinct to be obvious to the preparator and is usually found in situ in first class ligamentary skeletons.

The ulna is broad, heavy and compressed proximally but rapidly tapers to its distal third which is subcylindrical and scarcely half the diameter of the radius. The outer side of its proximal half is deeply channelled. Its inner side is smooth and convex except an excavated area on the side of the olecranon. The upper lip of the greater sigmoid cavity is unusually broad, especially its inner division which extends well beyond the inner base of the olecranon from which it stands out prominently. Its inner division has the usual position opposite the lesser sigmoid cavity. This broad upper lip articulates with the whole of the posterior articular surface of the humerus and in doing so brings the ulna into a very oblique or nearly transverse position with its broad grooved outer side directed nearly forward. Except at its distal end the ulna is caudad of the radius. The olecranon is relatively thick and heavy with a rhomboidal roughened extremity for various muscle attachments.

The radius has a rounded ovoid head supported on a distinct neck below which is a prominent tubercle having a rounded roughened face quite two-thirds as large as that of the head. The proximal third of the radius is rather abruptly flexed forward leaving a considerable space between it and the ulna. For the remainder of its length it is in close

contact with the ulna, so close that more than a slight degree of rotation is not probable.

While the heavy humerus and the broad elbow joint of *Cænolestes* seem adapted for digging or scratching habits, the character of the forearm distinctly tends toward the cursorial type of structure. Aside from a few detailed peculiarities, it may be said that the humerus has considerable general resemblance to that of *Peramys* and *Marmosa* while the radius and ulna are clearly nearer those of *Antechinomys* and *Sminthopsis* or even *Perameles*.

BONES OF THE HAND.

Plate XVI, Fig. 1.

The carpal bones are eight in number, four in the distal row and four in the proximal row, including the pisiform. In the distal row the trapezium forms the seat of the first metacarpal and is in contact also with the inner proximal extremity of the second metacarpal. Its inner surface articulates with the trapezoid and its inner palmar surface with the distal projection of the scaphoid. It is a short longitudinally compressed bone with broad grooves on its distal and proximal surfaces respectively. No sesamoids could be detected in relation with it. The trapezoid and magnum are about equal in size. The former is rounded distally for articulation with the concavity of the proximal end of the second metacarpal. It articulates also with the trapezium, magnum, and scaphoid. The magnum is subtriangular in its anterior or dorsal aspect and articulates with the third metacarpal, the trapezoid, scaphoid and unciform. The unciform is large and roughly pentagonal in outline. It supports the fourth and fifth metacarpals. Its contact with the fifth however is limited to its radial side. The unciform articulates with five bones, viz., the two metacarpals, the magnum, lunar, and cuneiform. Its dorsal surface has two deep pits. The cuneiform receives the head of the ulna in its deeply concave proximal surface. Distally it articulates with the unciform and medially with the lunar. The lunar is quite distinct and presents a triangular dorsal surface about half as extensive as that of the magnum. With the scaphoid it supports the broad head of the radius. The scaphoid is nearly as large as the unciform. It lies between the head of the radius and the trapezoid and sends a curving process beneath the palmar surface of the trapezium. Its medial or ulnar surface is in contact with the lunar. No sesamoids were found associated with it. The pisiform is exceedingly well developed and broadly expanded distally. It is nearly as long as the fifth metacarpal and relatively longer than in any other marsupial which has been

examined. It articulates principally with the outer lateral surface of the cuneiform but also has slight contact with the ulna.

The metacarpals are slender and rounded with somewhat expanded heads and bases, the former slightly the larger. The medius is the largest. The indicis and annularis are slightly shorter and of about equal length. The pollicis is about three-fifths as long as the indicis and the minimus is scarcely more than half as long as the annularis. The first three are of approximately equal diameter, the annularis is very slightly more slender, and the minimus is slightly broader. The outer pairs are evenly and slightly divergent from the middle. The phalanges are regular in number and form. The basal ones of the second, third and fourth fingers are nearly equal in length; that of the fifth finger is slightly shorter, and that of the pollex is shortest. The claw of the middle finger slightly exceeds the others. The claw of the pollex reaches to the middle of the first phalanx of the index finger and the claw of the fifth finger extends to the first phalangeal joint of the fourth finger. The ungual phalanges of the first and fifth fingers are pointed but less curved and clawlike than those of the middle fingers. The usual paired sesamoids are present on the palmar sides of the metacarpo-phalangeal joints.

THIGH AND LEG.

Plate XII, Figs. 1, 3.

The femur is slightly shorter and decidedly more slender than the humerus. The shaft is cylindrical and nearly straight but the extremities are considerably expanded. The head is supported on a slight neck and is definitely directed forward as well as inward. Its smooth articular surface is partially incomplete behind. The outer or greater trochanter rises to the level of the top of the head and is inclined slightly inward so that its apex is practically in the long axis of the shaft. Laterally it is much compressed and it is only superficially separated from a broad ridge extending distad a short distance to a small rounded protuberance having a roughened surface and clearly representing a third trochanter. This has approximately the same relations as the third trochanter in *Phascolomys*, the only other marsupial except the extinct *Wynyardia* in which there is any obvious development of the kind. The great trochanter is bounded on the inner side by a deep slitlike digital fossa which has as its other boundary a high thickened intertrochanteric ridge almost as prominent as the great trochanter. This ridge runs from the apex of the great trochanter distad with slight inclination toward the lesser trochanter and then turning sharply and losing most of its eleva-

tion joins the lesser trochanter. A considerable depression lies between its lower half and the base of the head of the femur. In many marsupials, this ridge is only represented by a tubercle between the great trochanter and the head. It is fairly well developed in *Didelphis* and in *Perameles*. The lesser trochanter forms a prominent triangular projection on the inner side of the femur separated from the head by a short rounded notch.

The distal end of the femur is of the usual character. The condyles are produced backward to a marked degree and the intercondylar notch is narrow and deep. The anterior rotular groove is broad and shallow being scarcely more marked than in forms with a ligamentary patella. The patella itself is rather short and broad and well ossified. This is an important peculiarity of *Cænolestes*, for among living marsupials the only others having an osseous patella are *Tarsipes*, *Perameles* and *Notoryctes*. It is recorded also as present in the extinct genera *Prothylacynus* and *Amphiproviverra* (Sinclair).

The tibia is very long and much curved, resembling that of the macropods and certain small saltatorial polyprotodonts, as *Sminthopsis* and *Antechinomys*. Its proximal half is bowed inward and forward in a long curve and is widely separate from the fibula but sweeps downward into the same longitudinal axis for its distal half. The proximal half is of the usual trihedral form, its wide inner surface being longitudinally channelled and its antero-internal surfaces convex. The distal half is nearly cylindrical. The internal malleolus is highly developed, only being equalled among forms examined in the macropods. Its perpendicular articular surface exceeds the horizontal surface of the end of the tibia.

The fibula is very slender, especially in its distal three-fourths. Proximally it is compressed and somewhat expanded antero-posteriorly, its inner surface being somewhat grooved. It is surmounted by a rather large ovoid fabella. The tibia and fibula are closely appressed but not fused in the distal half of their length. The hind leg of *Cænolestes* is thus of a cursorial or saltatorial type and closely resembles that of *Antechinomys*, although even in this notably saltatorial form the internal malleolus is relatively small. A leg of similar character with tibia and fibula practically joined distally is found in *Sminthopsis*, *Perameles*, and in the Macropodidae. In almost all other marsupials the tibia and fibula are widely separate throughout.

The relative length of the lower leg in *Cænolestes* is very great. In proportion to the length of the femur, the length of the tibia in *Cænolestes* is only exceeded, among forms examined, by that of the Giant Kangaroo. The proportions in various genera are indicated by the following table,

the ratio given being that of the femur to the tibia. From this table it is seen that this ratio is less in *Cænolestes* than in any other forms examined except the Giant Kangaroo and that it is approximated only by saltatorial forms.

	Length of Femur	Length of Tibia	Ratio of Femur to Tibia
<i>Macropus giganteus</i>	265	465	57
<i>Cænolestes obscurus</i>	14.3	22.5	63.5
<i>Sminthopsis crassicauda</i>	13.6	19.8	68
<i>Antechinomys laniger</i>	19	27	70
<i>Petrogale xanthopus</i>	153	195	78
<i>Phascologale cristicauda</i>	23.3	28	81
<i>Marmosa mexicana</i>	23	26.8	85.8
<i>Dasyurus viverrinus</i>	65	73	89
<i>Myrmecobius fasciatus</i>	45.5	51	89
<i>Perameles nasuta</i>	64.5	69	93.4
<i>Peramys domesticus</i>	21	22	95.4
<i>Philander philander</i>	35.4	37	95.6
<i>Didelphis virginianus</i>	76.8	78	98.4
<i>Sarcophilus harrisi</i>	91.5	91.5	100
<i>Trichosurus vulpecula</i>	94.5	92.5	102.1
<i>Thylacinus cynocephalus</i>	195	190	102.6
<i>Phascolomys ursinus</i>	175	133	131
<i>Phascolarctos cinereus</i>	155	115	134

BONES OF THE FOOT.

Plate XVI, Figs. 2-5.

The foot is long and narrow, the tarsus being relatively short and the metatarsus very long. It has general resemblance to the foot in *Dasyurus*, *Sminthopsis*, *Antechinomys* and similar forms, but its structure shows decided approach to that of *Perameles* and the macropods. Although the hallux is present and relatively well developed (longer than in *Dasyurus*, for example) the relations of the ankle joint are more adapted to cursorial life than in some of the forms in which the hallux has nearly or quite disappeared. This is shown especially by the astragalus which is somewhat intermediate between that of the pedimanous marsupials and that of the highly saltatorial forms. Although it retains general similarity to the astragalus of *Didelphis*, *Sarcophilus*, *Myrmecobius*, *Phascologale* and various other polyprotodonts, it shows a distinct approach to the condition in the Macropodidae in which there is one large trochlea and facets on either side, nearly at right angles, for the outer malleolar parts of the tibia and the fibula. The body of the astragalus stands at right angles to the neck and head and scarcely exceeds them in size. The neck is not constricted to differentiate it

from the head. The inner malleolar facet is extensive and sharply deflected from the medial tibial facet from which it is separated by a high trenchant ridge which extends over the neck to the head at the navicular facet. In all other forms examined, this ridge is discontinuous or confined to the trochlea. The formation of a depression in this ridge between the trochlea and the head would seem to be the first step in the separation of the head from the trochlea. Hence, in this respect the astragalus of *Cænolestes* may be primitive. The tibial facet is very slightly concave and merges imperceptibly with the fibular facet which is narrow and considerably deflected. The sharp division of the two principal tibial facets virtually permits the tibia and astragalus to be locked together almost as effectively as in the macropods. Lateral movement of the tibia is practically precluded and the general freedom of the ankle joint found in most polyprotodonts is quite impossible. Even such forms as *Sminthopsis* and *Antechinomys* have much looser ankle joints than *Cænolestes*.

The inferior surface of the astragalus in *Cænolestes* is largely occupied by the sustentacular facet which is unusually broad and convex. It is nearly at right angles but confluent with the large navicular facet which covers the head of the bone and runs up the inner side for a considerable distance. The ectal facet is crescentic in outline and is separated from the sustentacular by a deep groove. The inner anterior margin of the navicular facet has slight contact with the cuboid bone but there is no distinct development of a cuboid facet. A small sesamoid is situated between the astragalus and the posterior distal extremities of the tibia and fibula.

The os calcis has the tuber laterally compressed and very slightly flexed downward and inward. The anterior sustentacular part of the bone is relatively large and broad. In addition to the usual ectal, sustentacular, and cuboid facets it has a prominent triangular projection on its antero-external surface for the attachment of the tarso-metatarsal and perhaps other ligaments.

The navicular is somewhat anvil-shaped with a smaller dorsal head articulating with the astragalus and slightly constricted from the larger plantar part which spreads inward to the middle of the sole, its plantar surface being broad and flat instead of narrow and ridgelike. Distally it articulates with the ectocuneiform, entocuneiform and mesocuneiform. The cuboid articulates distally with the two outer metatarsals and with the proximo-external surface of the mesocuneiform. Its plantar surface has the usual groove for the tendon of the *peroneus longus* muscle. The ectocuneiform, entocuneiform and mesocuneiform are rhomboidal bones, the first supporting the hallux, and the others the second and third

metatarsals respectively. The ectocuneiform is about twice as large as the mesocuneiform and nearly four times the size of the entocuneiform.

The hallux is supported on the distal end of the ectocuneiform more especially toward its plantar side and has considerable freedom of movement, although it is but slightly divergent. It is much more slender than the other digits and its terminal phalanx reaches only to the distal extremity of the second metatarsals. The proximal bases of the metatarsals form a very uneven row. The fifth reaches farthest proximad, the second and third are considerably short of it, and the first and fourth are still more so. The irregularity in this respect is greater than in any other forms examined. The second and fifth extend distad about the same distance and are slightly exceeded by the third and fourth which also end evenly. The fourth however is somewhat longer than the third by actual measurement since the uneven bases give it greater extension proximad. In a male specimen it is 9 mm. long. The first and also the second phalanges are approximately equal in length from the second to the fifth digits. The paired sesamoids on the metatarsophalangeal joints are as usual. There is no prehallux.

SUMMARY OF SKELETAL CHARACTERS.

The skeleton of *Cænolestes* has the broad general features found in most other marsupials. With the exception of modifications correlated with cursorial limbs, most of its peculiarities are of a primitive or generalized nature rather than of high specialization. The movable cervical rib and the very short pubic symphysis are examples. The general form of the pelvis is remarkably similar to that of some insectivores. In fact the whole skeleton in its general aspects is so closely paralleled by that of one species of insectivore (*Nesogale dobsoni*), which chances to be available, that the two have been illustrated together (Pl. XIX).

Skeletons of the following marsupials have been available for comparison: *Antechinomys laniger*, *Sminthopsis crassicaudata*, *Phascologale* (*Chaetocercus*) *cristicauda*, *Dasyurus viverrinus*, *Thylacinus cynocephalus*, *Sarcophilus harrisi*, *Didelphis virginianus*, *Marmosa marica*, *Peramys domesticus*, *Philander laniger*, *Myrmecobius fasciatus*, *Perameles nasuta*, *Petaurus australis*, *Trichosurus vulpecula*, *Phascolarctos cinereus*, *Macropus giganteus*, and *Phascolomys mitchelli*. A larger representation would have been desirable and it is possible that some conclusions will need modification when wider comparison is made. Numerous minor peculiarities of general characters of the skeleton have been discussed in preceding pages.

The features which seem most noteworthy, although of widely varying significance, are the following:

1. Atlas with open grooves or notches instead of closed foramina for passage of cranial nerves and vertebral arteries; agreeing only with *Perameles*, *Marmosa*, and *Peramys*.
2. Axis with movable cervical rib; agreeing only with certain species (or individuals?) of *Perameles* and *Phascologale*.
3. Seventh cervical tending to become imperforate; intermediate between diprotodonts and polyprotodonts.
4. Spine of first thoracic vertebra scarcely exceeding that of last cervical but greatly exceeded by that of second thoracic; agreeing only with *Perameles*.
5. Vertebral spines converging to a center of motion; disagreeing with Didelphiidae.
6. Costal element of first rib longer than vertebral; agreeing only with *Trichosurus* and *Petaurus*.
7. Ventral surface of xiphisternum not continuous with that of mesosternum; agreeing only with diprotodonts.
8. Pubic symphysis exceedingly short; unique among marsupials.
9. Ischial rami not flaring.
10. Marsupial bone with very long base; unequalled among marsupials.
11. A large heavy humerus combined with a relatively slender forearm; unique among marsupials.
12. Trochlea of humerus extended inward to form a partially separate articular surface; unique among marsupials.
13. A sesamoid in the supinator tendon at the external epicondyle of the humerus; unique among marsupials.
14. Radius flexed forward in its proximal third, in contact with ulna distally; agreeing with saltatorial or semi-saltatorial forms.
15. Pisiform large, relatively larger than in other marsupials examined.
16. Patella ossified; agreeing with *Perameles*, *Notoryctes* and *Tarsipes*; also with extinct forms *Amphiproviverra* and *Prothylacynus*.
17. Femur with a rudimentary third trochanter; agreeing only with *Phascolomys* and *Wynyardia*.
18. Tibia and fibula relatively long and in close contact distally; agreeing only with saltatorial forms.
19. Internal malleolus of tibia large; agreeing with Macropodidae.

20. Astragalus with a large inner malleolar facet nearly perpendicular to the medial tibial facet; agreeing with saltatorial forms.
21. Prehallux not present; agreeing with *Macropus*, *Trichosurus*, and *Phascologomys*; disagreeing with *Didelphis*.

SKULL.

Plates XVII, XX.

General form.—The skull of *Cænolestes* is characterized superficially by the great elongation of the facial region, by the deep and extremely thin-walled braincase, the swollen interorbital and lacrymal regions, the general absence of crests or ridges, the slender zygomata, and by the modifications of the maxillary by which the anterior teeth are at a different level from the posterior.

The general resemblance to the skull of *Perameles* was immediately noticed by Thomas, who says (1895, p. 872): "Skull in its general proportions something like that of a *Perameles*, although thinner and more delicately built, with a similarly elongated muzzle, smooth and rounded brain-case, and obsolete supraorbital and cranial crest and ridges; the zygomata are, however, so much more boldly expanded as somewhat to spoil the resemblance, which in any case does not apply to details." A still stronger statement of this resemblance to the skull of *Perameles* might be made and, as appears later, it does apply at least to a certain number of details. The zygomata are more spreading than in some of the peramelids but their sweep is practically identical with that of others (e. g. *Isoodon*). Certainly the general form of the skull is nearer that seen in the peramelids than that of any other living marsupial. The resemblances of general shape to the smaller dasyurids noted by Miss Dederer (1909) seem mostly to be associated merely with small size and are found also in *Acrobates* and *Dromicia*.

The ratio of zygomatic breadth to total length of skull in various marsupials is as follows: *Perameles nasuta* 39.7, *Perameles bougainvillei* 43.2, *Cænolestes* 47.7, *Isoodon barrowensis* 51, *Philander* 54, *Phascolarctos* 54.3, *Myrmecobius* 56.4, *Peramys* 56.9, *Didelphis* 57, *Chironectes* 57.1, *Phascologale* (*Chætocercus*) *cristicauda* 63.7, *Petaurus australis* 66.4, *Dactylopsila* 67.5, *Dasyurus viverrinus* 69.7, *Trichosurus vulpecula* 71.7, *Sarcophilus* 81.2. In this series, *Cænolestes* falls with the peramelids. The didelphids occupy a position midway between these dolichocephalic forms and the broader types presented among the phalangiers and dasyurids.

Braincase.—The braincase is relatively large and its walls are

exceedingly thin and papery. Viewed in longitudinal section (Pl. XVII) its depth is seen to be proportionately greater than in other marsupials. The ratio of the depth to the length measured inside the braincase in several forms is found to be as follows: *Cænolestes* 53.9; *Perameles nasuta* 48.8; *Petaurus australis* 47; *Myrmecobius* 44.4; *Trichosurus* 42.2; *Phascogale flavipes* 41.7; *Marmosa cinerea* 41.6; *Didelphis virginiana* 39.7; *Peramys domesticus* 38.6.

The dorsal outline of the braincase is broadly arched, agreeing in this respect with *Perameles* and with most diprotodonts and contrasting with the didelphids and dasyurids in which the dorsal outline is more nearly straight. The olfactory fossa is large and full, especially in its lateral parts, but it is relatively short antero-posteriorly as compared to didelphids and dasyurids. In these polyprotodonts its median section is subtriangular, often with an acute angle in front, whereas in *Cænolestes* and *Perameles*, and the several diprotodonts examined, it is broadly rounded anteriorly. It is separated from the cerebral fossa by a light ridge, the position of which is plainly indicated externally by the furrow running upward and slightly backward from the ethmoid foramen.

The cerebral fossa is very full, especially in its lower anterior part where it is extended forward on either side of the median part of the cribriform plate. The floor of the braincase is thus wider and flatter anteriorly than in other marsupials and it meets the cribriform plate more abruptly. This is in the region occupied by the tuberculum olfactorium which, as elsewhere noted, is unusually large (see p. 159). In front of the occipito-sphenoid suture is a shallow pituitary fossa. The sphenoidal fissure and the foramen rotundum open posteriorly on the floor of the braincase and have no lateral exposure like that in other marsupials. Externally the sphenoidal fissure and the foramen rotundum have what is practically a common opening, the bony septum between them being quite internal and not visible in lateral view. In this, the condition seems nearer that of the diprotodonts than of the polyprotodonts in which the foramen rotundum opens decidedly posterior to the sphenoidal fissure with the septum between them broadly exposed in lateral view.

The cerebellar fossa is smooth and without obvious division into lateral and median parts. It is separated from the cerebral fossa by an exceedingly slight tentorial ridge. The outline of the cribriform plate as seen in section is more similar to that of *Perameles* than to that of any other form examined. It is more nearly vertical than in polyprotodonts generally and more abruptly divided between its upright and its posterior basal extension in the presphenoid region. This posterior extension is broader and higher than in any other marsupial examined.

The foramina leaving the braincase include the sphenoidal fissure and foramen rotundum, which already have been mentioned, the foramen ovale, transverse canal, entocarotid foramen, and ethmoid foramen, in all of which the relations are generally similar to those in other marsupials although there is much variation, especially in diprotodonts.

Ethmoid and turbinals.—The cribriform plate of the ethmoid, as previously stated, is more nearly upright than in the didelphids and resembles that of *Perameles*. Its exact relations with the orbitosphenoid and presphenoid are not distinguishable in the adult, but the bony floor of the olfactory fossa is high and has a more rounded surface than in other marsupials. It extends exactly parallel to the palate and hence nearly at right angles to the anterior wall of the braincase. The principal ethmo-turbinals visible in a simple section (Pl. XVII, *et.*) are four in number. The lowermost is relatively larger than in any other form examined and is most nearly approached in *Perameles nasuta*. The uppermost and longest of the ethmo-turbinals is relatively short, extending forward only to the plane of the front of the last premolar. This seems to be a resemblance to diprotodonts since it agrees with the condition in *Trichosurus* and *Petaurus*. In the didelphids and in *Phascologale* this upper ethmo-turbinal is very long, reaching as far forward as the canines and almost completely inclosing the fenestrated part of the maxillo-turbinals. In *Perameles* and *Myrmecobius* an intermediate condition appears.

The naso-turbinal offers no especial peculiarities. The maxillo-turbinal springs from the nasal wall at a somewhat higher level than in polyprotodonts and the nasal floor is narrower.¹ In these respects, there is some variation among polyprotodonts but in general the resemblance seems to be to diprotodonts. The maxillo-turbinal, as in diprotodonts, is not largely enclosed by the anterior extension of the ethmo-turbinal. The mesethmoid is a thin vertical plate of the usual character.

Nasals.—These are long and slender, coming nearly to a point anteriorly and being abruptly expanded posteriorly. Their anterior extremities exceed the dorsal edges of the premaxillae and overhang the anterior nares, reaching nearly as far forward as the lower anterior extension of the premaxillae. In this they differ markedly from the Dasyuridae, Thylacynidae, and Myrmecobiidae but are approached by most diprotodonts, by the didelphids and especially by the peramelids. The posterior expansions of the nasals include a slender curved prong on each side projecting across the extraordinary vacuity which lies

¹ This is contrary to the observation of Broom (1911, p. 318), whose material may have been imperfect.

between the nasals, frontals and maxillary. They do not exclude the maxillaries from contact with the frontals, however, and despite their great length their posterior endings are in advance of the orbits.

Premaxillae.—The premaxillae are longer than high and reach quite to the bases of the canines. The maxillary suture approaches the perpendicular as in *Perameles* and most diprotodonts. A narrow posterior ascending branch extends backward between the nasals and the maxillary to the plane of the front of the middle premolar⁶. The palatine processes of the premaxillae do not extend the full length of the anterior palatine vacuities as in most other marsupials, but meet similar processes of the maxillary which extend forward. The bony division of the anterior vacuities thus consists of premaxillary elements only in its anterior three-fifths. The only other marsupials in which I have found this arrangement are the peramelids; in all others the premaxillary processes extend nearly or quite to the posterior boundary of the vacuities. In every respect, therefore, the premaxillae are more similar to those of the peramelids than of any other marsupial.

Maxillae.—The slender processes of the maxillaries projecting forward in the anterior palatine vacuities are only paralleled among the peramelids. The posterior fenestration of the palatal aspect of the maxillae also resembles closely that of *Perameles*. In their upward extension, the maxillae are not greatly narrowed between the antorbital vacuities and the lacrymals, but meet the frontals broadly at a point about even with the front of the orbits and somewhat caudad of the posterior nasal endings. In these respects they are conspicuously different from those of the didelphids.

In uniting with the jugals, the maxillaries send out a slender prong which takes a larger part in the formation of the anterior root of the zygoma than is usual in marsupials and contrasts especially with the condition in most polyprotodonts in which the jugal is expanded on the face below the lacrymals. In general, however, the relation of the anterior root of the zygoma to the maxillary is as in polyprotodonts, that is, the principal part of the maxilla, including all the molars except the last two, is anterior to the zygoma. This is in decided contrast to the diprotodonts in which that part of the maxillary bearing the molars is largely posterior to the anterior root of the zygoma. In some forms, especially among the Phalangeridae, the root of the zygoma rises from a point as far forward as the last premolar and consequently the entire molar series is posterior to it. In polyprotodonts, on the contrary, it is rare for more than the last reduced molar to lie posterior to the zygoma. This constitutes one of the most striking resemblances to polyprotodonts shown by the skull of *Cenolestes*.

Frontals.—The frontals are nearly as wide anteriorly as posteriorly. They are smoothly rounded at the orbital border and have neither ridges nor post-orbital processes, thus differing from most polyprotodonts. There is no post-orbital process in *Phascologale* and some allied forms but in these the frontals are more flattened and the orbital border is more angular. Within the orbit, the frontals of *Cænolestes* are less deflected toward the midventral line than in other marsupials and thus provide for the unusual space required by the large olfactory lobes of the brain. A distinct venous foramen perforates the orbital border in its upper anterior part.

Lacrymals.—The lacrymal is of moderate size and not largely extended beyond the orbit. It is slightly gibbous but not ridged between its facial and orbital parts. It is well separated from the nasals by the frontal process of the maxillary. The lacrymal canal has a single opening, thus differing from the didelphids and many other marsupials. It opens on the orbital border, not within the orbit, but not so definitely outside it as in other forms.¹

Parietals.—The parietals are broad and smooth with no suggestion of a sagittal ridge. The suture between the two parietals and that between them and the supraoccipital is practically obliterated in adults although easily visible in most other marsupials except the peramelids. Apparently the parietals reach laterally to the vicinity of the lambdoid ridge as in other forms and are not shortened as in the figure published by Miss Dederer (1909). A minute postparietal foramen is usually present.

Zygomata.—The anterior root of the zygoma, as previously stated, rises at the level of the third molar in marked contrast to the position general among diprotodonts. The jugal is sharply narrowed anteriorly and comes to a point just below the lacrymal canal. The arch has a general sweep somewhat as in *Phascologale* and the bone is similarly flattened but the anterior base is narrower and there is no trace of a postorbital process, this last character being a resemblance to *Perameles*. The jugal reaches the glenoid fossa as usual and its ending is truncate to form a slight anterior boundary for the fossa, thus agreeing more nearly with polyprotodonts than diprotodonts. The posterior root of the zygoma is not inflated as in most diprotodonts, but it is more slender than in dasyurids and didelphids and more nearly resembles the condition in the peramelids.

Squamosal.—The cranial part of the squamosal is not peculiar, being

¹ In Gregory's exhaustive comparative study of the lacrymal (Bull. Am. Mus. Nat. Hist., XLII, p. 235, 1920), made since the above was written, it is concluded that "The evidence of the lacrymal region is in harmony with the view that the cænolestoids are allied with the Peramelidae."

smaller than in *Perameles* and *Sarcophilus*, smaller also than in some diprotodonts, but relatively quite as large as in others. Its proportions are nearly as in the didelphids and smaller dasyurids. The glenoid fossa is considerably above the basicranial axis and is distinctly longer than wide. Except on its extreme outer edge where the end of the jugal bounds it, it is open in front. Posteriorly it is bounded on the outer side by a papillate bony elevation which is unconnected with the tympanic and has only very slight transverse extent. The general features of the glenoid fossa are thus somewhat intermediate between those in polyprotodonts and in diprotodonts. The transverse extent and the broad platelike posterior boundary usually found in polyprotodonts are changed in the direction of the narrower more open condition of the diprotodonts which allows of greater antero-posterior motion of the jaws in conformity with the requirements of their dentition. The arrangement of the foramina at the base of the squamosal is slightly different from that of the Didelphiidae and similar to that in the smaller Dasyuridae and the Peramelidae. As in these, the postzygomatic foramen is lacking and the postglenoid and subsquamosal are closely associated. The subsquamosal foramen opens laterally just behind the postglenoid process and the postglenoid foramen antero-ventrally between the postglenoid process and the superior end of the tympanic. A thin bony plate bounds the subsquamosal foramen ventrally and extends from the postglenoid process to a short flat mastoid process of the squamosal.

Occipitals.—Contrary to the usual condition in marsupials, supraoccipital, exoccipitals and basioccipital are completely ankylosed in adult skulls making it impossible to determine their exact boundaries. In other forms, especially among polyprotodonts, the sutures separating these bones are usually obvious. The supraoccipital shows no clear indication of a forward extension to the dorsal surface of the skull and the median occipital region is smooth without any sharp angle between the dorsal and occipital surfaces. Laterally a slight ridge rises on each side suggesting a lambdoid crest at the union of the supraoccipital and the parietals.

The exoccipitals are small and extend but little beyond the condyles. Probably they do not meet above the foramen magnum. Small paroccipital processes lie prostrate between the mastoid and the petiotic and do not form projecting points. Their condition is nearly the same as in *Perameles* and *Myrmecobius* but different from other forms examined. The exoccipitals are perforated by two small condylar foramina near the apparent line of union with the basioccipital.

The basioccipital is deeply cleft by the inferior lip of the foramen

magnum. Its ventral surface is smooth or even slightly concave medially without suggestion of a central ridge. This constitutes another slight and perhaps unimportant, but still practically exclusive, resemblance to the peramelids.

Sphenoid bones.—The basisphenoid is decidedly concave on its exposed surface but has no sharp lateral ridges. It is perforated as usual by the transverse canal and the entocarotid foramen. The suture between it and the basioccipital is closed in adults although usually indicated by a faint transverse ridge. On its superior or cranial surface it is excavated posteriorly for the reception of the hypophysis but there is no distinct sella turcica nor clinoid process.

The presphenoid is only slightly exposed ventrally, being largely inclosed by the thin irregular extensions of the palatine, vomer, and pterygoids. The suture between it and the basisphenoid, usually so distinctly open, is but faintly indicated. Within, the vertical part of the presphenoid is slightly thickened basally but soon becomes exceedingly thin. The orbitosphenoid occupies a small area between the lacrymal and the sphenoidal fissure, its exact boundaries being uncertain. The alisphenoids show no peculiarities beyond those connected with the unusual fullness of the braincase in the region covered by them. As stated elsewhere, the sphenoidal fissure and foramen rotundum are practically confluent just before reaching the exterior but those of the two sides of the skull are more distinctly separated from each other than in any other form examined. The alisphenoid bulla forms a simple subhemispherical capsule similar to that of the didelphids and dasyurids.

Palatine.—Owing to the very large posterior vacuities the palatine bones are relatively small. In the median line they send a pair of slender processes forward to the plane of the middle of the second molar where they meet similar processes from the maxillaries. Posteriorly they are bounded by a slight transverse ridge which is perforated on each side by a small postero-lateral foramen. In close proximity to this foramen is the slightly larger sphenopalatine foramen piercing the upright plate. This plate forms the side walls of the posterior part of the nasal passage, but anterior to the sphenopalatine foramen the walls become discontinuous since the turbinals are fenestrated and exposed much farther caudad than usual and the narrow vomer does not reach the side walls. Posteriorly the palatines form part of the sloping ridge which supports the pterygoids. The pterygoid processes are essentially similar to those of polyprotodonts although they take the form of slender rounded prongs rather than flattened subtriangular plates.

Vomer.—The vomer is narrow in front and expanded posteriorly

enclosing somewhat more of the ventral surface of the presphenoid than usual. It is not emarginate posteriorly and extends slightly beyond the posterior nares. Its ventral surface has a low median ridge but this has no contact with the palatines nor with the maxillaries except anteriorly.

Tympanic.—The tympanic is a simple ring open behind and attached to the alisphenoid bulla by cartilage. It is wider and heavier than in the didelphids but is essentially the same as in Australian polyprotodonts.

Periotic.—The petrous part of the periotic is exposed ventrally as a subtriangular bone between the basioccipital, the tympanic and the alisphenoid bulla. It is thus almost wholly exposed as in polyprotodonts and not partly or almost completely covered by the alisphenoid bulla and the tympanic as in diprotodonts. It is less inflated than in the dasyurids but projects below the surface of the basioccipital slightly more than in the didelphids. Internally it shows the usual openings, including a large floccular fossa. The foramina surrounding it are much as in the other marsupials except for the so-called carotid canal which forms a long open slit between the petrous periotic and the basioccipital which are actually in contact scarcely more than enough to separate this opening from the jugular foramen behind and the entocarotid foramen in front. Such an opening has not been found in any other marsupial examined, although in polyprotodonts it may be represented by the small foramen which Gregory (1910, p. 223) has called the posterior carotid foramen.

The mastoid is relatively larger than in any other form examined and is especially characterized by its great lateral extent. Its occipital surface is not more than one-third its lateral and between them there is only a gentle curve or an exceedingly feeble ridge. The lateral surface is slightly convex and meets the squamosal evenly. The short flattened paroccipital process and the mastoid process of the squamosal clasp it on either side making the entire mastoid region relatively smooth and continuous with the general outlines of the skull. This lack of prominences is in marked contrast to the condition in the didelphids and is nearer to that of *Antechinomys* and other primitive dasyurids. A very large mastoid foramen is situated at the upper end of the occipital part of the mastoid separating it from the supraoccipital for a considerable distance. A mastoid foramen occurs in the didelphids but in adults it is very small or practically closed, whereas in *Canolestes* it is as large in the adult as in the very small pouch young of *Didelphis*.

Hyoid.—The hyoid apparatus is of the usual marsupial type. The basihyal is flat and subcircular with its posterior edge free but its anterior and lateral boundaries united with the ceratohyals and thyrohyals. The ceratohyals are slightly arched forward and only ossified

basally, becoming cartilaginous and slender in their distal two-thirds. The thyrohyals are flattened bars standing nearly at right angles to the general axis and ossified to their tips which are united by a short cartilage to the osseous wing of the thyroid cartilage.

Vacuities.—The palate is highly fenestrate, its bony floor being relatively less extensive than in any other living marsupials with the possible exceptions of some of the peramelids. There are two pairs of vacuities of constant occurrence and fairly regular form. The anterior pair extend from the front of the second lateral incisor to the front of the middle premolar and occupy nearly all the space between the tooth-rows. They are divided by the median processes of the maxillaries and premaxillaries which are flattened ventrally to an extent about equalling one-third the width across the paired vacuities. These vacuities are thus about equal to the combined length of the two separate pairs found in peramelids and they are much longer than the single pair of most other polyprotodonts although they are rather closely approached by *Sminthopsis*. The posterior vacuities begin at the front of the last premolar and reach nearly to the transverse ridge bounding the palate. Their endings are almost exactly at the level of the back of the last molars. They are divided by a narrow median extension of the maxillaries and a similar one from the palatine, the two meeting midway of the length of the vacuities and forming a slight expansion at their union. Laterally the boundaries of the vacuities are slightly irregular but anteriorly they are always evenly rounded and the general shape and extent in different specimens is very constant.

The palatal vacuities are regarded as secondary developments and, since they vary so widely throughout the marsupial group, it is not probable that their character in *Cænolestes* has any special significance. The palate in extinct cænolestids is only imperfectly known.

A prominent vacuity on each side of the face directly above the infraorbital foramen is one of the most unusual features of the skull of *Cænolestes*. It is bounded by the nasal, frontal and maxillary bones and opens into the large sinus between the naso-turbinal and the maxillary. Its relations to the overlying dermal tissues are simple and no glandular or other specialized development is apparent. As noted by Thomas (1895), a vacuity in this part of the skull is found in other mammals only among ungulates.

Auditory ossicles.—The malleus has the general features usual in marsupials. The neck is relatively short, the lamina very thin, and the processus muscularis is but feebly indicated. The manubrium is about two-thirds as long as the processus gracilis and forms a thin flattened blade with a pointed apex and a slightly falcate outline when seen from

the flat side. The processus brevis at the base of the manubrium is relatively insignificant. The processus gracilis has two thin longitudinal ridges bounding a long v-shaped groove which seats the tympanic ring. The extremity of the process is somewhat expanded and obtuse. The union with the tympanic is close and considerable care is required in forcing the parts away from each other. The malleus differs from that of *Perameles* (*P. bougainvillei*) principally in its shorter neck; from that of *Didelphis* it is distinguished by its generally frailer construction and its closer union with the tympanic.

The incus is practically identical with that of *Perameles*, differing mainly in the processus brevis which is somewhat compressed and truncate instead of long and conical. The processus longus is very slender distad of the sharp right angle at which it is bent. The Sylvian apophysis is elliptical. The incus in adults is rather firmly attached to the malleus and does not separate except after slight pressure.

The stapes is strictly columelliform without trace of an opening and there is not even an appreciable difference in its diameter from the head to the base. The head is very small and the base relatively large and elliptical. In adults, the stapes is well ossified.

The auditory ossicles of *Cænolestes* are clearly of a type closely approaching that of *Perameles*, which was regarded by Doran (1879) as the lowest among marsupials. They also show resemblance to some of the dasyures, which, as described by Doran, have the stapes columelliform. Whether this simple stapes is in truth a primitive character or not, it serves at least to distinguish *Cænolestes* sharply from the didelphids, in which the stapes is always bicurrate. So far as the auditory ossicles are concerned, therefore, *Cænolestes* resembles Australian rather than American forms and especially approaches the peramelids and dasyurids.

Mandible.—The mandible is slender and relatively straight anteriorly, its greatest curvature being just below the base of the coronoid process. The symphysis is long but weak. The rami have exceptionally little divergence. The coronoid is large, broad, and nearly upright. Its anterior border is gently curved but approaches the vertical more closely than in any polyprotodonts. The broad masseteric fossa is rather shallow and is indistinctly divided into an anterior and a posterior portion by a weak elevation between the tips of the coronoid and the condyle. Near the lower edge of the masseteric fossa is a tiny foramen, indistinct in some specimens, but obvious in others. A similar foramen is found in many Australian diprotodonts but does not occur in polyprotodonts. Its presence in *Abderites* and *Garzonia* has been noted by Ameghino and Sinclair. Mental foramina of the usual character are present, a larger

one below the posterior root of the middle premolar and a smaller below the anterior root of the first molar.

On the lingual side of the mandible the dental foramen is situated as usual at the base of the angular process. In some specimens a shallow groove runs along the inner side of the proximal half of the jaw but in no case does its position or character seem to warrant any assumption that it is a true Meckelian groove.

The angular process is moderately inflected and subtriangular in shape with a fairly wide base and a rounded but not decidedly obtuse tip. The inflection is slightly less than in the majority of polyprotodonts and the process is not produced into a slender prong like that in the smaller dasyurids. The condyle is relatively high in position, slightly more than halfway from the angle to the tip of the coronoid and extending well beyond a line drawn between them. In this respect, therefore, it is more like *Phascologale* and other dasyurids than like the didelphids in which the condyle is seated on a relatively short base.

The length of the bony lower jaw is proportionately less than in most polyprotodonts and this is especially true if it is considered in relation to the upper jaw. In two specimens of nearly equal size, the length of the bony part of the jaw in *Cænolestes* is 77 per cent of the total length (including terminal incisors) and that in *Phascologale* is 95 per cent.

Owing to the wide variation in the lower jaws of marsupials, it is difficult to draw any conclusions from the findings in regard to *Cænolestes*. Miss Dederer (1909) has regarded the jaw of *Cænolestes* as indicative of polyprotodont affinity, believing it to be very similar especially to *Antechinomys* and *Sminthopsis*. It differs from these in many respects, as in the broader more upright coronoid, the less inflected and less attenuated angle, the less divergent rami, and in the presence of a foramen in the masseteric fossa, this last being a diprotodont character.

Among the didelphids, the mandible shows considerable variation and comparison is difficult. It might be said that as respects the angle, *Cænolestes* is somewhat intermediate between *Didelphis* and *Philander* and as respects other characters it shows no especial indications of relationship to the didelphids more than to other marsupials.

SUMMARY OF CRANIAL CHARACTERS.

The great range of variation among both polyprotodonts and diprotodonts leaves relatively few characters that are diagnostic of the two groups. Without a complete representation of all known forms it can only be said that certain characters are regarded as polyprotodont or diprotodont because they are mutually exclusive in the majority of forms examined. On this basis, it is clear that the skull of *Cænolestes*

shows more polyprotodont than diprotodont characters. These are as follows:

1. Anterior root of zygoma only slightly in advance of end of toothrow.
2. Posterior root of zygoma not inflated.
3. Tympanic narrow and annular.
4. Petrous part of periotic largely exposed ventrally.
5. Alisphenoid bulla not fused with paroccipital process.
6. Foramina of squamosal arranged in general as in Dasyuridae.

Practically all of these characters relate to the otic bones or to the backward extension of the palate with relation to the zygomata. In the extinct *Palæothentes*, a close relative of *Cænolestes*, the zygomata and palate are practically as in diprotodonts, so the polyprotodont characters of the skull in cænolestids are reduced to those of the tympanic region. The additional characters supposed to indicate polyprotodont affinity and cited by Miss Dederer (1909) and Broom (1911) are found in the smaller diprotodonts, as *Dromicia* and *Tarsipes*. Hence their importance is doubtful.

A few characters are at least in agreement with those of some diprotodonts, and with no polyprotodonts so far as known. These are the following:

1. Upper anterior ethmo-turbinal short and enclosing but little of the maxillo-turbinal.
2. Maxillo-turbinal arising relatively high up on the nasal wall.
3. Glenoid fossa relatively long and with a small postglenoid process which is not extended transversely.
4. A small foramen in masseteric fossa of mandible.

Distinctions in cranial characters between *Cænolestes* and the didelphids are numerous; in fact the only characters in common are of quite a general nature.¹ On the other hand, resemblances to Australian forms, especially peramelids, are abundant. In the following respects there is agreement with some or all Peramelidae or Dasyuridae and disagreement with all Didelphyidae:

1. Rostrum slender and elongate.
2. Palatine processes of premaxillae not extending full length of incisive foramina.
3. Basioccipital concave medially.
4. Maxillo-premaxillary suture nearly upright anteriorly.
5. Cribriform plate approaching the vertical.
6. Fronto-maxillary suture broad.

¹The mastoid foramen may be an exclusive resemblance, but in the absence of immature specimens I am unable to learn whether or not it occurs in dasyurids.

7. Sagittal and lambdoid crests absent.
8. Postorbital processes absent.
9. Paroccipital and mastoid processes small and prostrate.
10. Lacrymal canal with a single opening.
11. Postzygomatic foramen lacking.
12. Stapes columelliform.

The first five of these characters are exclusive resemblances to the peramelids while the others are common to the peramelids and certain of the dasyurids. The total number of resemblances to the peramelids therefore, is very large. Moreover, important distinctions between *Cænolestes* and *Perameles* are comparatively few, being confined mainly to the region between the sphenoidal fissures and to the proportions of the processes of the mandible.

In a number of respects the skull of *Cænolestes* differs from all other living marsupials. The most important of these are as follows:

1. A preorbital vacuity between the nasals, maxillary, and frontal.
2. A large mastoid foramen persisting in adults.
3. A long narrow carotid canal between the petrous periotic and the basioccipital.
4. Floor of braincase very wide between sphenoidal fissures.
5. Mastoid large and broadly exposed laterally (approached by peramelids).
6. Olfactory fossa relatively large and wide.

DENTITION.

Plate XVIII.

The dentition and the cranium of *Cænolestes* have been known since the original discovery of the animal and have been variously described, figured, and subjected to comment by different authors, notably Thomas, Bensley, Sinclair, Dederer, and Gregory. The diprotodont modification of its lower incisors combined with a polyprotodont upper incisor formula and quadrate bunolophodont molars at once furnished conditions not found elsewhere among living marsupials and not typically representing either the diprotodont or the polyprotodont group. The teeth have been well described in considerable detail by Thomas (1895), but his specimen was somewhat affected by wear, which obscured a few important points. For this reason and for the sake of completeness and convenience of discussion full description of the teeth is given in the following pages. Unfortunately it is still impossible to give any information as to the succession of the teeth, since all the specimens at hand are fully adult.

NUMBER AND HOMOLOGIES OF TEETH.

The teeth are 46 to 48 in number. In the upper jaw there are one pair of terminal incisors and three pairs of lateral ones, a pair of sharp, well-differentiated canines, three pairs of more or less triconodont premolars, and four pairs of molars, the first and second of which are quadrate and the third and fourth subtriangular. In the lower jaw there is a pair of long terminal chisel-like incisors resembling those of the most pronounced diprotodont dentitions. Behind these in lateral position are four and sometimes five pairs of small unicuspid, practically undifferentiated from each other, which are interpreted as two (or three) incisors, one canine, and one premolar. Next are two pairs of double-rooted premolars and four pairs of molars. The dentition, therefore, is a rather highly modified one but retains approximate numerical agreement (at least in certain specimens) with the polyprotodont dentition usually regarded as generalized. With the exception of the didelphids, which have five upper incisors, and *Myrmecobius*, which has supernumerary molars, no living polyprotodont has more teeth than *Cænolestes*. The formula may be written:

$$I. \frac{4}{3-4}; C. \frac{1}{1}; Pm. \frac{3}{3}; M. \frac{4}{4}; x2 = 46-48.$$

This is essentially the classification made by Thomas (1895) who divided the four unicuspid behind the median incisor into two incisors, one canine, and one premolar. "Any other determination," he says, "would involve the presence of four incisors or four premolars, each equally unlikely." It now appears that, although the normal number of lower incisors is three as decided by Thomas, the presence of four would by no means have been unlikely. The number of lower unicuspid or intermediate teeth is normally four, but a variation in which there are five is not infrequent. In one specimen (Field Museum No. 18603), there are five on the right side of the jaw and four on the left. Bensley (1903, Pl. 5, fig. 38) has figured a specimen belonging to the British Museum in which there are five on each side of the jaw. Others, so far as examined, have four unicuspid on each side, but since the total number of specimens is very small the percentage of variation is high. It is possible, therefore, that these aberrant specimens represent a condition once normal and in which the lower antemolar formula was equal to that of the didelphids and in excess of other living marsupials. This is further suggested by the occurrence of the same lower antemolar formula in the extinct genus *Halmarhiphus*, regarded by Sinclair as the direct ancestor of *Cænolestes*. A closely allied form, *Garzonia*, has as

many as six unicuspid in the lower jaw, giving it a total antemolar formula of nine, exceeding that of any living mammal. While this formula, as observed in only one specimen, may be abnormal, as suggested by Sinclair, it is evident that the formula of *Cænolestes*, large though it be, has already undergone a reduction. A further and gradual reduction may be traced in Australian diprotodonts to those in which only the median or terminal pair of incisors and one pair of differentiated premolars remain, two pairs of antemolar teeth in all, as in *Phascolarctos* and *Phascolomys*.

Representatives in this series with the number of antemolar teeth in each would be as follows: *Garzonia*, 9; *Halmarhiphus*, 8; *Cænolestes*, 7-8; *Phalanger*, 6-7;¹ *Acrobates*, 5; *Dromicia*, 4-5; *Distaechurus*, 4; *Trichosurus*, 3-4; *Macropus*, 3; *Phascolarctos*, 2. When the form and character of the teeth as well as their number is considered, the probability that the series is an expression of homologies seems very great. Among the living forms of the *Phalangeridae* the reduction is going on at present, as indicated by the great instability of the number of the "small intermediate teeth" in various species. This was shown especially in the genus *Phalanger* by Jentink (1885) who examined a considerable series and found great variation. The subject has been further elaborated by Bateson (1894). In the extinct forms *Palæotheres*, *Acdestis*, and *Callomenus*, a reduction also has occurred, these having only five or six antemolar teeth. Hence, *Cænolestes* is more primitive than these. In fact, so far as mere number of teeth is concerned, *Cænolestes* and the allied extinct forms like *Halmarhiphus* and *Garzonia* are almost as primitive as the didelphids. Assuming their common ancestry, it is necessary to believe that the diprotodont modification preceded any numerical reduction. Moreover, it is not impossible that all living and extinct polyprotodonts are more reduced than these primitive diprotodonts. The case of *Garzonia* with its nine lower antemolar teeth is of interest in this connection. At least five of these must be regarded as incisors unless reduplication with no reference to homologies be assumed. This is a larger number of lower incisors than is possessed by any known polyprotodont. Further, if the specialized median incisor is not the first but the second as shown by embryology in the macropods (Woodward 1893), we must assume the ancestral formula to be six. Such an assumption is further justified by Woodward's (l. c.) discovery of vestiges of six upper incisors in the macropods and by the occurrence as an abnormality of six upper incisors in *Didelphis* (Bateson 1894, p. 247; Allen 1901, p. 158). If there were six in the upper jaw there may well have been six in the lower, although of course it is not yet known that the cænolestids

¹In abnormal cases only.

agree with the macropods in the ontogeny of the upper jaw. The dental formula of the primitive diprotodont would then have been as follows:

$$I. \frac{6}{6}; C. \frac{1}{1}; Pm. \frac{3}{3}; M. \frac{4}{4} \times 2 = 56.$$

The earliest known mammals, whether they were pro-marsupials or not, furnish no indication that such a large mammalian incisor formula ever existed and it must be admitted that the evidence in favor of it is little more than suggestive. Still, if early mammals only were considered, it would be almost as difficult to believe in an incisor formula of $\frac{6}{6}$ as one of $\frac{8}{8}$, for the usual number of lower incisors in the Triconodonta and Trituberculata was four. Both Thomas (1888) and Winge (1895) have presented evidence that the primitive formula was at least $\frac{6}{6}$ and the case for it must be regarded as considerably stronger than that for one of $\frac{8}{8}$ so to that extent the testimony of the little known "pro-marsupials" is weakened. In fact, in this as in some other respects, it is evident that the long known Mesozoic mammals are extremely liable to misinterpretation. Evidence is accumulating to show that even these early triconodonts were not altogether generalized and many connectant forms are needed before the main mammalian stem will be definitely revealed. The highly specialized nature of the dentition in some of the theriodont reptiles, including both secodont and crushing types and even showing a mammalian succession, suggest that some basic lines of the general evolution of mammalian teeth may have been laid down in these early reptiles and, although lost in the triconodonts, persisted in still later forms the immediate ancestors of which are unknown. Although most of these theriodonts had only four upper incisors on a side, there were some (e. g. *Pristerognathus*) which had as many as six. Hence a mammal with six is theoretically open to no serious objection. In any case it is clear that as early as Miocene times *Canolestes* were at least as primitive as polyprotodonts in the number of the teeth in the lower jaw. The number in the upper jaw is one less than in the didelphids and certain of the peramelids, which have five upper incisors, most other polyprotodonts having four.

The lateral lower incisors of *Canolestes* are small and while not wholly functionless are apparently approaching that condition, so variations in their number are more to be expected than would be the case with fully functional teeth. It is possible that the occasional appearance of an extra unicuspid has no reference to ancestral conditions and therefore that all the specimens of the extinct forms *Halmarhiphus* and *Garzonina* are abnormal. Evidently this would be the conclusion of Bateson (1894) who places no confidence in "reversion" or ancestral influences as the

explanation of such variations. However, he regards "the variability of a form as much a part of its specific character as any other feature of its organization." Without dissenting from this, it may be added that specific and especially generic characters (regardless of how they were initiated or established) are themselves the result or sequence of ancestral conditions. Whether or not characters are "presence or absence" characters, that is, "meristic" rather than "substantive," matters little. In specialized dentitions the loss of a series of several teeth can plainly be traced from species or genera in which the whole series is constantly present to those in which all of it is constantly absent. In such a series, the intermediate forms, or those in which the process of reduction is going on, are notoriously variable. How else, it may be asked, could a reduction take place, especially if discontinuous variation be assumed? No better example of a variable intermediate type could be had than *Phalanger*, which is treated by Bateson at considerable length. In the artificial series from the extinct cænolestid *Garzonia* to the modern *Phascolarctos* the lower antemolar teeth are reduced from nine to two and the "small intermediate" teeth from seven to none. *Phalanger* lies between the two extremes. Among 76 specimens of one species (*P. orientalis*) examined (Bateson, p. 253), 57 had three intermediate teeth on each side of the mandible. The remaining 19 varied from those having only one on each side to one specimen having five on one side and four on the other. Of these, 13 had at least three intermediate teeth on one side of the jaw. This variability is what might be expected as the result of interbreeding after the occurrence of a discontinuous variation in certain individuals. Or, as seems more probable to a taxonomist and comparative anatomist, it may be the result of the interbreeding of individuals which have in varying degrees gradually developed tendencies toward the loss of certain teeth. Such tendencies are illustrated by the familiar example of the tardily developed and frequently absent third molar or wisdom tooth of *Homo*. The vestigial incisors (many of which are partly calcified) which have been found in all marsupials investigated by embryologists would doubtless be admitted by everyone as indications of ancestral conditions. These vestigial incisors are shown by Woodward and Wilson and Hill to be, at least in some cases, not members of a pre-existing series (lacteals or prelacteals) but persistent rudiments of suppressed members of the existing series. Hence it seems not improbable that there may be cases in which these vestiges arise in place and become functional as "super-numerary" incisors. An embryological study of an extremely variable form like *Phalanger* should yield interesting results bearing on this point.

Therefore, while it is undeniable that some cases of supernumerary teeth are teratological with a basis that may be largely physiological and mechanical, it cannot be admitted that there are no cases of which phylogenetic influence is the chief determining factor. The objectionable term reversion need not be applied to them but it might be justified to refer to them as examples of *persistence* of ancestral characters. Aside from what has been said above, it is evident to anyone familiar with the dentition of many species of mammals, that cases are innumerable in which a nonfunctional or slightly functional tooth is present in one species and normally absent but occasionally present in a closely related species. Among bats and rodents instances are particularly abundant. The homology of such a tooth is invariably too clear to be questioned for a moment.

That the fifth lower unicuspid sometimes found in *Cænolestes* is evidence of an ancestor in which this tooth was normally developed may not now be proved conclusively but the evidence from paleontology, from embryology, and from phylogenetic series is sufficient to render it exceedingly probable. That the primitive diprotodont incisor formula was at least $\frac{5}{5}$ is also strongly indicated. Hence the main points in the present connection seem fairly certain, that is, that primitive diprotodonts had numerous teeth and that the diprotodont modification may have arisen prior to any reduction from the primitive number.

UPPER INCISORS.

Description.—The median incisors are set near the end of the premaxillae in nearly terminal position, only a slight shelf of bone extending in front of them. Their alveoli are more elevated than those of the succeeding teeth and they are separated from the next incisors by a slight space. They are inclined forward slightly away from the perpendicular and their points exceed the lateral incisors. They are separated at the base sufficiently to permit the insertion of a needle or bristle but their points are closely in contact. Their cutting edges are beveled and receive the extreme tips of the lower incisors. They are closely similar in shape to corresponding teeth in the Australian diprotodonts, especially some of the smaller macropods. They have relatively broad faces which suggest the undifferentiated median incisors of *Perameles* quite as much as they do those of the didelphids. The median upper incisors of typical polyprotodonts (dasyures and didelphids) are usually slender and peglike and in most cases the tips are not in contact although occasional specimens may be found having some resemblance to those of *Cænolestes*.

The three succeeding incisors (second, third, and fourth) are wholly lateral in position and the series on one side is almost exactly parallel

with that on the other, a condition approached in no polyprotodonts except *Myrmecobius* and *Perameles*. The second and third are in contact with each other, the second being somewhat larger than the third. They are laterally compressed into relatively broad blades, their cutting edges are truncate and their bases narrow. The blades, which are distinctly hatchet-shaped, are somewhat produced anteriorly. These teeth, although small, are completely functional since they engage the sharp, beveled, outer edges of the median lower incisors with which they have a very effective shearing action. Similar action is seen between the incisors of some of the macropods, and especially of certain of the Phalangeridae, as for example, *Dactylopsila*, in which it is very pronounced. Somewhat similar relations occur in *Perameles*.

The fourth upper incisor is abruptly smaller than the preceding ones and is usually separated from them by a slight space. In superficial view it appears to be pointed but closer examination shows that this is due to a partial forward rotation which has elevated the posterior angle of its cutting edge. This cutting edge is thus inclined forward so that it is parallel to the unbeveled edge of the lower incisor and in this relation it is possible for it to be slightly functional.

Discussion.—All the upper incisors are definitely adapted for relation with the long procumbent lower incisors. The result is the modification of the lateral upper incisors from simple seizing or grasping teeth to cutting or shearing blades. Such a modification, as shown by Bensley, is easily derived from the conical tooth by lateral compression accompanied by the movement of its anterior edge to a horizontal position and its posterior edge to a vertical one continuous with the root. The original apex of the tooth thus becomes the posterior angle. In less marked degree this change has been noted and figured in *Myrmecobius* and *Perameles* by Bensley (1903, p. 105) who speaks of it as a "curious appearance, due to a subcaniniform modification of their tips," and makes a rather obscure comparison with a condition in the degenerate teeth of the Madagascan viverrine *Eupleres*.

In viewing these teeth comparatively in *Myrmecobius*, *Perameles*, and *Cænolestes*, one is at once impressed with the practical identity in form and function in all three forms and it is evident that the modification of the lateral upper incisors is correlated with the elongation and specialization of the median lower incisors. In other words, this form of lateral upper incisor is an indication of beginning diprotodonty and is only found in the Australian diprotodonts and forms conceivably ancestral to them. This subject is discussed elsewhere. It may only be said further that the reciprocal relations of the median lower incisors and the lateral upper ones are exceedingly well-developed in *Cænolestes*.

In fact they appear so efficient for cutting that their relation to the animal's insectivorous habits is not obvious. Functionally, they are quite as well adapted for cutting vegetation as those of strictly herbivorous forms and the possibility that they may have passed through a herbivorous or partly herbivorous stage is to be considered.

LOWER INCISORS.

The median pair of lower incisors are long, slender, and only slightly curved, merely continuing the inferior outline of the mandibular ramus. They are strongly beveled on the upper side anteriorly and their outer edges are sharp and almost bladelike. They are separated from each other by a considerable space and in this and in the other foregoing respects they are closely similar to the lower incisors in the macropods. Their inner edges are not highly developed for cutting and seem to show a slight tendency to imitate even this character of the macropods. The lower incisors of most phalangiers are close-set and decidedly up-curved. Among those examined, the only form showing general resemblance to *Cænolestes* is *Petauroides*.

Of the small unicuspidate, single-rooted, "intermediate" teeth situated between the terminal incisors and the two-rooted premolars, there are two, and in some cases three, which are to be regarded as incisors. They are indistinguishable by size or shape from the supposed canines and anterior premolars except that the foremost one shows an extreme of pronation which is slightly less pronounced in the succeeding ones. Their crowns are turned forward and lie overlapping each other, the first being in contact with the base of the median incisor and the second with the upper exposed surface of the root of the first. Thus the original apex of each tooth is directed forward and the posterior side of the tooth including a part of the exposed root has come into horizontal position. This arrangement appears to be unique among living marsupials although slight approaches to it are seen in some of the Phalangieridae. It is duplicated in every feature, however, in the extinct Patagonian *cænolestids*.

CANINES.

The upper canines in *Cænolestes obscurus* and apparently in most specimens of *C. fuliginosus* are simple slender prongs slightly more curved but otherwise similar to the canines of other omnivorous or carnivorous marsupials. In the male they are much more prolonged than in the female, the exposed part measuring 2.5 mm. in a male and only 1.3 mm. in a female. In a specimen of *C. fuliginosus* figured by Bensley (1903, pl. V, fig. 38) the canine is shown to have a deep lateral

groove extending from the middle of its outer surface to the broad single root. In the closely allied genus *Orolestes* recently described by Thomas, the canine is said to be double-rooted, and shaped like a premolar. Hence we have in the cænolestids now living a graded series showing a change in the character of the canines from a two-rooted premolariform condition to a single-rooted caniniform condition. That this is the direction of the change and not vice versa seems fairly clear. In the first place, the two-rooted condition is the more primitive. Moreover, although it might be reacquired through "retrogression," as held by Bensley, it is very unlikely that the reversal would be represented by these particular stages. Thus, while the grooved root is the obvious intermediate state in change from a double to a single root, it is scarcely what would be expected if the direction of change were reversed. The logical intermediate stage, if we were dealing with a retrogression from a single to a double-rooted condition, would be a lengthening of the crown with possible appearance of antero-posterior cusps before tendencies to division of the root began. Hence it seems that the change in this case is toward rather than away from a long single-rooted piercing type of canine. It is plainly a very recent specialization and is quite in accord with the animal's present insectivorous and predaceous habits. Of still further interest in this connection is the fact that this change has progressed farthest in the northernmost form, that is, the one presumably most distant from the parent stock.

The lower canines have no intrinsically distinctive characters but are quite like the incisors preceding them upon which they lie in semi-prostrate position.

UPPER PREMOLARS.

The anterior premolar is small, greatly compressed laterally, and its crown in side view is nearly triangular, its anterior outline being practically straight and its posterior slightly concave. In wholly unworn condition, traces of a posterior cusp are evident but none of an anterior one. It is situated about midway between the canine and the middle premolar and is separated from each of these by a space equal to approximately one and one-half times its own length.

The middle premolar is somewhat larger than the anterior, but is similarly compressed. Its posterior outline is concave and sweeps backward to a definite posterior cusp. It is separated from the posterior premolar by a slight space about half as long as that between it and the anterior premolar.

The posterior premolar, which is probably a replacing tooth, is separated by a slight space from the middle premolar. It is situated

somewhat obliquely with its small anterior cusp slightly internal to its high triangular posterior blade. Posteriorly it stands closely against the first molar and, although its principal cusp is higher than any of the molars, it is functionally a member of the molar series and fairly well separated from the anterior teeth as in diprotodonts generally. It is slightly thickened posteriorly but has no internal cusps although the cingulum is well-developed. Of its two roots, the anterior one beneath the small antero-internal cusp is the longer and heavier.

UPPER MOLARS.

Description.—The first and second molars are essentially alike in structure and size, the first being only very slightly longer relative to its width. They are quadrate and have four principal cusps, two inner and two outer, i. e., paracone, metacone, protocone, and hypocone.¹ At the inner base of the metacone is a small but high and very distinct intermediate conule which in worn teeth is sometimes obliterated. In lateral view, the paracone and metacone appear as nearly equilateral triangles rising from the cingulum. The angle between these two cusps is more acute in the first molar than in the second and both cusps of the first are slightly higher than those of the second. The protocone is well developed and more thickened but is somewhat lower than the outer cusps. At its antero-internal base it is distinctly bulging. A distinct internal cingulum bounds its base and extends to the anterior base of the hypocone. The hypocone is about half as high as the protocone and its apex scarcely reaches the height of the intermediate conule, so it is much the smallest of the four principal cusps. Protocone and paracone are connected anteriorly by a commissure which forms the front boundary of the deep central depression of the tooth. A low commissure connects the protocone and the hypocone and near the middle of this is a slight ridge somewhat imperfectly connecting the posterior base of the protocone with the intermediate conule. Between the hypocone and the intermediate conule there is an open channel to the back of the tooth.

The third upper molar is subtriangular and the hypocone is practically undeveloped, although a very slight shelf exists between and slightly posterior to the intermediate conule and the protocone. The first, second, and third molars have three strong roots each and these are situated respectively beneath the protocone, paracone, and metacone. The external cingulum in all these teeth is well-marked, but shows no differentiation into styles. The fourth molar is very small, scarcely

¹ These terms and others are used without reference to homologies to designate the cusps to which they have been commonly applied in connection with the Cope-Osborn tritubercular theory.

one-sixth as large as the third, and it is turned slightly inward from the toothrow. It is roughly triangular and carries three small cusps apparently representing paracone, protocone, and metacone. They are scarcely higher than the commissures which connect them and inclose a central depression. There is still retained an external cingulum, a minute anterior shelf, and two roots, one of which is larger and obviously formed by coalescence. This tooth is plainly undergoing reduction and is in about the same stage as the fourth molar of *Dromicia*. In *Acrobates* and *Distaechurus* it has quite disappeared.

Discussion.—The upper molars of *Cænolestes* show an obvious resemblance to those of the diprotodont family Phalangeridae. Two principal types of molars are found in this family: (1) the bunoid and (2) the selenoid, respectively characterizing the two subfamilies Phalangerinae and Phascolarctinae. In general, the bunoid group are without styles or intermediate conules while the selenoid group usually have at least traces of both styles and conules, but in these respects there are exceptions in both groups.¹ Several theories have been advanced as to the origin and mutual relationships of the two groups. Winge considered them derivative the one from the other and regarded *Phascolarctos* ancestral to the bunoid Phalangerinae with *Pseudochirus* as a connecting form. On the other hand, students of placental molars have usually inclined to the belief that, among ungulates especially, bunoid types were ancestral to selenoid. Bensley, however, comes to the conclusion that "the molars of the Phascolarctinae and Phalangerinae have been derived by a divergent evolution from a common insectivorous secodont type," that is, independently from a tritubercular or post-tritubercular type similar to that seen in the Peramelidae. This divergent evolution took place, according to Bensley, as follows:

1. The selenoid type evolved by the simple addition of a hypocone and the reduction of the styles, no change in the shape of the principal cusps being required since they are already selenoid in the tritubercular molar. This method was recognized also by Winge. For the production of the molar of *Pseudochirus* the addition of the intermediate conules was further required and that these were secondary elements which arose de novo seems to have been taken for granted, although, as shown later, there is some evidence that they are not.

2. The bunoid type evolved also by reduction of styles but accompanied by a lowering of the principal cusps and a gradual change in their form from crescentic to subconical. These and other changes involved

¹ In the Phalangerinae, traces of a metaconule occur in *Petaurus* and a large persistent style is present in *Dactylopsila*. On the other hand, the Phascolarctinae include some species of *Pseudochirus* in which styles are lost and *Phascolarctos* in which conules are scarcely evident.

with them are foreshadowed in the molar of *Philander*. As in the selenoid group the quadrate form was produced by the addition of a hypocone.

It is in their relation to these theories of the origin of bunodont and selenodont molars and also to some of the still more debatable theories of the general evolution of mammalian teeth that the molars of *Cænolestes* must be viewed. The essential difference between the bunoid and selenoid groups of *Phalangeridae* is in the shape of the outer labial cusps of the upper molars, these being in the one convex in their external aspect and in the other concave. Judged by this character alone, the molar of *Cænolestes* at once classifies itself with those of the bunoid *Phalangerinae*. Among these, however, it stands out as a relatively primitive type. This is evidenced by (a) the high lanceolate labial cusps, (b) the difference in height between the labial and lingual cusps, (c) the practical absence of a hypocone in the third molar (d) the presence of a distinct metaconule in the first, second and third molars.¹ When, in addition to these characters of the molars themselves, the primitive nature of the antemolar formula is considered, it is clear that there are strong reasons for considering the molar of *Cænolestes* as possibly prototypal to those of the *Phalangerinae*. It is of course conceivable that it is now becoming specialized in an insectivorous direction secondary to a previous bunodont stage. This possibility is suggested by the cursorial adaptations in the skeleton of *Cænolestes*, but, as shown by the *Peramelidae*, these may accompany relatively slight modifications of molar structure.

Assuming that the molar of *Cænolestes* represents a late stage in the development of bunodontology, it becomes difficult to harmonize it with current theories. Such a molar would scarcely be produced by the lowering and thickening of cusps which appear in *Philander*. These changes are only relatively slight deviations from the conditions prevailing among all *Didelphiidae*. They are seen to some extent also in *Didelphis* where, as in *Philander*, they are probably related to habits which are more frugivorous and omnivorous than those of *Peramys* and *Marmosa*. It is conceivable that they might have led either to the bunoid or the selenoid section of the *Phalangeridae* or to some of the conditions seen in the *Peramelidae*. But they cannot be accepted as any indication of the genesis of the quadrate bunolophodont molar of *Cænolestes*.

How, then, was this molar derived? Considerable evidence points to a possible evolution by persistence of styler elements rather than by their reduction and total elimination. The theories of Bensley and others, with the exception of Winge (1882), invariably give a minor role

¹ The interpretation of the metaconule as primitive is discussed on a later page.

to the styles, although they are so highly developed in the tritubercular molar. They are supposed to grow up from the cingulum and then to shrink back into it to complete extinction. In some cases this may occur, especially in placentals, but in the marsupials there is much to show that the styles have played an important part and that they may be the principal elements preserved in the outer cusps of the molars of *Cænolestes* and the bunoid *Phalangerinae*. In some cases they may have wholly supplanted the paracone and metacone and in others they may have fused wholly or partly with them, the resulting cusp usually preserving more of the character of the style than of the so-called primary cusps. This hypothesis favors Winge's theory that the styles are the original elements of the molar crown persistent from the triconodont stage, but it is only facts bearing on the evolution of the bunoid molar from the tritubercular that are of present importance. Some of these are the following:

1. The styler elements of the molar crown are deep-seated structures, being recognizable in many early mammals and in developmental stages of recent ones.

2. They occupy the outer or labial side of the teeth which is phylogenetically older than the inner or lingual.

3. They are well developed in polyprotodont marsupials and show a high degree of consistency in general relations (Bensley, 1906).

4. Morphologically they are bunoid, at least conical or compressed conical in section.

5. The cusps of the trigon in the tritubercular molar are selenoid, essentially alike and essentially different from the styles.

6. Hence the crown of this molar consists of two elements one of which is fundamentally bunoid and probably primitive while the other is selenoid and secondary.

7. The styles are situated directly over the buccal roots and in line with the primitive longitudinal axes of the premolars in the identical position of the outer cusps of the quadrate bunoid molar.

8. On the contrary, the paracone and metacone are midway between the lingual and buccal roots and in order to become the outer cusps in the bunoid molar would require not only a change in form but a migration from their position.

9. Aside from the confusing factors introduced by minor adaptations, therefore, the general relations in the tritubercular molar are favorable for positive participation of styler elements in the formation of the bunoid molar.

With these broader facts in view, evidence may be sought as to detailed methods of change other than that of downward reduction

and elimination of styles. The high bladelike cusps in the molar of *Cænolestes* suggest an evolution along lines similar to those which have produced the carnivorous dentitions. In the Dasyuridae, the carnivorous evolution obviously has proceeded by lateral compression and fusion of styles and cusps. In this process it is difficult in all cases to demonstrate to what extent one element dominates the other, but the styles are quite as consistent as the so-called cusps of the trigon. In *Sarcophilus*, the style dominates the paracone in the first and second molars and the two are evenly divided in the third. Metacone and style, on the other hand, are evenly divided in the first molar, while the metacone dominates in the third. The character of the outer aspect of the teeth in all cases seems determined by that of the styles. In *Thylacinus*, styler elements persist in reduced condition anteriorly and posteriorly but in the intervening region have completely fused with the metacone, doubtless somewhat after the manner indicated in *Sarcophilus*. Even after these extensive changes, the resulting cusp, which is derived principally from the metacone, is situated in the same relation to the roots as the metacone of the tritubercular molar of the didelphids and dasyurids. In the carnivorous evolution, therefore, although the tendency to consolidation of cusps is very great, the styles have been reduced only when their corresponding cusps were reduced and in cases of fusion they have given their character to a large part of the combined product.

In the formation of the molar of *Cænolestes* with its decidedly convex labial surfaces it seems probable that fusion of styles has played quite as large a part as in carnivorous forms. In the bunoid Phalangerinae it is possible that fusion and reduction may have been combined in some cases. As compared with the upper molars of other living marsupials, those of *Cænolestes* are unique in the possession of a marked undifferentiated external cingulum. It forms a continuous faintly crenulate ledge along the bases of all the molars. A similar structure is found in various degrees of perfection in the extinct cænolestids but nothing exactly comparable occurs in modern forms. It is not necessarily evidence of reduction of styles and may easily have been developed independently. In fact an exactly analogous structure is found at the inner base of the protocone of the first upper molar and cingula of the same character appear on the lower molars to which no suspicion of reduced styles attaches. The irregular swelling at the bases of the upper molars in primitive Phalangeridae, as *Dromicia* and *Petaurus*, is regarded by Bensley as a vestigial external cingulum which "appears to be equivalent to that bearing the external styles in polyprotodont forms." In a certain sense this is apparent enough, but I believe that at most only the minor styler elements are represented and that the principal

ones (styles *b* and *c*) are no more reduced than they are in the anterior molars of *Sarcophilus* in which there is a similar slight basal ridge combined with large obvious styles.

Evidence of combined fusion and reduction is seen in the first upper molar of *Dactylopsila*. In this the quadrate form is complete so far as the principal cusps are concerned, but there remains on the antero-external border of the tooth a single well-marked style, which is well distinguished from the paracone. Apparently the other styles have fused, but this one has changed only little and that by reduction. It is to be noted also that this style retains a primitive relation to the root. Most interesting is the fact that the antero-external surface of the paracone, that is, the part opposite the style, is concave or semi-selenoid while its postero-external surface is convex like that of the succeeding molars. In other words, it seems that where fusion takes place, the outer surface of the style is added to the cusp making it convex and where fusion does not take place, the outer surface remains concave as at first.

To produce a bunoid molar from a tritubercular one by this method requires less change than by reduction of styles. In the primitive Dasyurinae and in *Marmosa* and *Peramys* the styles are very well-developed and their convex external surfaces and sublanceolate form closely parallel the conditions in *Cænolestes* and some of the primitive Phalangerinae. Moreover, the paracone in these forms is obviously derived from the styles, in most cases being subordinate to them and in none reaching a stage of development which overshadows them. In the first upper molar of *Phascologale cristicauda* the paracone is not yet differentiated and in the corresponding tooth in *Dasyurus viverrinus* nearly the same condition obtains. This tooth, on account of its anterior position, is probably more primitive than those posterior to it. In these forms, its antero-external cusp has all the characteristics of a style (style *b*). A very slightly bifid tip indicates the beginning of the separation of the paracone and its complete differentiation is seen progressively emphasized in the succeeding third and fourth molars. This first upper molar in the dasyurids is at least superficially similar to the deciduous molar of the didelphids in which, as suggested by Bensley, the antero-external cusp appears to be the result of fusion of style and paracone and may be retrogressive. However this may be, the tendency to persistence of the styles is evident and the conclusion seems justified that the same elements are still represented in the quadrate bunoid molars of diprotodonts. If this be true it is possible that the small metaconule in the molar of *Cænolestes* is not a new element but a persistent unfused vestige of the metacone of the ancestral tritubercular tooth. This at first seems improbable on account of the large size and

functional importance of the metacone in polyprotodonts. Gregory (1916, p. 248) has suggested even that the metacone in these forms represents the apex of the original two-rooted tooth. It is to be noted, however, that with the reduction of the metacone spur and the growth of the hypocone the relative importance of the metacone decreases and in some cases (e. g. *Dactylopsila*) the postero-external cusp, presumably the metacone element, is exceeded by the hypocone which was non-existent when the metacone dominated the tritubercular crown. In *Phascologale*, *Antechinomys* and other generalized dasyurids, the stylar element (style *c*) is very well-developed and only slightly exceeded by the metacone. According to Sinclair, style *c* is well-developed also in *Microbiotherium tehuelchum* of the Patagonian Miocene. To totally eliminate this style and change the selenoid form of the metacone to bunodont, meanwhile transposing the relations of the two to the root, would be a much more extensive process than fusion of style and metacone or even more than increased growth of the style and elimination of the metacone. Moreover, it would be contrary to general evidence on cusp migration, most of which is negative (Gidley, 1906; Tims, 1903).

LOWER PREMOLARS.

Description.—The single-rooted, unicuspid, anterior lower premolar lies in series with the canine and the lateral incisors and is essentially like them in size, form, and implantation. It is separated from the middle premolar by a slight space. In the abnormal specimen having five unicuspid, this space is unaffected, indicating that the extra tooth is an anterior rather than a posterior one.

The middle premolar is two-rooted and tricuspid but it is in a rather advanced stage of reduction toward the unicuspid condition. The principal or middle cusp has taken an anterior position directly above the anterior root leaving a long sweeping curve between it and the posterior cusp. The anterior cusp is practically eliminated, although its position is still indicated by the angle between the root and the front of the main cone. A further reduction involving the loss of the small posterior cusp and root would produce a tooth like the unicuspid anterior premolar.

The posterior lower premolar, like the corresponding upper tooth, is functionally associated with the molars rather than with the anterior premolars. A slight space separates it in front from the middle premolar but posteriorly it is closely engaged with the front of the first molar. It rises considerably above the anterior teeth and even exceeds slightly the level of the molars. It has one large upright cusp rising above its anterior root and a short depressed heel which is over-

lapped by the anterior shelf of the first molar. Thus, although higher, it is slightly shorter than the middle premolar. On its antero-external surface is a faint indication of a cingulum.

Discussion.—The lower premolars, like the upper ones, show tendencies toward the condition prevalent in diprotodonts. This is evidenced principally by the reduction of the anterior teeth and the separation of the posterior one from them to a position at the head of the row of grinding teeth. The only approach to this among polyprotodonts is seen in the Peramelidae.

LOWER MOLARS.

Description.—The first lower molar has a relatively small trigonid and an enlarged talonid. The protoconid, which is only very slightly larger and higher than the other cusps of the trigonid, is connected by crests with the paraconid and metaconid. The talonid is broad and extended and the lateral angle between the protoconid and the entoconid is relatively shallow. No hypoconulid is distinguishable. A distinct external cingulum is present and continues forward to form a slight anterior shelf. The second lower molar is similar in general to the first, but is larger and its trigon is relatively wider, although still somewhat narrower than the talonid. There is a slight indication of a hypoconulid and the paraconid is flattened or even emarginate anteriorly where it meets the entoconid of the preceding tooth. The third molar is smaller than the second and approximately equal to the first. The talonid exceeds the trigon in width only very slightly and the general outline of the tooth from above is rectangular. The flattening of the paraconid is more pronounced and almost divides it into two distinct tubercles.

The fourth molar is compressed and greatly reduced in size, being narrower and only slightly longer than the trigon of the preceding tooth. The metaconid persists nearly as large and well-differentiated as in the other molars but the other cusps are rudimentary. The entoconid is slightly elevated and distinguishable; the hypoconid is represented merely by the postero-internal angle of the short blunt talonid; the protoconid has disappeared; and the imperfectly divided paraconid is vestigial. Despite its small size, this tooth is two-rooted, agreeing in this respect with the extinct form *Halmariphus*, but disagreeing with *Garzonia*.

Discussion.—The lower molars, as compared with those of primitive polyprotodonts, show three conspicuous tendencies.

(1) The reduction of the protoconid and the general reduction in size and height of the outer cusps as compared to the inner ones. (2) The elevation of crests connecting all the cusps. (3) The production of a

subrectangular tooth by the extension of the talonid and the antero-posterior compression of the paraconid. These characteristics are seen to a somewhat lesser extent in the Peramelidae and in general they lead to conditions found among diprotodonts. The talonid still remains slightly wider than the trigon, the individual cusps are still distinctly elevated above the crests, and the molars as a whole are relatively long and narrow. These appear to be primitive features and it is evident that the lower molars are less advanced than the upper. Among the extinct *cænolestids*, as noted by Sinclair, some are even slightly more primitive in these respects than *Cænolestes* while others show increased development of transverse crests making the evidence quite clear for the derivation of buno-lophodont types from tuberculo-sectorial.

THE ORIGIN OF DIPROTODONTY.

The diprotodont modification of the antemolar teeth has arisen in so many diverse groups of mammals that it may well have had different causes and courses of development in different cases. In some, however, at least the general factors involved must have been the same. It occurs in multituberculates, modern marsupials, insectivores, primitive edentates, rodents, ungulates, bats and the lemuroids. In its most specialized condition it consists in the complete suppression of all the antemolar teeth except one pair of incisors in each jaw, these being functionally modified and situated terminally. Among living marsupials, the progressive steps of this modification are illustrated by a series which is so nearly complete that, except for the incipient stages, it could scarcely be more convincing if it were in reality a linear phylogenetic series, which of course it is not. Of living forms in which diprotodonty is unmistakably developed, *Cænolestes* undoubtedly has the most generalized dentition. From *Cænolestes* to the more primitive phalangers, as *Dromicia* and *Distachurus*, the step is very slight; thence to the herbivorous phalangers of the genus *Trichosurus* and on through *Hypsiprymnodon* and *Bettongia* we come to the more typical macropods; and in *Phascodomys*, although in other respects it is not a true terminal member of this series, we see diprotodonty highly developed. This has been traced in great detail and described by Bensley (1903) in the important paper to which reference is so frequently necessary. Beginning with *Cænolestes*, the gradation from one form to another is almost as complete as could be desired.

The idea that *Cænolestes* is not a true diprotodont, but has only paralleled the diprotodonts in dentition, need not here be considered at length, since it is only dentition that is important in this connection

and so far as that alone is concerned there could be no possible objection to considering *Cænolestes* directly prototypal to the Australian diprotodonts. It may be said, however, that the ancestor of diprotodonts by merely a priori reasoning would at some point have the general characters of *Cænolestes*, that is, it would have a diprotodont dentition while retaining a number of polyprotodont characters. This will be discussed elsewhere, but in the series of diprotodont dentitions we can at least place that of *Cænolestes* as the first and most generalized among living forms.

The uncertainty lies in the steps preceding the development of the cænolestid type of dentition, for the gradations between generalized polyprotodont dentitions and the cænolestid stage of diprotodonty are not recognizable with certainty among known living or extinct forms. Hence the most constructive stages in the development of diprotodonty in this case must be supplied theoretically. Bensley, whose opinion is most important, states that "the diprotodont modification, although characteristic of the herbivorous section of the Marsupials, is the result of an insectivorous adaptation which must have been developed in the minute ancestors of the Phalangeridae during the incipient stages of the omnivorous evolution, but after the separation of the peramelid stem." He therefore finds the beginnings of diprotodont tendencies in small polyprotodonts such as *Phascologale* and *Peramys* which have undifferentiated lower incisors but show a slight differentiation of the median upper ones. Since primitive diprotodonts like *Dromicia* have upper incisors of this same character he concludes that such incisors are part of the course toward diprotodonty. With enlargement of the median lower incisors in a form like *Phascologale*¹ he would expect a shortening of the lower jaw and consequent disturbance and reduction of the "intermediate" antemolar teeth. These conditions he finds exemplified in *Dromicia* and *Cænolestes*. This explanation by Bensley may be the true one, for it is difficult to gainsay; but if applied to *Cænolestes* it practically assumes a direct didelphid ancestor and therefore needs careful examination. Before granting that diprotodonty arose exclusively in this manner, it may be inquired whether there is any other way in which it might have originated among marsupials. The didelphids and cænolestids were already well distinguished in the Patagonian Miocene. Hence the origin of the diprotodonty of *Cænolestes* is still farther back, perhaps in the Mesozoic, and it is certain that the ancestral form was not the present day didelphid but an ancient generalized type, all the characters of which are not preserved in any recent forms.

¹ A much greater development of the anterior incisors is found in *Oxygomphius frequens*, a European Miocene form usually referred to the Didelphiidae.

Besides the didelphids and dasyurids there are at least two other groups which may have retained some of the characters of the generalized marsupial and which may throw some light on the origin of diprotodonty. These are the Myrmecobiidae and the Peramelidae, the former group with only one living species and the latter with a considerable number. *Myrmecobius*, as is well known, has a very remarkable dentition combining an excessive number of molars and a relative degeneration of all its teeth. This apparently incompatible condition has been explained in at least three ways. Owen, Thomas, Leche and others believed it to be a primitive survival from ancestral forms having numerous post-canine teeth such as *Dryolestes* of the North American upper Jurassic. Winge has advanced the hypothesis that it is due to the retention of teeth originally belonging to the deciduous series. In this he is followed by Gidley (1915), who points out that "it requires but the addition of two more permanently retained deciduous molars to equal the greatest number of post-canine teeth found in this species, namely, nine." A third explanation is offered by Bensley who believes that the extra molars of *Myrmecobius* are due to a "simple reduplication of teeth from the posterior position of the dental lamina" induced by "the favorable conditions of increased space in the molar region." This conclusion is obviously influenced by the reduced incisor formula of *Myrmecobius* which he believes is derived through the dasyurids from the didelphids which have a more primitive number of incisors. But he fails to take into account the fact that similar reductions of incisors have taken place in the early history of marsupials in Tertiary or even Mesozoic times.

To the writer, these different explanations seem equally theoretical and the original one of Owen, if stated in general terms, is not less probable than the others; that is, the resemblances between *Myrmecobius* and Mesozoic forms seem quite as great as those between it and the modern forms and, since other considerations point to great antiquity for several different types of marsupials, there would seem to be no great improbability in the belief that the extra molars are a survival rather than a modern development.

At the present time *Myrmecobius* is obviously in process of losing rather than gaining teeth. This has been noted by Gidley who says: "The variability in the molar series in *Myrmecobius* seems due to the presence or absence of the last molar, probably a disappearing tooth. In the upper jaw the last molar seems to be normally wanting, while the second is apparently in the process of disappearing, being sometimes present and sometimes wanting." It seems highly improbable that the causes which originally brought about an increase in the tooth formula are the same as those now producing a decrease. A more reasonable

assumption would be that the increase took place before degeneration of individual teeth began and for a different reason. The process of reduction now beginning in *Myrmecobius* may perhaps be compared to the more advanced condition in *Tarsipes* which retains but three or at most four post-canine teeth and is evidently well on the way to an edentulous state.

Gidley makes the unequivocal statement that "There is in our present knowledge nothing to support Owen's hypothesis regarding the derivation of *Myrmecobius*." But in succeeding pages of the same paper he argues for an ancient origin of the Myrmecobiidae, which he supposes to have been well differentiated from the didelphids and dasyurids as early as the Paleocene. Evidently he believes in a Mesozoic ancestor for *Myrmecobius* and to that extent at least agrees with Owen. Whatever the case may have been with respect to the history of the extra molar teeth of *Myrmecobius*, the view that all the living families of marsupials were well differentiated early in the Tertiary seems to be well founded. Therefore, without reference to possibly archaic characters other than teeth, it is still possible to believe in an early predidelphid origin for *Myrmecobius*. As a possible forerunner of diprotodonts, the *Myrmecobius* line is thus entitled to some consideration. It has the same antemolar formula as *Cænolestes*, its upper canines are somewhat reduced, and its median lower incisors are greatly specialized and relatively larger than in any other polyprotodont. These incisors are completely terminal in position and, although slightly recurved at the tips, are essentially proclivous with their roots nearly in the principal longitudinal axis of the mandible and set at a decidedly different angle from the lateral incisors. Thus the median lower incisors of *Myrmecobius* are essentially like those of diprotodonts. If these teeth are as much reduced as the rest of the dentition, they must have been derived from a type which might easily have led to the diprotodonts. The median upper incisors of *Myrmecobius* are reduced in size and slightly smaller than the lateral ones. They are set very far apart and are slightly procumbent, obviously in a secondary condition which differs from that of polyprotodonts and diprotodonts alike. Hence no conclusion is to be drawn from them although it is to be noted that the median upper incisors are widely separated at the base in diprotodonts more generally than in polyprotodonts. The first pair of upper lateral incisors are modified to meet the median lower ones as in diprotodonts and all the lateral incisors are laterally compressed. The differentiation of the median upper incisors is scarcely to be regarded as a diprotodont character for it is not pronounced except in a few extreme cases, as *Phascolarctos* and *Phascalomys*, and in many it is distinctly less than in

some polyprotodonts. In *Trichosurus*, for example, and in *Cænolestes*, the median upper incisors are less differentiated than in some species of *Phascologale*.

For the development of diprotodonty from such a form as *Marmosa* the following are among the changes which would be necessary: Reduction in number of incisors; enlargement of median lower incisors; reduction in size of canines and anterior premolars; decrease in the depth of the mandible; increase in craniofacial length; formation of short diastemmata between at least some of the teeth; assumption of proclivous and fully terminal position by median lower incisors; assumption of wholly lateral position by posterior incisors; lateral compression of incisors and development of long and forwardly directed cutting edges, especially in those of the upper jaw. All of these modifications are found to a considerable degree in *Myrmecobius* and in these respects this form resembles the diprotodonts more closely than do any of the didelphids or dasyurids. Whether these characteristics were more or less pronounced before the present degeneration of teeth began cannot be said, but at least it must be conceded that *Myrmecobius* has some claims to a position in or near the line of the diprotodonts whether it be regarded as an ancient type derived from Jurassic forms or a more recent one proceeding from the dasyurids.

Leaving *Myrmecobius* for further discussion later, it may now be inquired whether the peramelids could have furnished the beginnings of diprotodonty. Although Bensley finds in the Peramelidae many indications of an ancestral relation to the modern diprotodonts and an advance over the primitive dasyurids and didelphids, he is not inclined to regard the diprotodont dentition as directly derived from them. In this he was influenced largely by the undifferentiated condition of the median upper incisors in the peramelids; by the retrogressive(?) character of the upper canines and premolars, the existence of primitive and advanced stages of the hypocone in both groups, and the hypsodonty of the peramelids. The development of the hypocone and the hypsodont modifications in the peramelids are progressions beyond the most primitive stages of the known diprotodonts. But they need not be considered wholly homoplastic and their inception may have preceded the divergence of the peramelid stem. In any case they do not preclude a belief in a common ancestry for the peramelids and the diprotodonts and one which was not very remote. Such an ancestry at most need only be removed to a stage in which the hypocone and the hypsodonty were approximately at the most primitive stage common to both peramelids and diprotodonts. This would carry us back to a brachyodont condition but not to a total absence of the hypocone. So of the two characters

involved, only one—hypsodonty—needs to be accounted for. This may well be regarded as a recent specialization for it is of a very peculiar nature, a sort of half-hypsodonty confined to one side of the teeth and not fairly comparable with the full hypsodonty of the most advanced diprotodonts.

The undifferentiated median upper incisors of peramelids and the supposed retrogressive canines and premolars, in the light of a study of *Cænolestes*, do not seem at all unfavorable to a hypothesis that a diprotodont dentition might have proceeded from the peramelid stem. The incisors, while not differentiated for seizing as in the dasyures and didelphids, are unquestionably in a secondary condition of which no further development is seen elsewhere, unless it be among diprotodonts. These incisors have become broad and somewhat bladelike with their cutting edges in the same horizontal plane as those of the adjacent lateral incisors. In fact they are practically indistinguishable in shape from the pair of lateral incisors situated next to them. Hence it seems that the same factors which have led to modification of the lateral teeth, may have affected the median ones. The principal of these factors is the functional interaction with the lower incisors. In *Perameles* the three pairs of lower incisors are collectively so modified in form and especially in position that their functional relation to the upper incisors is much like that of the elongated single pair of lower incisors in diprotodonts, certainly much more so than in any other polyprotodont. They are set so closely together that their cutting edges are practically continuous. In addition they are exceedingly proclivous with their roots almost as near the long axis of the mandible as in diprotodonts. Thus the functional effect of the three incisors in *Perameles* is practically the same as that of the one in diprotodonts and it is probable that specialization in the lower jaw has conditioned that in the upper rather than vice versa.

Another interesting character correlated with the proclivity of the lower incisors in *Perameles* is found in their exposed roots, those of the first and third especially being without bony covering in front. The second incisor is more fully rooted, but the other two, although well-developed, are very weakly attached to the jaw. Perhaps enlargement of any one of the three might lead to the loss or reduction of the others which would result in a diprotodont dentition. This is uncertain, but it is probably fair to assume that the proclivity and the exposed roots of these teeth are indications at least of some instability, possibly following the loss of the fourth pair or foreshadowing a changed relation of those remaining. The reduction that has already taken place in *Perameles* corresponds to the reduction in the diprotodonts, that is, the first pair of lower incisors has been lost and thus one of the stages preliminary to

diprotodonty has been passed. Embryological evidence so far adduced points to the conclusion that this is the case. Woodward's (1893) studies of developing teeth in the Macropodidae led him to believe that the large functional lower incisor is a persistent member of a series in which it formerly stood in second place. Wilson and Hill (1897), from extensive work on the development of *Perameles*, find that the first lower incisor has been lost and that the one now found in terminal position is homologous with the large lower incisor of the macropods. These authors say (p. 509): "It is interesting to compare the condition in *Perameles* with that described by Mr. Woodward for *Petrogale*. In that form he found two small calcified vestigial incisors. The anterior of these was entirely in front of the large permanent lower incisor, which latter he therefore regarded as really i_2 And may it not be that Woodward's second vestigial incisor (his i_3) to which he was unable to recognize any ancestor, is really the homologue of our di_1 whose legitimate successor is Woodward's i_2 , the almost certain homologue of our ' i_1 ' of the adult *Perameles*?"

The dasyurids have the same number of lower incisors as *Perameles*, but whether they also have lost the first pair has not as yet been demonstrated. Embryological evidence in their case seems lacking. It was formerly supposed (Cf. Thomas, 1887) that the dasyures had lost the fourth incisor and therefore that the maximum of three incisors in higher mammals represented the first, second, and third of the primitive series. The evidence for this supposition adduced by Thomas now seems inadequate, but in connection with theories of the origin of diprotodonty it would be important to know whether the dasyurids have lost the same incisor as the peramelids.

The canines and premolars, which Bensley interprets as retrogressive, constitute a similarity between *Perameles* and *Canolestes* and would seem to favor a common ancestry rather than otherwise. Whether they are in reality retrogressive is another question. Retrogression in this sense is a sort of reversed homoplasy which can be recognized only if the immediate ancestry of the form in question is known. That is, in the case of *Perameles*, a didelphid ancestry is essential to a belief in the retrogressive nature of its canines. On the other hand, if *Perameles* had no distinct didelphid ancestry but descended from a more primitive form also ancestral to the didelphids, it may have retained a primitive premolari-form canine. Highly specialized canines appeared very early, in fact they were well differentiated among some early reptiles, and various Mesozoic mammals had simple single-rooted canines, but others, like the Jurassic *Triconodon*, possessed double-rooted or grooved canines similar to those now found in *Perameles*, *Canolestes*, and *Myrmecobius*—

all of which it should be noted are forms having certain primitive characters not retained by the didelphids. Double-rooted canines are not unknown in other groups than marsupials but are very rare except in insectivores, another primitive group. Among these, premolariform canines show no evidences of retrogression. Double-rooted canines are said to occur also in the Miocene pig, *Hyotherium* (Beddard). In *Antechinomys laniger*, one of the most primitive polyprotodonts, the canine is single-rooted but has its crown distinctly premolariform. To interpret this as primitive seems quite as reasonable as to call it retrogressive (see p. 119). Canines with lateral or antero-posterior cusps are found also in certain bats (e. g. *Cynopterus*, *Harpionycteris*). In the Miocene cænolestids the canine is mostly unknown, but in *Paraëpanorthus*, as figured by Ameghino (1903, fig. 62), the canines are small and single-rooted, not unlike those of many phalangers. These teeth may have been derived from double-rooted ones but indicate no tendency toward them. Such a form as *Paraëpanorthus*, however, is evidently farther advanced in some respects than *Cænolestes*, for its incisors are more reduced in number.

On the whole, it appears that double-rooted, groove-rooted, or premolariform canines are found almost exclusively among forms exhibiting various primitive characters. The Peramelidae are primitive in certain other respects and, although it is not unlikely that their canines may have passed through various changes prior to recent times, it seems quite as possible that they may have retained an early condition like that shown in the Jurassic *Triconodon*. The same may be true of *Myrmecobius* and of *Cænolestes*.

The objections to a diprotodont evolution from the peramelids therefore, are not insuperable. In fact it is clear that many of the changes necessary to a transition from a didelphid to a phalanger or cænolestid are already present in *Perameles* and it does not seem demonstrable that these changes took place subsequent to the separation of the peramelid and diprotodont stems. In other words, the early peramelids may have become considerably differentiated from the generalized didelphoid marsupial and later divided into one line leading to the modern peramelids and another to the diprotodonts. At least, the conclusion is unavoidable that among living forms those most suggestive of what the ancestor of *Cænolestes* and other diprotodonts was like, are the Peramelidae. Moreover, if the early Tertiary form *Myrmecoboides* is in reality a "properamelid," as suggested on another page (p. 142), belief in the origin of diprotodonty from this group is greatly strengthened. The following resemblances between *Cænolestes* and *Perameles* are to be noted: Six or more exclusive cranial characters (see p. 110), tooth

formula, molariform teeth with hypocone, lower incisors proclivous, lateral upper incisors compressed, reduced canines and premolars, specialized posterior premolar, movable cervical rib, osseous patella, large median vagina, cursorial structure of legs, first and fifth toes of forefeet reduced, caecum small, tail not prehensile. Many of these characters, while not diagnostic of the diprotodonts as a group are present in the majority of that group and absent in the majority of polyprotodonts.

It may be concluded, therefore, that diprotodonty in the cænolestids may have originated in one of three ways:

(1) from didelphid pre Miocene types through undiscovered intermediate forms existing prior to the differentiation of the peramelids.

(2) from an ancient line now represented by *Myrmecobius*.

(3) from primitive peramelids after their separation from the dasyurid-didelphid stem.

The first of these is the theory of Bensley and in general of those who have followed him. The second is perhaps less entitled to consideration, but is still not wholly baseless. The third is to the writer more probable than either of the others.

RELATIONSHIPS OF WYNYARDIA.

The extinct marsupials known from the Australian region are nearly all of Pleistocene age and unequivocally diprotodont or polyprotodont. The most conspicuous exception to this is *Wynyardia bassiana* Spencer (1900) from beds at Table Cape, Tasmania, which are regarded as late Eocene or early Oligocene. This form, the remains of which include parts of limb bones and an imperfect skull without teeth, exhibits such a curious mixture of characters that it has so far been omitted from formal classifications. Spencer summarizes his study of it as follows: "A consideration of all the features would appear to lead to the conclusion that the fossil is the representative of a now extinct series of forms which were more nearly allied to ancestral Polyprotodonts than are any of the existing Diprotodont forms. It may, in fact, be regarded as intermediate between the former and the latter, and as indicative of a stage in the development of Australian marsupials when the ancestors of the recent diprotodontia were beginning to diverge from the original Polyprotodontid stock from which they have been developed within the limits of the Australian region."

Bensley (1904, p. 200) is somewhat more definite, saying: "While it would be difficult to add to the excellent comparisons presented by Spencer, it is probable that the reference to the animal as an inter-

mediate form must be excluded, if for no other reason, on account of its large size. . . . All the forms which approach the hypothetical intermediate type are of small size. Not only this, but the diprotodont modification itself may, as already explained, be shown to represent an insectivorous adaptation which could only have taken place in comparatively small animals. The relations of *Wynyardia* are more probably with one of the advanced genera, such as *Pseudochirus*, *Phascolarctos*, or *Phalanger*."

Osborn (1907) says: "Among fossil forms the gap between the two suborders is largely bridged over by the extraordinary genus *Wynyardia* of Baldwin Spencer, which presents a perfect mélange of characters seen elsewhere only in the Opossums and Dasyures (Polyprotodonts), and in the Phalangers and Kangaroos (Diprotodonts)." Gregory (1910, p. 214) speaks of *Wynyardia* as "another form that helps to bridge over the structural gap between the Polyprotodontia and the Diprotodontia." Later (Osborn, 1910) it fails to receive formal classification.

Spencer, in summarizing the important characters of *Wynyardia* finds five which he interprets as indicating polyprotodont affinities. These, as numbered and stated by him are as follows:

- "(1) Proportionate length to breadth of the skull, 100:67. This approximates most nearly to *Dasyurus*, and shows a decidedly greater width than in the Phalangeridae.
- "(2) Lambdoidal crest well developed, as in *Dasyurus*.
- "(3) Sagittal crest strongly developed, resembling that of Dasyuridae and species of *Didelphys*.
- "(5) The wide sweep and upward curvature of the zygomatic arches, as in Dasyuridae.
- "(7) The transverse elongation of the glenoid cavity, the downward produced plate of bone which forms the boundary is not connected with any structure forming part of the auditory passage. In this respect it agrees with Dasyuridae and *Perameles*, and differs markedly from the Phalangeridae, amongst which it forms the anterior part of a bony auditory canal."

At least the first three of these characters cannot be regarded as exclusively polyprotodont. Taking them up numerically, it is found as to (1) that, although *Dasyurus* and *Sarcophilus* have very wide skulls, polyprotodonts in general have relatively narrow ones. The ratio runs from only 39.7 in *Perameles* to 81.2 in *Sarcophilus* (see p. 99). On the other hand the average relative width is greater in diprotodonts than in polyprotodonts, the only very narrow type seen among diprotodonts being in the primitive *Cenolestes*.

As to (2), it is to be noted that a high lambdoid crest is no more than a generic character among polyprotodonts. Within the limits of the

Dasyuridae and the Didelphiidae it is extremely variable. Various diprotodonts, including most of the macropods and some phalangers have considerable indications of a lambdoid crest and in *Phascolarctos* it is highly developed. *Thylacoleo* also is described as having a high crest.

As to (3), much the same is true of the sagittal as of the lambdoid crest. In the Didelphiidae we find a high sagittal crest in *Didelphis* and none in the closely related *Peramys* and *Philander*. A sagittal crest of some prominence is seen in the skull of *Trichosurus*. Its presence is less frequent than its absence in both polyprotodonts and diprotodonts and cannot be regarded as of great significance.

Spencer's characters (1), (2), and (3), therefore, cannot be taken as evidence of the polyprotodont affinity of *Wynyardia*.

Characters (5) and (7) seem somewhat correlated and it is evident that in the region of the glenoid fossa and the posterior base of the zygoma we have a more pronounced resemblance to the condition in polyprotodonts than in any living diprotodont with the exception of *Cænolestes*. In the relatively high position of the glenoid fossa, however, there is approach to the phalangers. It seems, therefore, that only one of the supposed resemblances to polyprotodonts is important and this is approximated in the primitive diprotodont *Cænolestes*.

The resemblances to diprotodonts are almost as variable and as difficult to regard as diagnostic, but there are more of them and on the whole they seem more convincing of relationship. This is especially true if comparisons are made only with typical polyprotodonts (as didelphids and dasyurids) and with typical diprotodonts (phalangers), leaving aberrant forms, as *Perameles* on the one hand and *Cænolestes* on the other, out of consideration. Thus the large squamosal is found in *Phascolarctos* much as in *Wynyardia* completely excluding the alisphe-noid from contact with the parietals; and in general the squamosal tends to forward extension among diprotodonts but, while this offers considerable contrast to the condition in the Dasyuridae and Didelphidae, it is fully equalled or even outdone in the Peramelidae. One of the characters suggesting diprotodont relationship of *Wynyardia* is the large premaxilla but here again the peramelids must be ruled out. The relatively large cranium, the high position of the glenoid fossa, the elevated mandibular condyle, the position and shape of the anterior root of the zygoma, the massive ischium, and the proportionately long hind limbs all seem to point to diprotodont affinity in *Wynyardia*. Hence, while it may be agreed with Spencer and others that it combines many features of the two groups, it seems best for practical purposes to classify it with the diprotodonts and in recognition of its various unique features to regard it as the type of a distinct family, the WYNYARDIIDAE.

A comparison of *Cænolestes* with Spencer's figures and description of *Wynyardia*, although rather unsatisfactory as a basis for positive conclusions, reveals a number of points in common. These may be enumerated as follows: (1) Cranial part of skull large as compared to facial. (2) Relations of glenoid fossa to auditory meatus similar. (3) Squamosal not inflated and posterior base of zygoma somewhat Dasyurine. (4) Glenoid fossa relatively high. (5) Nasals extended anteriorly and expanded posteriorly. (6) Premaxillae large and separated from the maxillae by a suture which is only slightly oblique. (7) Mandibular condyle relatively high. (8) Femoral trochanters similar and a third tuberosity present as in *Phascolomys*.

Comparison of *Wynyardia* with extinct American forms allied to *Cænolestes*, as *Palæothentes* and *Abderites*, also shows several interesting points. Thus *Palæothentes* has lambdoid and sagittal crests combined with zygomata of a type similar to those of *Wynyardia* and the frontals are widest in their cranial part. In *Abderites*, as previously noted by Spencer, the seat of the large lower sectorial tooth forms a deep transverse furrow in the mandible which seems closely comparable to the condition in *Wynyardia*.

Present material is not adequate for the positive conclusion that *Wynyardia* is intimately allied to the American forms but the above similarities are most suggestive and must be regarded at least in some degree as strengthening belief in extensive community of relationship between Australian and American marsupials, both polyprotodont and diprotodont.

RELATIONSHIPS OF *MYRMECOBOIDES*.

The genus *Myrmecoboides* was established by Gidley in 1915 for an incomplete lower jaw from the Paleocene (Fort Union) of Montana. It was tentatively classified as belonging to the family Myrmecobiidae and characterized as follows: "Canine semipremolariform, being irregularly triangular in cross section and but slightly curved; canine and the three simple premolars evenly spaced with short intervening diastemae. There is also a short diastema between the canine and i_3 (the position of the other incisors is not known). Fourth tooth behind the canine (probably dp_4 retained) completely molariform; true molars tritubercular, with well-developed basin heel, but with inner cusps of trigonid (paraconid and metaconid) as high or higher than main outer cusp (protoconid)."

The principal characters regarded by Gidley as suggesting possible affinity with *Myrmecobius* are (1) the diastemata between the premolars,

(2) the indicated straightness of the jaw anterior to the canine, (3) the premolariform canine, (4) the compressed premolars, (5) the absence of cingula, (6) the relatively high inner cusps (metaconid and entoconid). He also points out that the extinct form differs from *Myrmecobius* in at least three important particulars (1) the possession of the normal number of post-canine teeth, (2) the approximation of the paraconid to the metaconid, (3) the more elevated and better defined trigonid.

Comparison of Gidley's figures and description with several species of Peramelidae shows at once that practically all the above enumerated points of resemblance are to be found in that family and none of the important points of difference. This is indicated by the following parallel:

Myrmecoboides and *Perameles*

Canine equal to or lower than premolars.
Protoconid and hypoconid well developed.
Paraconid and metaconid approximated.
Trigonid elevated and distinct.
Post-canine teeth seven.

Myrmecobius

Canine higher than premolars.
Protoconid and hypoconid small or rudimentary.
Paraconid and metaconid not approximated.
Trigonid not differentiated.
Post-canine teeth more than seven.

The important points of difference between *Myrmecoboides* and *Perameles* are (a) the unicuspid anterior premolar in the extinct form, (b) the large last molar with its well-developed hypoconulid, (c) the exceptionally large paraconid of the first molar, and (d) the somewhat more brachyodont condition of the molar series. The unicuspid premolar is a specialization such as might be expected in an ancestral diprotodont but the other characters are primitive and doubtless hark back to a tritubercular stage. The resemblances to *Perameles* are pronounced, however, and cannot be overlooked, although the classification of such an ancient form on fragmentary material has much possibility of error. In view of the many similarities to *Perameles* shown by *Cænolestes*, the occurrence in the early American Tertiary of a form with distinct leanings to *Perameles* becomes especially significant. The hypothesis can scarcely be avoided that, provided *Myrmecoboides* is a marsupial, it may well be ancestral to the cænolestids. A point of especial interest and significance is the unicuspid premolar which is paralleled among marsupials only in cænolestids and diprotodonts.

PHYLOGENY AND TAXONOMY.

The great range of variation in structure and adaptive characters shown by existing marsupials makes it possible without reference to extinct forms to construct linear series in which the sequence from

generalized to specialized types is remarkably free from interruptions. If their present distribution also be disregarded, that is, if both American and Australian forms be included, the problem is simplified. As worked out in great detail by Bensley (1903), the American Didelphiidae stand in a prototypal morphogenetic relation to the remaining groups of marsupials. This conclusion is based upon their possession of a combination of various generalized features which are distributed over at least three different Australian groups. They have the primitive tritubercular molar as in the Dasyuridae, the upper incisor formula and upper molar stylar characters as in the Peramelidae, and the pedimanous foot-structure (including incipient syndactyly in one form) as in the Phalangeridae.

As the probable ancestor of the modern Didelphiidae, Bensley finds the Oligocene *Peratherium* fulfilling practically all theoretical requirements including a distribution in both Europe and North America which might permit its dispersal either to South America or Australia or to both. This dispersal he supposes to have taken place in at least three radiations, the first being represented by the Australian fauna, the second by the Miocene fauna of South America and the third by the present day Didelphiidae of South America, this last being in its incipency. The evolutionary series so thoroughly expounded by Bensley is an exceedingly plausible one in spite of its having been based largely upon existing forms. The various modern polyprotodonts are obviously reducible to a common type and, although the diprotodonts offer more difficulty, they may at least be traced from one to another. Between the two groups is a rather definite hiatus but there is little room for doubt that diprotodonts proceeded from polyprotodont ancestors. Indications of some of the steps leading from one group to the other are seen principally in the Peramelidae, the Wynyardiidae, the Myrmecoboididae and the Palæothentidae. No one of these aberrant groups is free from objection as an ideal transitional type, yet taken together they serve to bridge quite thoroughly the gap between the generalized polyprotodonts and the more primitive diprotodonts. Bensley dealt chiefly with modern Australian forms, giving relatively little attention to *Wynyardia* or *Cænolestes*, and *Myrmecoboides* was unknown to him. But the Peramelidae he regarded as proceeding from a hypothetical stem-group called Properamelidae, one branch of which led to modern peramelids and another to the diprotodonts. This group presumably had begun to acquire subquadrate upper molars with incipient hypocones as in modern peramelids but had not developed a terrestrial foot structure. The Properamelidae themselves were supposed to have evolved from the generalized polyprotodont having the characters of the Didelphiidae

and giving rise through another line to the Dasyuridae and modern polyprotodonts. Interesting in this connection is the fact noted by Bensley (1903, p. 111) that the modern peramelids agree with the Oligocene *Peratherium* more closely in the character of the important styles of the molars than they do with the Australian dasyurids.

The fossil record is very incomplete but its testimony accords in a surprising degree with the hypothetical requirements of Bensley's scheme. *Myrmecoboides* might without serious objection be looked upon as in the direct line between the "Properamelids" and the diprotodonts, since it has begun to reduce its premolars, the first being small, unicuspid, and single-rooted as in many diprotodonts. Furthermore, *Cænolestes* shows numerous resemblances to *Perameles*, so many that a common ancestry subsequent to the generalized polyprotodont stage is by no means impossible. The material representing *Myrmecoboides* is too imperfect to warrant more than a provisional conclusion that it may be an early stage in the line of the cænolestids. Nevertheless the evidences of affinity between *Myrmecoboides*, *Perameles*, and *Cænolestes* cannot be overlooked. Moreover, if *Myrmecoboides* is in truth a representative of the early differentiation of a diprotodont stem then it is the first form of the kind to be discovered outside the southern continents and goes far toward explaining the present day occurrence of diprotodont types in both Australia and South America.

Further points of interest in connection with the resemblances of *Cænolestes* to the peramelids are found in the slight indications of common relationship to *Notoryctes*. This peculiar form is so highly specialized that its affinities are very problematical, but Bensley is inclined to regard it as an offshoot of the "properamelids." Certain features of the myology (see p. 60) and osteology of *Cænolestes* and *Notoryctes* are similar and favor belief in a common (properamelid) derivation rather than otherwise. The actual course of evolution must be inferred to a considerable extent even under the best of circumstances. The view that *Cænolestes* is a primitive diprotodont is not proved but is strongly supported by its resemblance to the peramelids which of all polyprotodonts are the ones most suggestive of the incipient stages leading from one large group to the other. It is probably not too much to say that if the cænolestids had been discovered in Australia instead of South America they would have been accepted without question as ancestral diprotodonts. This geographical difficulty is at least partly overcome by the presence of *Myrmecoboides*, also with peramelid affinities, in North America. Hence it is a hypothesis of considerable probability that the cænolestids, like the Australian diprotodonts, developed

from "properamelid" ancestors and that the characters or tendencies fundamentally distinguishing them from polyprotodonts were a heritage which they had in common with the Australian diprotodonts.

Increased knowledge frequently, perhaps usually, leads to increased difficulty in defining taxonomic concepts. This is especially true if it be attempted to express both morphogenetic and phylogenetic relationships in one and the same scheme of nomenclature. The division of modern marsupials into polyprotodonts and diprotodonts was made originally upon a morphological basis and until the discovery of extinct forms, which not only were connectant themselves but drew attention to connectant characters in certain modern forms, the two groups were clearly defined. In the case of the peramelids, a position with the polyprotodonts was generally conceded, notwithstanding their syndactyly which was otherwise characteristic of diprotodonts; *Wynyardia* was left without definite allocation; and *Cænolestes*, as detailed in preceding pages (pp. 6-15), was regarded by some authors as a diprotodont and by others as a polyprotodont.

The foregoing study of *Cænolestes*, while adding much to knowledge of fact, does not make classification easier. It is believed that the cænolestids and the peramelids, possibly with *Notoryctes* also, proceeded from the same stem as the phalangeroid diprotodonts. Hence these might be brigaded as one group. To characterize such a group on the basis of the morphological characters exhibited by modern forms, especially with relation to degree of specialization, however, seems practically impossible. After trying various alternatives and considering the present state of knowledge, no more satisfactory primary division of marsupials offers itself than the old one into Polyprotodontia and Diprotodontia. In such a classification, as already shown, *Cænolestes* and *Wynyardia* may be placed with the diprotodonts. The line must be drawn somewhere and the least objectionable place for it seems to be between the cænolestids and the modern peramelids. The position of *Myrmecoboides* must remain doubtful until better material is forthcoming, but its close alliance with *Perameles* is scarcely to be doubted unless it should prove to have placental affinities. The recognition of a superfamily Cænoles-toidea, as in the classification of Osborn and Gregory, serves to divide American and Australian diprotodonts and has its advantages as a matter of convenience. The modern genus *Cænolestes*, however, need not take separate family rank but may be included in the *Palæothentidae* with the very closely allied extinct forms from the Santa Cruz beds of Patagonia. To rank it as a subfamily, Cænolestinae, would be convenient.




DIAGRAM OF PHYLOGENETIC AND TAXONOMIC RELATIONSHIPS OF PRINCIPAL GROUPS OF MARSUPIALS.

The accompanying diagram (p. 144), like others of its kind, has many shortcomings and some inconsistencies. It is offered only provisionally as an aid in indicating the relative position of *Cænolestes*. Various problems in the phylogeny of marsupials must somehow be met in such a diagram even though they seem properly beyond the scope of the present study. So far as any conclusions are indicated, they are to be regarded mainly as suggestive rather than decisive, as indeed they would be in any case. *Myrmecobius* is separated from the main polyprotodont stem and from the dasyurids in order to indicate its possible direct descent from mesozoic types, but a position nearer the dasyurids is almost equally probable.¹ The problem of the relationship of *Propolymastodon* has not been studied and its uncertain position is merely suggested (see Gregory, 1910, p. 212). The same is true of *Pedimys* and *Didelphops*, while the multituberculates (Allotheria), which Broom contends are monotremes rather than marsupials, are indicated as probably proceeding from the unknown forms indefinitely termed "generalized marsupials" and in reality meaning very little.

DISPERSAL OF MARSUPIALS.

Broadly speaking there are only two theories of the dispersal of marsupials, one that the group originated in Holarctica and spread southward, the other that it had its beginnings in Antarctica, the hypothetical southern continent which formerly may have connected part of present-day South America and Australia. Advancing knowledge has from time to time favored one or the other hypothesis but opinions are still divided. A review of the subject with many references is given by Osborn (1910, pp. 64-80).

Before the discovery of the extensive extinct fauna of Patagonia, the distribution of marsupials offered no serious objection to a theory of northern origin for the group. Practically the only fossils known were didelphids from the Upper Miocene and Oligocene of Europe and North America, and the conclusion was fairly obvious that these might represent the ancestral stock from which both Australia and South America received their marsupial populations. The best known advocate of this theory was Wallace. A reference to his work on the Geographical Distribution of Animals (1876), however, shows at once that the information

¹ Although this disposition of *Myrmecobius* is somewhat heretical, I am led to it through the cumulative effect of encountering in this study and in the published work of others the continual necessity of special theory or explanation to account for first one and then another peculiarity of this animal. Much of this would be obviated by carrying its origin farther back. The possibilities for longer lines of descent than usually assumed is evidenced by *Cænolestes* itself in its persistence from Miocene times to the present in practically unchanged condition.

on which his conclusions were based was vastly inferior to that now available.

Although it may be admitted that some of the Mesozoic mammals were generalized marsupials, it is hardly possible to consider them in efforts to follow the course of differentiation and dispersal of the modern groups. For present purposes the multituberculates also may be disregarded. From the early Tertiary we now know relatively primitive types of marsupials from North America, Europe and South America. Correlation of these different forms has been a matter of much difficulty but according to the most recent authorities, there is little difference in age between the northern and the southern types. Thus in North America and Europe we have the didelphoid forms collectively placed in the genus *Peratherium* and regarded as of upper Eocene and Oligocene age. We also have in North America *Myrmecoboides*, which, although a less primitive form, is found in the Basal Eocene (Fort Union). Of still earlier age, but more doubtful relationship, are *Pedimys* and *Didelphops* from the North American Upper Cretaceous. From South America the most primitive type known is *Proteodidelphys* from the Notostylops beds the age of which has been variously placed from Lower Cretaceous to Upper Eocene. From Australia the earliest known form is *Wynyardia* which is of at least Oligocene age and, although not primitive, has a combination of diprotodont and polyprotodont characters which at this early period is significant.

Subsequent to these primitive types there are three large marsupial faunas the members of which show various similarities to each other and any or all of which may have been developed from the early generalized forms. These faunas are represented by (1) the Miocene fossils from South America including the cæolestids, the borhyaenids and the so-called Microbiotheridae which have didelphid characteristics; (2) the present day Australian fauna with its diversity of specialization; and (3) the present day American fauna all referred to the single family Didelphiidae. The first and second of these faunas, broadly speaking, are mutually representative, that is, the South American cæolestids are at least to some degree counterparts of certain Australian diprotodonts; the borhyaenids stand in the same or even closer relation to the Australian thylacine; and probably it is not too much to say that the Microbiotheridae, in spite of didelphid similarities, are not far removed, structurally, from the generalized Australian polyprotodonts.

This correspondence between the South American Miocene forms and the Australian recent forms has not unnaturally led to the conclusion that a direct interchange between the faunas actually took place

perhaps in Cretaceous times by means of an Antarctic land bridge. Further evidence of the former existence of such a bridge is afforded by plants, mollusks, crustaceans, insects, amphibians, reptiles, and other forms of life of which closely allied species are now found inhabiting both Australia and southern South America. Enumeration of these forms has been made at great length by various authors and need not be repeated. It is sufficient to admit that the evidence is very extensive and if the southern continent be supposed to have been one of no great permanence, but, as Hedley (1895) has indicated, may have been attenuated, irregular and unstable at many points, belief in its existence is greatly strengthened and the objections based upon geological and isostatic data do not weigh so heavily against it. To cite only a few of the believers in Antarctica, it may be said that among its most recent adherents have been Sinclair (1906), Ortmann (1905), Scott (1913), and Osborn (1910). One of its earliest and most distinguished advocates was Huxley. In general it seems to have been conceded that with so much and so varied evidence there could be little doubt that some sort of connection must have existed. The northern origin of many and very important groups, however, could scarcely be denied and the general theory of mammalian dispersal was thus a bipolar theory. Nevertheless the Argentine paleontologist Ameghino went so far as to regard South America as the original home and center of dispersal of practically all the larger groups of mammals.

In contradistinction to these views and in reversal of his own previous acquiescence to belief in Antarctica, Matthew (1915) has presented a powerful argument for the dispersal of mammalian life exclusively from Holarctic centers.¹ His conclusions are based upon a broad and comprehensive survey of geological and biological data and the similarities between certain southern faunas are regarded as exceptions to the general rule to be explained either by the imperfection of the paleontological record, by natural raft transportation, or by convergent evolution. The case of *Cænolestes* is mentioned by Matthew and treated as one of convergence, the views of Broom and Dederer as to its polyprotodont affinities being accepted. A similar view of *Cænolestes* and its allies has been taken by Gregory. The present study does not favor a great degree of convergence, but it tends to substantiate the opinion that an Antarctic land connection is not required to explain the presence of *Cænolestes* in South America.

The many resemblances which *Cænolestes* shows to the Australian

¹The north polar theory had been presented also by Haacke in 1887 and by Haseman in 1912.

peramelids would be difficult to interpret from the distributional standpoint were it not for the occurrence in the Eocene of North America of *Myrmecoboides*, the imperfect remains of which are so suggestive of peramelid relationship. This form seems to show that differentiation along diprotodont lines was well advanced in North America during the basal Eocene at an earlier period than the appearance of any of the South American cœolestids, all of which are of Miocene age. Since all modern peramelids are Australian while *Cœolestes* and *Myrmecoboides* are American, the simplest explanation of their distribution is found in a common northern origin subsequent to the generalized polyprotodont stage; in other words, it may be assumed that the main lines of divergence between polyprotodonts and diprotodonts were laid down in the northern hemisphere. In this case the supposed convergence or parallelism exhibited by *Cœolestes* may be merely of a superficial nature, all its fundamental characters having been derived directly from its northern ancestor.

The absence of fossil marsupials from Asia leaves us without direct evidence of the history of the group in that region, but a wide distribution of the early Tertiary fauna throughout the Holarctic region is generally admitted. Hence it may be assumed that if a division of the marsupial stem had taken place in the early Tertiary of North America the same division or one arising in a similar manner may have extended to Eurasia and thence to Australia. The lack of ancient fossil types from Australia is almost as complete as that from Asia. Still it is important to note that *Wynyardia*, from the Upper Eocene or at latest Oligocene, is the oldest known Australian form and has many diprotodont characters. Moreover, the diprotodonts exhibit various primitive features, in some respects (e. g. auditory ossicles) being more primitive than the didelphids. They reached a very high degree of specialization in the Pleistocene of Australia when such forms as *Diprotodon* and *Thylacoleo* were abundant. At the present time the group is decidedly the dominant one in Australia.

The time of the introduction of marsupials into Australia has been variously estimated from the Jurassic to the Eocene, recent authors being inclined to the later period. Bensley concludes that it is "unlikely that the marsupial radiation could have begun until well on into the middle of the Tertiary period." Elsewhere (1903, p. 86) he speaks of the possibility "that the ancestors of the Australian fauna passed the incipient phases of their evolution in another country, such as Asia or even South America." The same idea is expressed by Matthew (1915, p. 267) as follows:

"There is nothing unlikely in the view that they (Diprotodonta) originated primarily in the North like their polyprotodont and allotherian relatives and were driven southward with the former group and somewhat more thoroughly extinguished in the north, while in Australia they blossomed out into a great adaptive expansion paralleling the absent ungulate mammals."

Therefore, the conclusion that the South American and Australian marsupials had a common Holarctic origin is in accord with the contentions of Matthew as to the origin of mammalian life in general. In fact the case of *Cænolestes* might be regarded as exemplifying the manner in which he has assumed that forms with irregular distribution are to be explained. Thus it is not only a case in which some convergent development may have occurred, but also one in which a slight addition to the fossil record (i. e. *Myrmecoboides*) is of great importance.

The evidence in this case may be divided as follows:

1. The many primitive features of *Cænolestes* indicating the possible development of diprotodont characters very early in the history of the order Marsupialia.
2. The existence of very primitive characters among Australian diprotodonts (see Gregory, 1910, p. 227).
3. The large number of minor resemblances between *Cænolestes* and the peramelids.
4. The probable affinity of *Myrmecoboides* to the modern peramelids.
5. The correspondence of *Myrmecoboides* with the hypothetical ancestor of diprotodonts, that is, the "properamelid" of Bensley.
6. The diprotodont characters shown by the earliest known Australian fossil, *Wynyardia*.
7. The lack of any special affinity between the cænolestids and the didelphids.
8. The probability of a more ancient origin for various marsupial groups than has been supposed (see Gidley, 1915, p. 400).
9. The fact that the Miocene cænolestids were specialized to a high degree and coeval with polyprotodont types which differed from them in essentially the same characters as those which separate modern diprotodonts and polyprotodonts.
10. The presumption of a northern origin of all mammalian groups as indicated by Matthew.

GENERAL SUMMARY.

The most significant results of the foregoing study may be summarized as follows:

1. *Cænolestes* is a surviving member of an ancient group and retains many primitive characters.

Among these may be mentioned: (a) small size and insectivorous habits; (b) in the brain—the very large olfactory bulbs and tubercles and the small simple cerebellum; (c) in the myology—the agreement in certain respects with edentates and insectivores, as in the divided *trapezius*, the *dorso-cuticularis*, the reduced *sartorius*, the sesamoid origin of the *supinator* and the absence of the *pronator quadratus*; (d) in the reproductive system—the possession of a median vagina of similar primitive type to that of *Perameles*; (e) in the alimentary canal—the short colon; (f) in the skeleton—the movable cervical rib and the short pubic symphysis; (g) in the skull—the smooth braincase, large olfactory fossa, proximally expanded nasals, broad frontals, small paroccipital processes, annular tympanic, large persistent mastoid foramen, and columelliform stapes; (h) in the dentition—the large upper incisor formula only exceeded by certain peramelids and didelphids, the large number of lower antemolar teeth, the premolariform canines.

2. It has numerous resemblances to modern peramelids.

These are not of equal importance, but collectively seem significant of affinity. Among them are the following: (a) in external characters—first and fifth toes of forefeet reduced and having nails instead of claws, ear with double antihelical folds, limbs cursorial, tail not prehensile; (b) in the brain—general primitive structure; (c) in the myology¹—general cursorial adaptations; (d) in the skeleton—movable cervical rib, imperforate atlas, short spine of first thoracic vertebra, ossified patella; (e) in the skull—elongate rostrum, short palatine processes of premaxillae, concave basioccipital, nearly perpendicular maxillo-premaxillary suture, long fronto-maxillary union, absence of postorbital processes, lateral exposure of mastoid, small paroccipital and mastoid processes, lacrymal canal with single opening, postzygomatic foramen lacking, stapes columelliform; (f) in the dentition—large incisor formula, quadrate molars with hypocone, proclivous lower incisors, compressed lateral upper incisors,

¹ Adequate comparison of the myology of *Cænolestes* and *Perameles* has not been made.

premolariform canines, specialized posterior premolar; (g) in the urinogenital system—presence of round ligament, median vaginae indicating parturition by direct median passage; (h) in the alimentary canal—small caecum.

3. *It has few non-marsupial characters and no great degree of specialization.*

The important characters not shared with at least some other marsupials are: (a) the preorbital vacuity found elsewhere only among ungulates; (b) the large persistent mastoid foramen; (c) the large carotid canal; (d) the very short pubic symphysis; (e) the extended articular surface of the trochlea of the humerus; (f) the sesamoid in the supinator tendon.

The cardiac gland of the stomach has a counterpart in *Phascolarctos* and *Phascolumys*; otherwise it is unique among marsupials.

4. *It has no especial affinity to the American Didelphiidae.*

Except for a few relatively unimportant myological characters concerning which data are not available for many Australian forms, *Cænolestes* shows practically no exclusive resemblances to the didelphids. On the other hand it differs from them in numerous characters including nearly all the above-mentioned resemblances to the Peramelidae.

5. *The ancestor of the cænolestids was probably a northern form which had already separated from the generalized polyprotodont stock.*

This conclusion is based chiefly upon the probable affinity of *Cænolestes* to modern peramelids and of both to the Eocene *Myrmecoboides* which partially fulfils the hypothetical requirements of the ancestral diprotodont. Many considerations favor this assumption.

6. *The North American ancestor of the cænolestids possibly extended throughout Holarctica and therefore may also have given rise to the Australian diprotodonts.*

This assumes that the main lines of divergence between diprotodonts and polyprotodonts were established in the north prior to the Australian and South American radiations. Hence an Antarctic land connection is not necessary to explain the resemblances between American and Australian marsupials.

7. *The phylogenetic and morphological relations of Cænolestes are best expressed by classifying it in the suborder Diprotodontia, family Palæothentidae, subfamily Cænolestinae.*

This does not emphasize its supposed phylogenetic relationship to the peramelids, but recognizes its advance beyond them to greater morphological similarity to the specialized diprotodonts. On the basis of present findings, the only logical alternative would be to remove peramelids from the Polyprotodontia and unite them with the cænolestids in a group co-ordinate with (1) the Australian Diprotodontia and (2) the remaining Polyprotodontia. This, however, could not be done consistently without further division, especially of polyprotodonts, which does not seem advisable in the present state of knowledge.

LITERATURE CITED.

ALLEN, J. A.

1901. A Preliminary Study of the North American Opossums of the Genus *Didelphis*. Bull. Am. Mus. Nat. Hist., XIV, pp. 149-188, June 15, 1901.

ALSTON, E. R.

1880. Biologia Centrali-Americana, Mammalia. 1879-1882.
1880. On *Antechinomys* and its Allies. Proc. Zool. Soc. Lond., pp. 454-461, 1880.

AMEGHINO, FLORENTINO.

1897. Mammifères crétacés de l'Argentine. Deuxième contribution à la connaissance de la faune mammalogique des couches à *Pyrotherium*. Bol. Inst. Geog. Argent., XVIII, pp. 1-117, 1897.
1899. On the Primitive Type of Plexodont Molars of Mammals. Proc. Zool. Soc. Lond., pp. 555-571, 1899.
1903. Los Diprotodontes del Orden de los Plagiolacoideos. Anales del Museo Nac. Buenos Aires, ser. 3, vol. II, pp. 81-192, July, 1903.

BARDELEBEN, KARL VON.

1894. On the Bones and Muscles of the Mammalian Hand and Foot. Proc. Zool. Soc. Lond., pp. 354-376, 1894.

BATESON, WILLIAM.

1894. Materials for the Study of Variation. Pp. 1-598.

BEDDARD, F. E.

1902. Mammalia. Camb. Nat. Hist. Pp. 1-590. London, 1902.
1895. The Visceral Anatomy and Brain of *Dendrolagus bennetti*. Proc. Zool. Soc. Lond., pp. 131-137, figs. 1-6, 1895.
1908. On the Anatomy of *Antechinomys* and Some Other Marsupials with Special Reference to the Intestinal Tract, etc. Proc. Zool. Soc. Lond., p. 561, 1908.

BENSLEY, B. ARTHUR.

1903. On the Evolution of the Australian Marsupialia; with Remarks on the Relationships of the Marsupials in General. Trans. Linn. Soc. London (2), IX, pp. 83-217, 3 pls., 6 figs., 1903.
1906. The Homologies of the Styler Cusps of the Upper Molars of the Didelphyidae. University of Toronto Biological Series, No. 5, pp. 149-159, 1906.

BROOM, R.

1911. On the Affinities of *Cænolestes*. Proc. Linn. Soc. N. S. W., XXXVI, p. 315, 1911.

COUES, ELLIOTT.

1872. The Osteology and Myology of *Didelphys virginiana*. Mem. Bost. Soc. Nat. Hist., II, pp. 41-154, figs. 1-35, 1872.

CUNNINGHAM, D. J.

1882. Report on Some Points in the Anatomy of the Thylacine (*Thylacinus cynocephalus*), Cuscus (*Phalangista maculata*), and Phascogale (*Phascogale calura*), collected during the Voyage of H. M. S. Challenger in the years 1873-1876; with an account of the Comparative Anatomy of the Intrinsic Muscles and Nerves of the Mammalian Pes. Voy. H. M. S. Challenger, Zool., V, pt. XVI, pp. 1-192, pls. I-XIII, 1882.

DEDERER, PAULINE H.

1909. Comparison of *Cænolestes* with Polyprotodonta and Diprotodonta. Amer. Nat., XLIII, pp. 614-618, figs. 1-2, Oct., 1909.

DORAN, A. H. G.

1879. Morphology of the Mammalian *Ossicula auditus*. Trans. Linn. Soc. Lond., (2), I, Zool., pp. 371-497, pls. 58-64, 1879.

GIDLEY, JAMES WILLIAMS.

1906. Evidence Bearing on Tooth-Cusp Development. Proc. Wash. Acad. Sci., VIII, pp. 91-110, July 10, 1906.
1909. Notes on the Fossil Mammalian Genus *Ptilodus* with Descriptions of New Species. Proc. U. S. Nat. Mus., XXXVI, pp. 611-626, 1 pl., 9 figs., 1909.
1915. An Extinct Marsupial from the Fort Union with Notes on the Myrmecobiidae and other Families of this Group. Proc. U. S. Nat. Mus., XLVIII, pp. 395-402, pl. 23, Jan. 28, 1915.

GREGORY, WILLIAM K.

1910. The Orders of Mammals. Bull. Amer. Mus. Nat. Hist., XXVII, pp. 1-524, Feb., 1910.
1916. Studies on the Evolution of Primates. Bull. Am. Mus. Nat. Hist., XXXV, pp. 239-355, June 16, 1916.
1920. Studies in Comparative Myology and Osteology; No. IV. A Review of the Evolution of the Lacrymal Bone of Vertebrates with Special Reference to that of Mammals. Bull. Am. Mus. Nat. Hist., XLII, pp. 95-263, Dec. 4, 1920.

HAACKE, W.

1887. Der Nordpol als Schöpfungscentrum der Landfauna. Biol. Centralblatt, VI, pp. 363-370, 1887.

HASEMAN, J. D.

1912. Some Factors of Geographical Distribution in South America. Ann. N. Y. Acad. Sci., XXII, pp. 9-112, May 31, 1912.

HEDLEY, C.

1895. Considerations on the Surviving Refugees in Austral Lands of Ancient Antarctic Life. Jour. & Proc. Roy. Soc. N. S. W., XXIX, pp. 278-286, 1895.

HILL, JAMES P.

1899. Contributions to the Morphology and Development of the Female Urogenital Organs in the Marsupialia. I. On the Female Urogenital Organs of *Perameles*, with an Account of the Phenomena of Parturition. Proc. Linn. Soc. N. S. W., XXIV, pp. 42-82, 12 pl., 3 figs., 1899.
1900. Contributions to the Morphology and Development of the Female Urogenital Organs in the Marsupialia. II. On the Female Urogenital Organs of *Myrmecobius fasciatus*. Proc. Linn. Soc. N. S. W., XXV, pp. 519-532, pls. XVII-XIX, 1900.

HUXLEY, T. H.

1880. On the Application of the Laws of Evolution to the Arrangement of the Vertebrata, and More Particularly of the Mammalia. *Proc. Zool. Soc. Lond.*, pp. 649-661, 1880.

JENTINK, F. A.

1885. A Monograph of the Genus *Cuscus*. *Notes Leyden Mus.*, VII, pp. 87-119.

JOHNSTONE, JAMES.

1898. The Thymus in the Marsupials. *Jour. Linn. Soc. Lond.*, XXVI, pp. 537-557, 3 pl., 2 figs., 1898.
 1899. On the Gastric Glands of the Marsupialia. *Jour. Linn. Soc. Lond.*, Zool., XXVII, pp. 1-14, 1 pl., 1 fig., 1899.
 1901. The Neck Glands of the Marsupialia. *Trans. Liverpool Biol. Soc.*, XV, pp. 354-362, 3 figs., 1901.

LECHE, W.

- 1874-1900. *Bronn's Klassen u. Ordn. Thierreichs*, Bd. VI, Abth. V, Mammalia, Bd. I., Leipzig, 1874-1900.

LYDEKKER, RICHARD.

1894. A Hand-Book to the Marsupialia and Monotremata. *Allen's Nat. Lib.*, London, 1894.
 1899. The Dental Formula of the Marsupial and Placental Carnivora. *Proc. Zool. Soc. Lond.*, pp. 922-928, pl. LXII, 1899.

MACALISTER, ALEXANDER.

1870. On the Myology of the Wombat (*Phascolomys wombata*) and the Tasmanian Devil (*Sarcophilus ursinus*). *Ann. & Mag. Nat. Hist.*, (4), V, pp. 153-173, March, 1870.
 1872. Further Observations on the Myology of *Sarcophilus ursinus*. *Ann. & Mag. Nat. Hist.*, (4), X, pp. 17-20, July, 1872.
 1872. The Muscular Anatomy of the Koala (*Phascolarctos cinereus*). *Ann. & Mag. Nat. Hist.*, (4), X, pp. 127-134, Aug., 1872.

MATTHEW, W. D.

1915. Climate and Evolution. *Ann. N. Y. Acad. Sci.*, XXIV, pp. 171-318, Feb. 18, 1915.

METCALF, M. M.

1920. Upon an Important Method of Studying Problems of Relationship and of Geographical Distribution. *Proc. Nat. Acad. Sci.*, VI, pp. 432-433, 1920.

MITCHELL, P. CHALMERS.

1905. On the Intestinal Tract of Mammals. *Trans. Zool. Soc. Lond.*, XVII, Pt. V, pp. 437-535, 1905.

ORTMANN, A. E.

1902. Tertiary Invertebrates. *Repts. Princeton Univ. Exped. to Patagonia*, IV, Paleon., pp. 45-332, pls. XI-XXXIV, April 19, 1902.

OSBORN, HENRY FAIRFIELD.

1907. Evolution of Mammalian Molar Teeth. Pp. 1-250. New York, 1907.
 1910. The Age of Mammals in Europe, Asia and North America. Pp. XVIII-635. New York, October, 1910.

OSGOOD, WILFRED H.

1912. Mammals from Western Venezuela and Eastern Colombia. *Field Mus. Nat. Hist.*, Zool., X, pp. 33-66, Jan. 10, 1912.

PARSONS, F. G.

1896. On the Anatomy of *Petrogale xanthopus*, Compared with that of Other Kangaroos. Proc. Zool. Soc. Lond., 1896, pt. 3, pp. 683-714, 10 figs., 1896.
 1898. The Muscles of Mammals with Special Relation to Human Myology. Journ. Anat. & Phys., XXXII, pp. 428-450, 721-752, 1898.
 1898. The Limb Myology of *Gymnura rafflesii*. Journ. Anat. & Phys., XXXII, pp. 312-324, 1898.
 1903. On the Anatomy of the Pig-footed Bandicoot (*Charopus castanotis*). Journ. Linn. Soc. Lond., Zool., XXIX, pp. 64-80, 10 figs., 1903.

POULTON, E. B.

1883. On the Tongues of the Marsupialia. Proc. Zool. Soc. Lond., pp. 599-628, pls. LIV-LV, 1883.

RUGE, GEORG.

1878. Untersuchung über die Extensorengruppe am Unterschenkel und Füsse der Säugethiere. Morph. Jahrb., IV, pp. 592-643, pls. XXXII-XXXV, 1878.
 1878. Zur vergleichenden Anatomie der tiefen Muskeln in der Fusssohle. Morph. Jahrb., IV, pp. 644-659, pls. XXXIV-XXXV, 1878.

SCOTT, W. B.

1913. A History of Land Mammals in the Western Hemisphere. Pp. 1-693.

SIDEBOTHAM, E. J.

1885. On the Myology of the Water-Opossum. Proc. Zool. Soc. Lond., pp. 6-22, 1885.

SINCLAIR, W. J.

1905. The Marsupial Fauna of the Santa Cruz Beds. Proc. Amer. Philos. Soc., XLIV, pp. 73-81, 2 pls., 1905.
 1906. Mammalia of the Santa Cruz Beds. Marsupialia. Repts. Princeton Univ. Expeds. to Patagonia, IV, Paleon., I, pp. 333-460, Sept. 14, 1906.

SPENCER, BALDWIN.

1900. A Description of *Wynyardia bassiana*, a Fossil Marsupial from the Tertiary Beds of Table Cape, Tasmania. Proc. Zool. Soc. Lond., pp. 776-794, pls. XLIX-L, 1900.

STIRLING, E. C.

1891. Description of a New Genus and Species of Marsupialia, *Notoryctes typhlops*. Trans. Roy. Soc. South Aust., XIV, pp. 154-187, pls. II-IX, 1891.

STONE, WITMER.

1914. On a Collection of Mammals from Ecuador. Proc. Acad. Nat. Sci. Phila., pp. 17-19, Mch. 31, 1914.

SWEET, GEORGIANA.

1904. Contributions to our Knowledge of the Anatomy of *Notoryctes typhlops* Stirling. Parts I and II. Proc. Roy. Soc. Victoria, N. S., XVII, pp. 76-111, 4 pls., 1904.
 1907. The Skin, Hair, and Reproductive Organs of *Notoryctes*. Contributions to our Knowledge of the Anatomy of *Notoryctes typhlops* Stirling. Parts IV-V. Quart. Jour. Micr. Sc., LI, pp. 325-344, 2 pls. 1 fig., 1907.

SYMINGTON, JOHNSON.

1898. The Thymus Gland in the Marsupialia. Jour. Anat. & Phys., XXXII, N. S. XII, pp. 278-291, figs. 1-4, 1898.

THOMAS, OLDFIELD.

1887. On the Homologies and Succession of the Teeth in the Dasyuridae. Philos. Trans. Roy. Soc. Lond., CLXXVIII, B, pp. 443-462, pls. 27-28, 1887.

1888. Catalogue of Marsupialia and Monotremata in the British Museum. Pp. 1-401, pls. I-XXVIII, 1888.
- 1895a. Descriptions of Four Small Mammals from South America, Including One Belonging to the Peculiar Marsupial Genus "*Hyracodon*" Toms. Ann. & Mag. Nat. Hist., (6), XVI, pp. 367-370, Nov., 1895.
- 1895b. On *Cenolestes*, a Still Existing Survivor of the Epanorthidae of Ameghino. Proc. Zool. Soc. Lond., p. 870, 1895.
1896. On *Cenolestes* n. g. of Marsupials. (Abstr.) Zool. Anz., XIX, no. 493, p. 31, 1896.
1917. Preliminary Diagnoses of New Mammals Obtained by the Yale-National Geographic Society Peruvian Expedition. Smiths. Misc. Coll., LXVIII, p. 3, Apr. 10, 1917.
1920. Report on the Mammalia Collected by Mr. Edmund Heller during the Peruvian Expedition of 1915 under the Auspices of Yale University and the National Geographic Society. Proc. U. S. Nat. Mus., LVIII, pp. 217-249, pls. 14-15, 1920.
- THOMPSON, PETER, and HILLIER, W. T.
1905. The Myology of the Hind Limb of the Marsupial Mole (*Notoryctes typhlops*). Jour. Anat. & Phys., XXXIX, pp. 308-331, pls. XXXVIII-XXXIX.
- TIMS, H. W. M.
1903. Evolution of the Teeth in the Mammalia. Jour. Anat. & Phys., XXXVII, pp. 132-140, 1903.
- TOMES, R. F.
1860. Notes on a Second Collection of Mammalia Made by Mr. Fraser in the Republic of Ecuador. Proc. Zool. Soc. Lond., p. 213, 1860.
1863. Notice of a New American Form of Marsupial. Proc. Zool. Soc. Lond., pp. 50-51, pl. VIII (col.), 1863.
- TROUESSART, E. L.
1898. Catalogus Mammalium, V, p. 1205, 1898.
- WATERHOUSE, G. R.
1846. Natural History of the Mammalia, Vol. I., Marsupialia, 1846.
- WEBER, MAX.
1904. Die Säugethiere. Pp. 1-866. Jena, 1904.
- WILSON, J. T.
1894. On the Myology of *Notoryctes typhlops*, with Comparative Notes. Trans. Roy. Soc. South Aust., XVIII, pp. 3-74, pls. II-XV, 1894.
- WILSON, J. T., and HILL, J. P.
1897. Observations upon the Development and Succession of the Teeth in *Perameles*; Together with a Contribution to the Discussion of the Homologies of the Teeth in Marsupial Animals. Quart. Jour. Micr. Sci., XXXIX, N. S., pp. 427-588, pls. 25-32, 1897.
- WINDLE, BERTRAM C. A., and PARSONS, F. G.
1898. On the Anatomy of *Macropus rufus*. Jour. Anat. & Phys., XXXII, N. S., XII, pp. 119-134, 1898.
- WINGE, HERLUF.
1882. Om Pattedyrenes Tandskifte, især med Hensyn til Tændernes Former. Vid. Medd. f. d. Naturh. Foren., Kjöbenhavn, pp. 15-67, 1882.
- WOODWARD, M. F.
1893. On the Development of the Teeth in the Macropodidae. Proc. Zool. Soc. Lond., pp. 459-473, 1893.
1896. On the Teeth of Certain Insectivora. Proc. Zool. Soc. Lond., pp. 557-594.

THE BRAIN OF *CÆNOLESTES OBSCURUS*.

BY C. JUDSON HERRICK.

Plates XXI-XXII.

These notes are based on a single female specimen which had been preserved in formalin and later transferred to alcohol. Since the cranial cavity had not been opened before the specimen was put into the hardening fluid, the preservation of the brain is not as perfect as might be desired. In particular, the swelling of the brain tissue produced by the formalin has caused considerable compression of the brain within the endocranial cavity, thus possibly exaggerating somewhat the superficial relief, so far as this conforms to the sculpturing of the endocranial cavity, and obscuring some other features.

The brain was very skillfully removed from the skull of the alcoholic specimen in the Anatomical Laboratory of the University of Chicago by Dr. G. W. Bartlemez and drawn under the Zeiss stereo-binocular microscope by Mr. A. B. Streedain. Since it seemed desirable to preserve the brain intact in the hope of a later opportunity to prepare it for microscopic examination, this report must of necessity be limited to the superficial anatomy.

Measurements.—The dimensions as measured on the alcoholic specimen are as follows:—

Total length, tip of olfactory bulb to first spinal nerve	14.1 mm.
Length, tip of olfactory bulb to rostral end of cerebral hemisphere	2.6 mm.
Length of cerebral hemisphere	10.0 mm.
Length of cerebellum on longitudinal axis of brain in median plane	3.0 mm.
Greatest width of both olfactory bulbs	7.6 mm.
Greatest width of both cerebral hemispheres	11.8 mm.
Total width of cerebellum and flocculi	11.0 mm.
Width of cerebellum exclusive of flocculi	8.8 mm.

For our most precise knowledge of the brains of lower mammals we are indebted to the researches of Elliot Smith. Throughout this account we shall make frequent references to his papers and base our interpretations to a large extent upon them. His papers cited in the appended bibliography give references to the other relevant literature.

The general form of the brain of this little marsupial is evident from the figures. It is apparent that we have here before us one of the simplest types of mammalian brain hitherto described. The brain is strongly macrosomatic, as shown by the enormous size of the olfactory bulbs and secondary olfactory area.

The *olfactory bulbs* are closely appressed to each other in the median plane, convex above and concave below. They are very slightly overlapped dorsally by the cerebral hemispheres. From the widely flaring lateral borders of the bulb the lateral olfactory tract passes spinalward and ventralward along the lateral border of the very large tuberculum olfactorium.

The *cerebral hemispheres* extend backward quite to the cerebellum, these two structures being in intimate contact for the entire width of the body of the cerebellum. There is a very slight divergence of the posterior borders of the hemispheres from the median plane, which is somewhat exaggerated in Figure 1 (Pl. XXI), so that even if the meninges were entirely removed from the median longitudinal fissure (as has not been done), but little if any of the mesencephalon would be visible from the dorsal surface. This is in contrast to the usual marsupial arrangement, for the corpora quadrigemina are in most cases well exposed dorsally. (*Petaurus* is another exception; see Elliot Smith, '95, p. 168.)

Superficially the cerebral hemisphere, exclusive of the olfactory bulb, exhibits three chief regions:—(1) the dorsal convexity; (2) the lateral convexity, or pyriform lobe; (3) the ventral convexity, or tuberculum olfactorium.

The dorsal convexity of the hemisphere is purely neopallial; that is, it is non-olfactory cortex. About one-fifth of the distance backward from the frontal to the posterior pole of the hemisphere there is a distinct, though shallow, transverse sulcus which probably represents the sulcus orbitalis of Elliot Smith's descriptions. Otherwise the dorsal convexity is smooth.

The dorsal convexity is bounded laterally by an imperfectly developed fissura rhinalis. This begins anteriorly as a sharp sulcus in the transverse fissure between the olfactory bulb and the cerebral hemisphere at the dorso-lateral angle of the tuberculum olfactorium (Pl. XXI, fig. 2) and extends backward. On the lateral aspect of the hemisphere it is obscurely confluent with the orbital sulcus, and behind this level it disappears in a depressed area on the lateral wall of the hemisphere. Two-thirds of the distance back from the anterior to the posterior pole of the hemisphere this depressed area is slightly deepened, thus marking more precisely the location of the fissura rhinalis in this region; and at the posterior end of the hemisphere there is a wide, shallow notch which marks the posterior end of this fissure (Pl. XXI, figs. 1 and 2).

The pyriform lobe (*lobus piriformis*) comprises the larger part of the lateral and ventral aspects of the hemisphere. As we have seen above, it is very imperfectly separated from the dorsal neopallial cortex, though the location of this boundary, the fissura rhinalis, is evident.

Antero-ventrally it is separated from the tuberculum olfactorium by a very sharp fissura endorhinalis. The ventral surface of the pyriform lobe shows two shallow depressions separated by an elevated ridge running obliquely backward and lateralward from the posterior margin of the tuberculum olfactorium. Further information regarding the internal structure is desirable before the signification of this sculpturing is explained. It may be due merely to the conformation of the brain to the wall of the cranium.

The tuberculum olfactorium is very large and strongly convex ventrally. The medial borders of the two tubercula are divaricated posteriorly, exposing a portion of the anterior perforated space in front of the optic chiasma which probably includes the diagonal band of Broca (Pl. XXII).

The lateral olfactory tract arises from the ventrolateral border of the olfactory bulb and can readily be seen as a clear white stripe accompanying the fissura endorhinalis. This band of fibers lies distinctly on the ventromedial side of the fissura, that is, within the tuberculum olfactorium, though we may infer by analogy with other mammals that many of the fibers are distributed within the pyriform lobe on the other side of the fissure. Microscopic examination will probably show that a portion of the tract lies in the floor and walls of the fissure.

The tuberculum olfactorium in mammals generally is a basal, that is, subcortical, reflex center, receiving olfactory fibers of the second order from the olfactory bulb. The pyriform lobe, on the other hand, is a structure of transitional type. So far as it and the underlying amygdala receive secondary olfactory fibers from the lateral olfactory tract it should be considered as a part of the basal secondary olfactory area (nucleus olfactorius lateralis). So far as its differentiated cortex is in physiological connection with this secondary olfactory area it should be regarded as archipallial in type, i. e., olfactory cortex of the same type as the hippocampal cortex. So far as its cortex is in physiological connection with the non-olfactory thalamic projection fibers it is neopallial, i. e., of the same type as the dorsal cortex. The surface characteristics suggest that in *Cænolestes* these three components of the pyriform lobe are very incompletely differentiated, thus resembling the still more generalized reptilian condition (cf. Elliot Smith, '10, and Crosby, '17). In higher mammals, on the other hand, these components of the pyriform lobe attain much more distinct spatial localization in the gyrus hippocampi formation, though even in the human brain we find the same transitional type of structure in the uncus region.

The cerebellum.—The cerebellum is smaller and simpler than in any hitherto described mammals with the exception of *Notoryctes* and

Perameles, though the elephant-shrew, *Macroscelides*, has a cerebellum but little more complex than that of *Perameles* (Elliot Smith, '02b).

The cerebellum is composed of a simply organized transverse body and the two lateral floccular lobes. (In the preparation of the brain the left flocculus was destroyed. In the figures it has been restored after the one on the opposite side.) Each flocculus is mushroom-shaped with a slender pedicle and a widely expanded cap or pileus, whose surface is somewhat lacerated but apparently was smooth or nearly so.

The body of the cerebellum, so far as it is visible from the surface, is a transverse bar with two shorter convolutions under its posterior border. The main bar consists of a plump median lobe and two lateral lobes, each of which is somewhat less than half the length in the transverse plane of the medial lobe. The median lobe and the two smaller posterior convolutions apparently correspond with the vermicular portion of higher cerebella and the lateral lobes with the hemispheres.

Since only a part of the cerebellum is visible in the undissected specimen, it is impossible to determine with certainty the homologies of the exposed structures. But comparison with the very similar cerebella of *Notoryctes* and *Perameles* as described and figured by Elliot Smith ('03a and '03b) suggests that in *Cænolestes* the lobus anterior and fissura prima are entirely concealed in the transverse fissure between the cerebellum and the cerebral hemispheres, the transverse bar is the lobus medius, the fissure limiting it posteriorly is the fissura secunda, the first of the two shorter convolutions is the uvula, the fissure behind the latter is the fissure postnodularis, and the posterior one of the shorter convolutions is the nodulus, these names being used as defined by Elliot Smith ('03a).

The brain stem.—On the ventral surface (Pl. XXII) the optic nerves are seen to be very slender. Behind the optic chiasma is a rather wide tuber cinereum whose surface is somewhat obscured by meninges which in view of the poor state of preservation of the specimen it seemed inexpedient to attempt to remove. The pituitary body is mushroom-shaped and elevated on a very short infundibular stalk. On account of the flexure of the brain in the isthmus region the cerebral peduncles are entirely concealed. Stumps of the third and fourth cranial nerves are seen in the usual relations.

The medulla oblongata under the cerebellum is very wide. The sculpturing of the ventral surface is obscured by pressure against the floor of the cranial cavity and posteriorly by a blood clot in the meninges. Nevertheless the roots of all of the cranial nerves were identified. The trigeminus is large, arising from the posterior border of the pons, which is very slender. The abducens is very minute and is recognized (with

some uncertainty) at the posterior border of the pons. Immediately below the pons is a broad, flat, transverse band which is probably the trapezoid body and at whose lateral ends are the stumps of the VIII nerves. The facialis root arises from the infero-lateral surface of the trapezoid body; the IX, X and XI nerves arise from the lateral surface of the medulla oblongata; and medially of these are the XII roots. The pyramidal tracts are visible as light colored bands near the mid-ventral line extending downward from the posterior surface of the pons. At the level of the first spinal roots there is a strong cervical flexure.

General considerations.—Comparing the brain as a whole with those of other mammals, it is seen to resemble most closely those of *Notoryctes* and *Perameles*. In all three cases the rhinencephalon is enormously developed, the olfactory bulbs and tubercles being very large. The cerebral cortex is nearly smooth and apparently relative to the total size of the brain less extensive than in any other mammals hitherto described. Elliot Smith's figures of *Notoryctes* ('95) indicate that in this genus the cerebral hemispheres are relatively smaller than in *Cænolestes*; his figure of the ventral surface of *Perameles* ('02, p. 171, fig. 52) suggests that here the hemisphere is relatively as large as in *Cænolestes* or larger. In absolute dimensions the brain of *Notoryctes* is about the same size as that of *Cænolestes*, while that of *Perameles* is more than three times as long. The simplicity of these brains cannot, therefore, be correlated directly with the size of the animals. In fact, our figure of the ventral view of the brain of *Cænolestes* resembles more closely Elliot Smith's figure of the ventral surface of the larger *Perameles* than of the *Notoryctes* of equal size. Both *Notoryctes* and *Perameles* belong to the Polyprotodontia. The larger members of the Diprotodontia, such as the kangaroos, have larger and more highly convoluted brains than any of the Polyprotodontia. The brain of *Cænolestes* is more simply organized than that of any Australian diprotodont.

LITERATURE.

- CROSBY, ELIZABETH CAROLINE. 1917. The Forebrain of *Alligator mississippiensis*. Jour. Comp. Neur., Vol. 27, pp. 325-402.
- SMITH, G. ELLIOT. 1894. A Preliminary Communication upon the Cerebral Commissures of the Mammalia, with special Reference to the Monotremata and Marsupialia. Proc. Linnean Soc. New South Wales, Series 2, Vol. 9, pp. 635-657.
1895. The Comparative Anatomy of the Cerebrum of *Notoryctes typhlops*. Trans. Roy. Soc. South Australia, 1895, pp. 167-193.
1899. Further Observations on the Anatomy of the Brain in the Monotremata. Jour. Anat. and Physiol., Vol. 33, pp. 309-342.

1899a. The Brain in the Edentata. Trans. Linnean Soc. London, Series 2, Zoology, Vol. 7, Part 7, pp. 377-394.

1902. Descriptive and Illustrated Catalogue of the Physiological Series of Comparative Anatomy contained in the Museum of the Royal College of Surgeons of England, 2 Ed., Vol. 2.

1902a. On a Peculiarity of the Cerebral Commissures in Certain Marsupialia, not hitherto Recognized as a Distinctive Feature of the Diprotodontia. Proc. Roy. Soc. London, Vol. 70, pp. 226-231.

1902b. Notes on the Brain of Macroscelides and other Insectivora. Jour. Linnean Soc. London, Zoology, Vol. 28, pp. 443-448.

1902c. The Primary Subdivision of the Mammalian Cerebellum. Jour. Anat. and Physiol., Vol. 36, pp. 381-385.

1903. Notes on the Morphology of the Cerebellum. Jour. Anat. and Physiol., Vol. 37, pp. 329-332.

1903a. Further Observations on the Natural Mode of Subdivision of the Mammalian Cerebellum. Anat. Anzeiger, Bd. 23, pp. 368-384.

1903b. On the Morphology of the Brain in the Mammalia, with special Reference to that of the Lemurs, Recent and Extinct. Trans. Linnean Soc. London, Series 2, Zoology, Vol. 8, Part 10, pp. 319-432.

1910. The Arris and Gale Lectures on Some Problems Relating to the Evolution of the Brain. The Lancet, Jan. 1, 15, and 22, 1910.



Fig. 1. *Amphioxus* (L.)
Amphioxus (L.)
Amphioxus (L.)

Length 1.5
 Breadth 1.5
 Height 1.5



Fig. 2. *Amphioxus* (L.)
Amphioxus (L.)

111. *Thysanotus* (Thysanotus) *Thysanotus* (Thysanotus) 215

112. *Thysanotus* (Thysanotus) *Thysanotus* (Thysanotus) 216

113. *Thysanotus* (Thysanotus) *Thysanotus* (Thysanotus) 217

114. *Thysanotus* (Thysanotus) *Thysanotus* (Thysanotus) 218

115. *Thysanotus* (Thysanotus) *Thysanotus* (Thysanotus) 219

116. *Thysanotus* (Thysanotus) *Thysanotus* (Thysanotus) 220

117. *Thysanotus* (Thysanotus) *Thysanotus* (Thysanotus) 221

118. *Thysanotus* (Thysanotus) *Thysanotus* (Thysanotus) 222

119. *Thysanotus* (Thysanotus) *Thysanotus* (Thysanotus) 223

120. *Thysanotus* (Thysanotus) *Thysanotus* (Thysanotus) 224

PLATE II.

EXTERNAL CHARACTERS.


Three times natural size.

Fig. 1. Head.

Fig. 2. Dorsal aspect of rhinarium.

Fig. 3. Sole of hind foot.

Fig. 4. Sole of fore foot.



1



2



3



4

EXTERNAL CHARACTERS.
Three times natural size.



GENERAL ASPECT OF OUTER MUSCLES AFTER REMOVAL OF SUPERFICIAL FASCIA.

a., anconeus;
ac.d., acromio-deltoides;
act., acromio-trapezius;
b.a., brachialis anticus;
b.b., biceps brachii;
b.c., biventer cervicis;
b.f., biceps femoris;
br., brachioradialis;
bu., buccinator;
c.c., cruro-coccygeus;
c.c.m., muscular bulb of corpus cavernosum;
cl.o., cleido-occipitalis;
Cp².Cp³., Cowper's glands;
c.s.m., muscular bulb of corpus spongiosum;
d.c., dorso-cuticularis (cut);
di., digastricus;
e.c.b., extensor carpi radialis brevis;

e.c.l., extensor carpi radialis longus;
e.d., extensor digitorum communis
 and lateralis;
f., fascia over rhomboideus;
f.c., femoro-coccygeus;
f.c.u., flexor carpi ulnaris;
f.d.f., flexor digitorum fibularis;
g., gastrocnemius;
g.m., gluteus maximus;
il., iliacus;
l.d., latissimus dorsi;
m.d.a., mandibulo-auricularis;
m.r., ramus of mandible;
ms.i., inferior masseter;
ms.s., superior masseter;
ob.e., obliquus externus;
ob.i., obliquus internus;
om., omohyoides;
per., peronei;
ps.m., psoas magnus;

pt., pectoralis;
r.a., rectus abdominis;
r.gl., rectal gland;
scd., scalenus;
scr., scrotum;
s.m., serratus magnus;
smt., semitendinosus;
sp., sphincter cloacae;
sp.d., spino-deltoides;
sp.t., spino-trapezius;
st.h., sterno-hyoideus;
st.m., sterno-mastoideus;
t.a., tibialis anticus;
t.c.l., triceps, caput laterale;
t.c.la., triceps, caput longum;
te., teres;
tm., temporalis;
v.e., vastus externus;
x¹, x², x³, zygomatikus.



GENERAL MUSCULATURE AFTER REMOVAL OF SUPERFICIAL FASCIA.

10

1. *What is the main purpose of the study?*

© 2000 Blackwell Science Ltd *Journal of Internal Medicine* 247: 395–401


PLATE IV.

MUSCLES AND GLANDS OF THE HEAD AND NECK.

Fig. 1. Muscles of right side of head and neck after removal of trapezius muscle.
Three and one-half times natural size.

<i>acd.</i> , acromio-deltoides;	<i>o.c.i.v.</i> , omo-cleido-transversarius
<i>bu.</i> , buccinator;	ventralis;
<i>clm.</i> , cleido-mastoideus;	<i>om.</i> , omohyoideus;
<i>clo.</i> , cleido-occipitalis;	<i>rh.c.</i> , rhomboideus capitis;
<i>di.</i> , digastricus;	<i>rh.v.</i> , rhomboideus vertebris;
<i>mda.</i> , mandibulo-aicularis;	<i>r.n.</i> , retractor naris;
<i>ms.i.</i> , inferior masseter;	<i>sp.</i> , splenius;
<i>ms.s.</i> , superior masseter;	<i>sth.</i> , sterno-hyoideus;
<i>ms.t.</i> , tendon of masseter;	<i>stm.</i> , sterno-mastoideus;
<i>o.c.i.d.</i> , omo-cleido-transversarius	<i>t.</i> , temporalis;
dorsalis;	<i>z¹, z², z³</i> , zygomaticus.

Fig. 2. Salivary glands. One and one-fourth times natural size.
pd., parotid duct;
pg., parotid gland;
sm.g., submaxillary gland.



MUSCLES AND GLANDS OF THE HEAD AND NECK.



PLATE V.

MUSCLES OF THE LEGS.



Two and one-half times natural size.

Fig. 1. Left hind leg from outside.

<i>b.f.</i> , biceps femoris;	<i>g.m.</i> , gluteus maximus;
<i>c.c.</i> , cruro-coccygeus;	<i>p.</i> , peronei;
<i>cf.</i> , caudo-femoralis;	<i>r.f.</i> , rectus femoris;
<i>e.c.</i> , extensor-caudae;	<i>sm.</i> , semimembranosus;
<i>f.c.</i> , femoro-coccygeus;	<i>st.</i> , semitendinosus;
<i>f.d.f.</i> , flexor digitorum fibularis;	<i>t.a.</i> , tibialis anticus;
<i>g.</i> , gastrocnemius;	<i>v.e.</i> , vastus externus.

Fig. 2. Right foreleg from inside.

<i>a.</i> , anconeus;	<i>o.c.t.d.</i> , omo-cleido-transversarius
<i>b.a.</i> , brachialis anticus;	dorsalis (atlanto-scapularis);
<i>b.b.</i> , biceps brachii;	<i>o.c.t.v.</i> , omo-cleido-transversarius
<i>br.</i> , brachioradialis;	ventralis (atlanto-acromialis);
<i>cl.</i> , clavicle;	<i>p.l.</i> , palmaris longus;
<i>cl.m.</i> , cleido-mastoideus;	<i>p.mi.</i> , pectoralis minor;
<i>cl.o.</i> , cleido-occipitalis;	<i>p.mj.</i> , pectoralis major;
<i>c.m.</i> , cutaneus maximus;	<i>p.t.</i> , pronator teres;
<i>d.</i> , deltoideus;	<i>sc.</i> , subclavius;
<i>e.c.r.</i> , extensor carpi radialis brevis;	<i>s.m.</i> , serratus magnus;
<i>el.</i> , epitrochlearis;	<i>ss.</i> , subscapularis;
<i>f.c.r.</i> , flexor carpi radialis;	<i>tr.</i> , trapezius;
<i>f.c.u.</i> , flexor carpi ulnaris;	<i>t.c.l.</i> , triceps brachii, caput laterale;
<i>f.d.p.</i> , flexor digitorum profundus;	<i>t.c.lo.</i> , triceps, caput longum;
<i>l.d.</i> , latissimus dorsi;	<i>t.c.m.</i> , triceps, caput mediale;
<i>l.s.</i> , levator scapulae;	<i>t.m.</i> , teres major.
<i>o.</i> , omohyoideus;	



MUSCLES OF THE LEGS.

Two and one-half times natural size.

THE UNIVERSITY OF CHICAGO
 LIBRARY

THE UNIVERSITY OF CHICAGO
 LIBRARY
 1215 EAST 58TH STREET
 CHICAGO, ILL. 60637
 TEL. 773-709-9300
 FAX 773-709-9301
 WWW.CHICAGO.EDU

THE UNIVERSITY OF CHICAGO
 LIBRARY
 1215 EAST 58TH STREET
 CHICAGO, ILL. 60637
 TEL. 773-709-9300
 FAX 773-709-9301
 WWW.CHICAGO.EDU

THE UNIVERSITY OF CHICAGO
 LIBRARY

THE UNIVERSITY OF CHICAGO
 LIBRARY
 1215 EAST 58TH STREET
 CHICAGO, ILL. 60637
 TEL. 773-709-9300
 FAX 773-709-9301
 WWW.CHICAGO.EDU

THE UNIVERSITY OF CHICAGO
 LIBRARY
 1215 EAST 58TH STREET
 CHICAGO, ILL. 60637
 TEL. 773-709-9300
 FAX 773-709-9301
 WWW.CHICAGO.EDU

THE UNIVERSITY OF CHICAGO
 LIBRARY

PLATE VI.

MUSCLES OF THE THROAT AND INNER SIDE OF HIND LEG.

Three times natural size.

Fig. 1. Muscles of the throat.


<i>cl.m.</i> , cleido-mastoideus;	<i>o.c.t.v.</i> , omo-cleido-transversarius
<i>cl.o.</i> , cleido-occipitalis;	ventralis;
<i>d.</i> , digastricus;	<i>sc.</i> , scalenus;
<i>gh.</i> , geniohyoideus;	<i>st.h.</i> , sterno-hyoideus;
<i>hg.</i> , hyoglossus;	<i>st.m.</i> , sterno-mastoideus;
<i>l.gl.</i> , lymphatic gland;	<i>st.t.</i> , sterno-thyroideus;
<i>o.</i> , omohyoideus;	<i>t.gl.</i> , thyroid glands;
<i>o.c.t.d.</i> , omo-cleido-transversarius	<i>t.h.</i> , thyro-hyoideus.
dorsalis;	

Fig. 2. Inner side of hind leg showing innervation of sartorius muscle by saphenus nerve.


<i>a.l.</i> , adductor longus;	<i>p.</i> , pectineus;
<i>a.m.</i> , adductor magnus;	<i>ps.m.</i> , psoas major;
<i>c.n.</i> , crural nerve;	<i>r.f.</i> , rectus femoris;
<i>g.</i> , gastrocnemius;	<i>sa.</i> , sartorius;
<i>g.m.</i> , gluteus maximus;	<i>sm.</i> , semimembranosus;
<i>gr.</i> , gracilis;	<i>s.n.</i> , saphenus nerve;
<i>l.p.</i> , ligamentum poupartii;	<i>st.</i> , semitendinosus;
<i>o.e.</i> , obliquus externus;	<i>t.a.</i> , tibialis anticus;
<i>o.i.</i> , obliquus internus;	<i>v.i.</i> , vastus internus.

Fig. 3. Same as Fig. 2 with sartorius muscle *in situ*.


1



2



3



MUSCLES OF THROAT AND INNER SIDE OF THIGH.
Enlarged.



The human liver is a large, fleshy organ, and is situated in the right hypochondrium, and extends to the umbilicus. It is divided into two lobes, the right and left, by a fissure, which is called the falciform ligament. The right lobe is larger than the left, and is covered by a thin membrane, called the peritoneum. The left lobe is smaller, and is covered by a thin membrane, called the peritoneum. The liver is connected to the gallbladder by a duct, called the common bile duct. The liver is also connected to the stomach by a duct, called the hepatic duct. The liver is supplied with blood by the hepatic artery, and with bile by the common bile duct.

The human liver is a large, fleshy organ, and is situated in the right hypochondrium, and extends to the umbilicus. It is divided into two lobes, the right and left, by a fissure, which is called the falciform ligament. The right lobe is larger than the left, and is covered by a thin membrane, called the peritoneum. The left lobe is smaller, and is covered by a thin membrane, called the peritoneum. The liver is connected to the gallbladder by a duct, called the common bile duct. The liver is also connected to the stomach by a duct, called the hepatic duct. The liver is supplied with blood by the hepatic artery, and with bile by the common bile duct.



PLATE VII.

PARTS OF ALIMENTARY CANAL.

Two to three times natural size.

Fig. 1. Spleen (x2).


Fig. 2. Ventral view of liver and adjoining organs. Anterior part of liver elevated and posterior lobes spread apart. Duodenum reverted and displaced to left (x2 $\frac{1}{4}$).

- | | |
|---|---|
| <i>c.d.</i> , cystic duct; | <i>p.v.</i> , portal vein; |
| <i>c.l.</i> , caudate lobe of liver; | <i>r.c.</i> , right central lobe of liver; |
| <i>d.</i> , duodenum; | <i>r.l.</i> , right lateral; |
| <i>d.ch.</i> , ductus choledochus; | <i>r.v.</i> , renal veins; |
| <i>g.b.</i> , gall bladder; | <i>sp.ca.</i> , caudal division of spigelian lobe; |
| <i>h.d.</i> , hepatic duct; | <i>sp.cr.</i> , cranial division of spigelian lobe; |
| <i>l.l.</i> , left lateral lobe of liver; | <i>st.</i> , stomach; |
| <i>oe.</i> , oesophagus; | <i>v.c.</i> , vena cava. |
| <i>p.d.</i> , pancreatic duct; | |

Fig. 3. Caecum or vermiform appendix (x3).

Figs. 4-5. Stomach (x2).

- | | |
|---|---|
| <i>c.g.</i> , cardiac gland; | <i>gl.</i> , glandular structure of stomach |
| <i>c.s.</i> , cardiac division of stomach, with | after removal of outer coat; |
| specialized glands; | <i>oe.</i> , oesophagus; |
| <i>d.</i> , duodenum; | <i>oe.s.</i> , oesophageal division of stomach; |
| | <i>p.s.</i> , pyloric division of stomach. |



PARTS OF ALIMENTARY SYSTEM.
Enlarged.

PLATE VIII.

REPRODUCTIVE ORGANS.


Fig. 1. Female reproductive organs. Ventral view (x5).

- | | |
|--------------------------------|--------------------------------------|
| <i>bl.</i> , bladder; | <i>r.l.</i> , round ligament; |
| <i>br.l.</i> , broad ligament; | <i>sp.c.</i> , sphincter cloacae; |
| <i>cl.</i> , cloacal opening; | <i>u.</i> , ureter; |
| <i>f.l.</i> , fallopian tube; | <i>u.b.</i> , uterine body; |
| <i>gl.</i> , lymphatic gland; | <i>u.n.</i> , uterine neck; |
| <i>m.v.</i> , median vagina; | <i>u.s.</i> , urogenital sinus; |
| <i>o.</i> , ovary; | <i>v.c.</i> , lateral vaginal canal. |

Fig. 2. Male reproductive organs. Ventral view (x2½).

- | | |
|--|--|
| <i>a.</i> , adrenal body; | <i>pr.</i> , prostate; |
| <i>ao.</i> , aorta; | <i>r.</i> , rectum; |
| <i>bl.</i> , bladder; | <i>r.a.</i> , renal artery; |
| <i>C¹</i> , Cowper's gland No. 1; | <i>r.p.</i> , retractor muscle of penis; |
| <i>C³</i> , Cowper's gland No. 3; | <i>r.v.</i> , renal vein; |
| <i>c.c.</i> , bulb of corpus cavernosum; | <i>t.</i> , testis; |
| <i>cl.</i> , cloacal opening; | <i>u.</i> , ureter; |
| <i>c.s.</i> , bulb of corpus spongiosum; | <i>u.a.</i> , anterior urethra; |
| <i>e.</i> , epididymis; | <i>u.p.</i> , posterior urethra; |
| <i>k.</i> , kidney; | <i>v.c.</i> , vena cava; |
| <i>l.p.</i> , Poupart's ligament; | <i>v.d.</i> , vas deferens. |
| <i>p.p.</i> , penis pouch; | |

Fig. 3. Ovary and adjacent parts (x7½).



REPRODUCTIVE ORGANS.
Enlarged.



PLATE IX.

DISSECTIONS OF MALE ORGANS.

Enlarged.

Fig. 1. Dorsal view of parts surrounding bulbous urethra. Two Cowper's glands removed from left side ($\times 3\frac{1}{2}$).

Fig. 2. Ventral view of bulb of corpus cavernosum and adjoining parts ($\times 6$).

Fig. 3. Longitudinal section of urethra with adjoining parts ($\times 5$).

b.u., bulbous urethra;

b.w., body wall;

c.c., corpus cavernosum;

c.c.m., muscular bulb of corpus cavernosum;

c.s., corpus spongiosum;

c.s.m., muscular bulb of corpus spongiosum;

gl., lymphatic gland;

g.p., glans penis;

i.t., tendon attaching corpus cavernosum to ischium;

l., median line;

m¹, m², m³, small paired muscles at anterior end of bulbous urethra;

p.p., penis pouch;

pr., prepuce;

p.t., tendon running to pubis;

r., rectum;




r.m., rectal muscle;

r.p., retractor muscle of penis;

sp.f., fascia to covering of bulb and to sphincter;

t.f., tendinous fascia connecting bulb and muscle in front of bulbous urethra;

u.a., anterior urethra.



DISSECTIONS OF MALE ORGANS.
Enlarged.

PLATE X.

PARTS OF REPRODUCTIVE, ALIMENTARY, AND RESPIRATORY SYSTEMS.

Enlarged.

Fig. 1. Dissection of base of bladder showing endings of ureters and vasa deferentia (x5).






Fig. 2. Ventral aspect of lungs and trachea (x2½).

Fig. 3. Lateral view of tongue (x3½).

Fig. 4. Base of tongue from above (x10).

Fig. 5. Transverse section of prostate gland (x5).

- az.*, azygos lobe of lung;
- bl.*, bladder;
- c.p.*, coronate papillae;
- c.v.p.*, circumvallate papillae;
- ep.*, epiglottis;
- fg.p.*, fungiform papillae;
- l.l.*, left lobe of lung;
- p.t.*, columnar tissue of prostate gland;
- r.l.a.*, anterior division of right lobe of lung;
- r.l.p.*, posterior division of right lobe of lung;
- u.a.*, anterior urethra;
- v.d.*, vas deferens.



PARTS OF REPRODUCTIVE, ALIMENTARY, AND RESPIRATORY SYSTEMS.
Enlarged.



THE JOURNAL OF THE ROYAL ANTHROPOLOGICAL INSTITUTE
 VOLUME 1, PART 1, 1871
 LONDON: PUBLISHED BY THE INSTITUTE, 1, BEDFORD SQUARE, W.C.
 1871

THE JOURNAL OF THE ROYAL ANTHROPOLOGICAL INSTITUTE
 VOLUME 1, PART 1, 1871
 LONDON: PUBLISHED BY THE INSTITUTE, 1, BEDFORD SQUARE, W.C.
 1871



THE JOURNAL OF THE ROYAL ANTHROPOLOGICAL INSTITUTE
 VOLUME 1, PART 1, 1871
 LONDON: PUBLISHED BY THE INSTITUTE, 1, BEDFORD SQUARE, W.C.
 1871



PLATE XI.

CERVICAL VERTEBRAE.

Five times natural size.

Fig. 1. Atlas, posterior aspect.

Fig. 2. Atlas, anterior aspect.

Fig. 3. Atlas, dorsal aspect.

Figs. 4-6. Seventh cervical from different individuals showing variation in form and occurrence of vertebrarterial foramina.

Fig. 7. Axis, left lateral aspect.

a.s., anterior zygapophysis;

h.t., hypapophysial tubercle;

i.v., intervertebral notch;

l.s., lamina of spinous process;

n.f., nutrient foramen;

o.p., odontoid process;

p.p., pleurapophysis or cervical rib;


p.z., postzygapophysis;

tr., transverse process;


v.c., vertebrarterial canal;

v.c.g., vertebrarterial canal reduced to a groove.


1




2




4




3




5



7



6



CERVICAL VERTEBRAE.
Five times natural size.




PLATE XII.

BONES OF STERNUM, HIND LEG, AND THORACIC REGION.

Five times natural size.

- Fig. 1. Right femur, posterior aspect.
Fig. 2. Sternum and attachment of ribs.
Fig. 3. Right tibia and fibula with adjoining part of femur, outer aspect.
Fig. 4. First and second ribs and attachments, outer aspect.
Fig. 5. Clavicle.


*c*⁷., seventh cervical vertebra;
f., fibula;
fb., fabella;
g.t., greater trochanter;
h., head of femur;
i.r., intertrochanteric ridge;
l.t., lesser trochanter;
m., manubrium of sternum;
p., osseous patella;
*r*¹., first rib;
s., shaft of femur;
t., tibia;
t.t., rudimentary third trochanter of femur;
*t*¹., first thoracic vertebra;
*t*²., second thoracic vertebra;
x., xiphisternum;
x.c., xiphoid cartilage;
2-7., costal attachments of second to seventh ribs.



BONES OF STERNUM, HIND LEG, AND THORACIC REGION.
Five times natural size.



THE JOURNAL
OF THE
AMERICAN MEDICAL ASSOCIATION
PUBLISHED WEEKLY
CHICAGO, ILL., MAY 1, 1919
Vol. 34, No. 19
CONTENTS
ORIGINAL ARTICLES
The Effect of the
Hormones of the
Adrenal Gland on the
Growth of the
Embryo of the
Rabbit
The Effect of the
Hormones of the
Adrenal Gland on the
Growth of the
Embryo of the
Rabbit
The Effect of the
Hormones of the
Adrenal Gland on the
Growth of the
Embryo of the
Rabbit



2



PLATE XIII.

SCAPULA, PELVIS, AND LUMBAR VERTEBRAE.

Five times natural size.

Fig. 1. Lumbar vertebrae, left lateral aspect.

Fig. 2. Lumbar vertebrae, ventral aspect.

Fig. 3. Scapula, right lateral aspect.

Fig. 4. Pelvis, right lateral aspect.


ac., acromion;

co., coracoid process;

i., ilium;

ip.t., iliopectineal tubercle;


i.t., ischial tuberosity.




1



2



3



4

SCAPULA, PELVIS, AND LUMBAR VERTEBRAE.
Five times natural size.



PLATE XIV.

SACRAL AND CAUDAL VERTEBRAE.

Five times natural size.

Fig. 1. Sacral and first six caudal vertebrae, dorsal aspect.

Fig. 2. Second to ninth caudal vertebrae, ventral aspect.

c^2, c^3, c^4, c^5 , caudal vertebrae;

ch., chevron bone;

s^1 , first sacral vertebra.



SACRAL AND CAUDAL VERTEBRAE.
Five times natural size.

100



Fig. 1.

Anterior view of the bone.

- 1. Head of the bone.
- 2. Neck of the bone.
- 3. Greater trochanter.
- 4. Lesser trochanter.
- 5. Shaft of the bone.
- 6. Epicondyle of the humerus.
- 7. Epitrochlear fossa.
- 8. Epicondyle of the ulna.
- 9. Epitrochlear fossa.
- 10. Epicondyle of the radius.
- 11. Epitrochlear fossa.
- 12. Epicondyle of the ulna.
- 13. Epitrochlear fossa.
- 14. Epicondyle of the radius.
- 15. Epitrochlear fossa.



Fig. 2.



Fig. 3.

PLATE XV.

HUMERUS, ULNA, AND RADIUS.

Five times natural size.

Fig. 1. Right humerus, posterior or caudal aspect.

Fig. 2. Right humerus, anterior or cranial aspect.

Fig. 3. Right radius and ulna, outer lateral aspect.

Fig. 4. Right ulna, anterior aspect.

ca., capitellum;

d.r., deltoid ridge;

e.f., epicondylar foramen;

g.s., greater sigmoid cavity;

g.t., greater tuberosity;

h., head of humerus;

i.a., inner articular surface of distal end of humerus;

i.c., inner condyle;

l.t., lesser tuberosity;

ol., olecranon and fossa;

r., radius;


s.r., supinator ridge;

tr., trochlea;


u., ulna.



1



2



3



4

HUMERUS, ULNA, AND RADIUS.
Five times natural size.



PLATE XVI.

MANUS AND PES.

Enlarged.

Fig. 1. Right manus, anterior aspect (x5).

Fig. 2. Right pes, anterior aspect (x5).

Fig. 3. Left astragalus, posterior aspect (x10).

Fig. 4. Left astragalus, anterior aspect (x10).

Fig. 5. Right os calcis, anterior aspect (x10).

a.l., facet for ligament to astragalus;

as., astragalus;

c., cuneiform;

cu., cuboid;

ec., ectocuneiform;

e.f., ectal facet;

en., entocuneiform;

f.f., fibular facet;

i.m., inner malleolar facet;

l., lunar;

m., magnum;

me., mesocuneiform;

na., navicular;

n.f., navicular facet;

o.c., os calcis;

p., pisiform;

r., radius;

sc., scaphoid;


s.f., sustentacular facet;

t.f., tibial facet;

tr., trapezium;

tz., trapezoid;

u., ulna.



1

2



3

4

5

MANUS AND PES.
Enlarged.

1870



Alfred H. Huxley

1870

1870


PLATE XVII.

LONGITUDINAL SECTION OF SKULL.

Five times natural size.

bo., basioccipital;
bs., basisphenoid;
c.c., carotid canal;
cb.f., cerebellar fossa;
cd.f., condyloid foramen;
c.f., cerebral fossa;
c.p., cribriform plate;
et., ethmoturbinal;
f., frontal;
f.c., carotid foramen;
f.f., floccular foramen;
f.o., foramen ovale;
f.s., frontal sinus;
j.f., jugular foramen;
m., maxilla;
me., mesethmoid;

m.f., mastoid foramen;
mt., maxillo-turbinal;
n., nasal;
n.c., nasal cartilage;
nt., nasoturbinal;
o.f., olfactory fossa;
p., palatine;
per., periotic;
p.f., pituitary fossa;
p.p.m., palatine process of premaxillary;
ps., presphenoid;
pl., pterygoid;
s.f., subsquamosal foramen;
t.c., transverse canal;
v., vomer.



LONGITUDINAL SECTION OF SKULL.
Five times natural size.


PLATE XVII.

LONGITUDINAL SECTION OF SKULL.

Five times natural size.

bo., basioccipital;
bs., basisphenoid;
cc., carotid canal;
cb.f., cerebellar fossa;
cd.f., condyloid foramen;
c.f., cerebral fossa;
c.p., cribriform plate;
el., ethmoturbinal;
f., frontal;
f.c., carotid foramen;
f.f., floccular foramen;
f.o., foramen ovale;
f.s., frontal sinus;
j.f., jugular foramen;
m., maxilla;
me., mesethmoid;

m.f., mastoid foramen;
mt., maxillo-turbinal;
n., nasal;
n.c., nasal cartilage;
nt., nasoturbinal;
o.f., olfactory fossa;
p., palatine;
per., periotic;
p.f., pituitary fossa;
p.pm., palatine process of premaxillary;
ps., presphenoid;
pt., pterygoid;
s.f., subsquamosal foramen;
t.c., transverse canal;
v., vomer.



LONGITUDINAL SECTION OF SKULL.
Five times natural size.

1. The first part of the paper is devoted to a general discussion of the problem of the existence of solutions of the system of equations (1) for arbitrary values of the parameters α and β . It is shown that the system has solutions for arbitrary values of the parameters α and β if and only if the condition $\alpha + \beta > 0$ is satisfied.

2. In the second part of the paper the problem of the existence of solutions of the system of equations (1) for arbitrary values of the parameters α and β is solved. It is shown that the system has solutions for arbitrary values of the parameters α and β if and only if the condition $\alpha + \beta > 0$ is satisfied.

3. In the third part of the paper the problem of the existence of solutions of the system of equations (1) for arbitrary values of the parameters α and β is solved. It is shown that the system has solutions for arbitrary values of the parameters α and β if and only if the condition $\alpha + \beta > 0$ is satisfied.

4. In the fourth part of the paper the problem of the existence of solutions of the system of equations (1) for arbitrary values of the parameters α and β is solved. It is shown that the system has solutions for arbitrary values of the parameters α and β if and only if the condition $\alpha + \beta > 0$ is satisfied.

5. In the fifth part of the paper the problem of the existence of solutions of the system of equations (1) for arbitrary values of the parameters α and β is solved. It is shown that the system has solutions for arbitrary values of the parameters α and β if and only if the condition $\alpha + \beta > 0$ is satisfied.

The author is grateful to the Ministry of Higher Education of the USSR for the financial support of this work.

PLATE XVIII.
DENTITION AND HARD PALATE.
Enlarged.

- Fig. 1. First upper molar, inner lateral aspect (x10).
- Fig. 2. Left lower molar series from above (x10).
- Fig. 3. Hard palate (x5).
- Fig. 4. Left upper molar series (x10).
- Fig. 5. Left lower molar series, outer lateral aspect (x10).



1




2



3



4



5

DENTITION AND HARD PALATE.
Enlarged.



Figure 1. A person in a small boat on a body of water, with a large, dark, rocky island in the background.



Figure 2. A person in a small boat on a body of water, with a large, dark, rocky island in the background.

PLATE XIX.

Fig. 1. Mounted Skeleton of *Nesogale dobsoni*.

Fig. 2. Mounted Skeleton of *Cænolestes obscurus*.



1. MOUNTED SKELETON OF NESOGALE DOBSONI.



2. MOUNTED SKELETON OF CAENOLESTES OBSCURUS.

XX 1914

January 1st
 January 2nd
 January 3rd
 January 4th

January 5th
 January 6th

January 7th
 January 8th

January 9th
 January 10th

January 11th
 January 12th

January 13th
 January 14th

January 15th
 January 16th

January 17th
 January 18th

January 19th
 January 20th

January 21st
 January 22nd

January 23rd
 January 24th

January 25th
 January 26th

January 27th
 January 28th

January 29th
 January 30th

January 31st
 February 1st

February 2nd
 February 3rd

February 4th
 February 5th

February 6th
 February 7th

February 8th
 February 9th

February 10th
 February 11th

February 12th
 February 13th

February 14th
 February 15th

February 16th
 February 17th

February 18th
 February 19th

February 20th
 February 21st

February 22nd
 February 23rd

February 24th
 February 25th

February 26th
 February 27th

February 28th
 February 29th

PLATE XX.


SKULL OF CÆNOLESTES.

Adapted from Miss Dederer.

Fig. 1. Palatal aspect.

Fig. 2. Lateral aspect.

<i>ang.</i> , angular process of mandible;	<i>m.f.</i> , mental foramen;
<i>a.o.v.</i> , antorbital vacuity;	<i>Ms.</i> , mastoid;
<i>a.pl.vac.</i> , anterior palatine vacuity;	<i>ms.f.</i> , masseteric foramen;
<i>As.</i> , alisphenoid;	<i>Mx.</i> , maxilla;
<i>As.bl.</i> , alisphenoid bulla;	<i>Na.</i> , nasal;
<i>Bo.</i> , basioccipital;	<i>Os.</i> , orbitosphenoid;
<i>Bs.</i> , basisphenoid;	<i>Pa.</i> , parietal;
<i>c.</i> , mandibular condyle;	<i>Per.</i> , periotic;
<i>car.can.</i> , carotid canal;	<i>p.gl.f.</i> , post glenoid foramen;
<i>car.f.</i> , carotid foramen;	<i>Pl.</i> , palatine;
<i>c.f.</i> , condyloid foramen;	<i>p.l.f.</i> , postero-lateral foramen;
<i>cor.pr.</i> , coronoid process;	<i>pl.r.</i> , palatal ridge;
<i>eust.f.</i> , eustachian opening;	<i>Pmx.</i> , premaxilla;
<i>Exo.</i> , exoccipital;	<i>p.pl.vac.</i> , posterior palatine vacuity;
<i>f.l.a.</i> , sphenorbital foramen;	<i>Ps.</i> , presphenoid;
<i>f.l.p.</i> , foramen lacerum posterum;	<i>Pt.</i> , pterygoid;
<i>f.ov.</i> , foramen ovale;	<i>pt.p.</i> , pterygoid process of palatine;
<i>Fr.</i> , frontal;	<i>s.sq.f.</i> , subsquamosal foramen;
<i>f.v.f.</i> , venous foramen in frontal;	<i>st.m.f.</i> , stylomastoid foramen;
<i>L.</i> , lachrymal;	<i>tr.can.</i> , transverse canal;
<i>l.d.</i> , lachrymal duct;	<i>Ty.</i> , tympanic;
<i>Ma.</i> , malar;	<i>V².</i> , infraorbital foramen.



SKULL OF CAENOLESTES.
Adapted from Miss Dederer.



PLATE XXI.

DORSAL AND LATERAL VIEWS OF BRAIN.


Five times natural size.

Fig. 1. View of the dorsal surface of the brain.

b.ol., bulbus olfactorius;
cb., body of the cerebellum;
floc., flocculus;
f.rh., fissura rhinalis;
s.orb., sulcus orbitalis.

Fig. 2. View of the brain from the right side.

b.ol., bulbus olfactorius;
cb., body of the cerebellum;
f.erh., fissura endorhinalis;
floc., flocculus;
f.rh., fissura rhinalis;
l.pir., lobus piriformis;
s.orb., sulcus orbitalis;
t.o., tuberculum olfactorium;
tr.ol.lat., tractus olfactorius lateralis.



1



2

DORSAL AND LATERAL SURFACES OF BRAIN.
Five times natural size.



PLATE XXII.

VENTRAL SURFACE OF BRAIN.

Five times natural size.

b.ol., bulbus olfactorius;

f.erh., fissura endorhinalis;

floc., flocculus;

hy., hypophysis;

II to XII., cranial nerves;


l.pir., lobus piriformis;

pyr., pyramid;

t.o., tuberculum olfactorium;

tr.ol.lat., tractus olfactorius lateralis;

trap., corpus trapezoideum.



VENTRAL VIEW OF BRAIN.
Five times natural size.

163
FIELD MUSEUM OF NATURAL HISTORY

PUBLICATION 222

ZOOLOGICAL SERIES

VOL. XIV, No. 2

REVIEW OF LIVING CAENOLESTIDS WITH
DESCRIPTION OF A NEW GENUS
FROM CHILE

BY

WILFRED H. OSGOOD

Curator, Department of Zoology

REPORTS ON RESULTS OF
THE CAPTAIN MARSHALL FIELD EXPEDITIONS



CHICAGO, U. S. A.

October 20, 1924

REVIEW OF LIVING CAENOLESTIDS WITH DESCRIPTION OF A NEW GENUS FROM CHILE

BY WILFRED H. OSGOOD

Among many interesting mammals obtained by the Captain Marshall Field Expedition to Chile in 1923, the most important novelty is a small marsupial representing an undescribed genus related to *Caenolestes* and *Orolestes*. It was found in the forests of the island of Chiloe nearly two thousand miles from the known habitat of its nearest relatives. Thus it not only provides valuable material for further study of an archaic group of mammals, but constitutes an important addition to the fauna of Chile.

In 1911, when an expedition from Field Museum found *Caenolestes obscurus* living in the mountains of Venezuela, the interesting group to which it belongs was represented in the museums of the world by only half a dozen imperfect native-collected specimens. The fresh material obtained at that time, including complete skeletons and soft parts in preservative for dissection, was considered so important it was made the subject of a special monographic study (antea, pp. 1-162). Continued work by trained American collectors in South American fields has resulted in an unexpectedly large increase in knowledge of the group. At the present time, including the form described below, there are recorded not less than seven species of living caenolestids belonging to three well-marked genera.

The original example of *Caenolestes fuliginosus* from Ecuador was described by Tomes under the generic name *Hyracodon* in 1863 and no further specimens appeared for more than thirty years, when the British Museum received four collected in 1895 by natives employed by Mr. G. D. Child, on the La Selva Hacienda, near Bogota, Colombia. One of these was acquired by the American Museum of Natural History and three remained in the British Museum, one being the type of the new species *Caenolestes obscurus*. Another long period ensued until 1911, when the present writer secured a series of eleven specimens in Venezuela and Colombia not far from the Bogota region. Later, in the same year, Mr. S. N. Rhoads collected for the Academy of Natural Sciences of Philadelphia two specimens at Hacienda Garzon, Mount

Pichincha, Ecuador, in the region from which came the original type of *C. fuliginosus*. Two additional specimens were collected at Gualea, Ecuador, by Mr. Walter Goodfellow and presented to the British Museum in 1914. Then, in 1915, Mr. Edmund Heller found caenolestids in abundance in a new region about the upper waters of the Urubamba River in Peru. He secured a large series numbering thirty or more specimens upon which was founded the new genus and species *Orolestes inca*. This series is in the U. S. National Museum at Washington. In 1920, Mr. H. E. Anthony, of the American Museum of Natural History, discovered a well-marked new species, *Caenolestes caniventer*, in the province of Oro, Ecuador, and in 1922 the same author described a third species from Ecuador, *Caenolestes tatei*, collected by Mr. G. H. H. Tate in the province of Azuay. Series of both these species, as well as additional specimens of *C. fuliginosus*, have been obtained recently in Ecuador by Mr. Anthony and Mr. Tate. Meanwhile, Dr. Einar Lönnberg, in 1921, recorded six specimens from various localities in Ecuador obtained through Consul L. Söderström. Finally, as stated above, a third genus has now been found in Chile.

In connection with the description of this genus, it has seemed desirable to present the brief statement of the systematic status of the whole group which follows. Specimens of forms not in the Field Museum, loaned by the American Museum of Natural History and the U. S. National Museum, are gratefully acknowledged.

Caenolestes Thomas.

(Plate XXIII, figs. 1, 1a, 1b.)

Caenolestes Thomas, Ann. & Mag. Nat. Hist., (6), 16, pp. 367-368, Nov. 1895—type, *Hyracodon fuliginosus* Tomes 1863, from Ecuador.

Characters. Size small; form soricine; tail longer than head and body, non-prehensile; feet pentadactyl eleutherodactyl; marsupium absent; mammae $\frac{3}{4}=4$, inguinal; habits cursorial and insectivorous; color uniform. Skull with elongate facial region; braincase large, deep and thin-walled, without external ridges or crests; interorbital and lacrymal regions swollen; nasals long and slender, abruptly expanded posteriorly; a marked vacuity between the nasals, frontals and maxillaries; zygomata slender, their roots rising at the plane of the third molar; palate highly fenestrate; tympanic annular; a carotid canal between the petrous periotic and the basioccipital. Teeth numerous and highly differentiated; anterior pair of incisors terminal and of the "diprotodont" type; second and third pairs of upper incisors in con-

tact with each other, lateral in position and laterally compressed in form, their functional edges entire, forming cutting blades slightly inclined forward; last pair of incisors similar to others but smaller and separated from them by a slight space; canine well-developed, falcate, single-rooted; anterior molariform teeth quadrate bunolophodont with a basal external cingulum; a small metaconule usually present. Tooth formula: I. $\frac{1}{1}$; C. $\frac{1}{1}$; Pm. $\frac{1}{1}$; M. $\frac{4}{4}$;=46.

Remarks. Four species of the genus *Caenolestes* have had nominal recognition, three from Ecuador and one from Colombia. All of these are obviously very closely related and the material has been so limited that characterizations of them have been rather unsatisfactory. *C. fuliginosus*, *C. obscurus* and *C. tatei* are so much alike in color and all general characters that further study with more material may prove them to be no more than subspecifically separable. Comparison has been made of a small series of *C. obscurus* from Paramo de Tama, Colombia-Venezuela boundary with a half dozen specimens from Ecuador, representing *fuliginosus*, *caniventer* and *tatei*. Collectively, the specimens of *obscurus* may be distinguished from those from Ecuador by the slightly more slender facial part of the skull and by the unnotched cutting edges of the second and third incisors. In one specimen of *obscurus*, the second incisor shows a faint indication of a notch, but in all others it is entire, whereas in all Ecuadorean specimens examined, this tooth is definitely notched and in some cases the third incisor is similarly notched. This character, to which attention has been called by Mr. Anthony, is accompanied by a somewhat greater forward rotation of the tooth and by a closer contact of the second incisor with the first. The supposed large size and the closing of the anteorbital vacuities mentioned by Thomas and Lönnberg as characters of *obscurus* are not appreciable in the material at hand. In the largest male of *obscurus* from Paramo de Tama, having a skull length of 32.5 mm., the combined length of the three anterior molars is only 5 mm. In the largest male from Ecuador (type of *caniventer*), with a skull length of 33.5, the three principal molars measure 5.4. In all specimens of *obscurus* from Paramo de Tama and, likewise, in all Ecuadorean skulls examined, the anteorbital vacuities are open. The type of *C. tatei* is a very small female and rather dark in color, but there is much variation in these respects among other Ecuadorean specimens. *C. caniventer* seems to be well characterized by average large size and by its more sooty, less brownish, coloration on the upperparts and its decidedly pale grayish underparts. A peculiar circumstance is the occurrence of both *C. tatei* and *C. caniventer* at the same locality (Mol-

leturo). It is greatly to be hoped that Mr. Anthony, with his field experience and the considerable series of *Caenolestes* which he is accumulating from Ecuador, will be able to clear up all uncertain points in connection with the variation, distribution, and pelage changes of these forms.*

Orolestes Thomas.

(Plate XXIII, figs. 2, 2a, 2b.)

Orolestes Thomas, Smiths. Misc. Coll., 68, No. 4, p. 3, Apr. 10, 1917; Proc. U. S. Nat. Mus., 58, pp. 244-247, pl. 14, fig. 5, 1920—type, *Orolestes inca* Thomas, from Torontoy, Peru.

Characters. External form, size and coloration much as in *Caenolestes*; cranial characters also generally similar to those of *Caenolestes*; jugal broader and more flattened anteriorly; external boundaries of infraorbital foramina slightly farther back. Antemolar teeth showing various differences from those of *Caenolestes*; fourth upper incisor sharply pointed and slightly retrorse, separated from the third and from the canine by a space equal to or greater than its alveolar length; canine double-rooted and premolariform, only slightly higher than incisors; first premolar a small blunt peg scarcely projecting above the gums, one third to one fourth as large as in *Caenolestes*; molariform teeth practically as in *Caenolestes*; fourth upper molar very small, scarcely more than half as large as in *Caenolestes*; lower canine smaller than incisors; fourth lower molar reduced in size.

Remarks. *Orolestes* is characterized principally by its low double-rooted canine and its minute first premolar. The food habits of *Orolestes* and *Caenolestes* are presumably the same and it is difficult to find teleological grounds for a pronounced difference in their canines.

Only one species (*O. inca*) of *Orolestes* has thus far been discovered. The original series from the upper Urubamba region of Peru has not been duplicated. Mr. Heller, who collected this series, has recently worked again in Peru somewhat north of the Urubamba valley in the upper Marañón and Huallaga region, but failed to obtain any caenolestids.

One of the specimens of *Orolestes* examined (No. 194948 U. S. N. M.) shows an interesting abnormality in the fusion of the lower canine and the first premolar, forming an imperfectly three-rooted tooth

*Since this was written a fourth species from Ecuador has been described by Mr. Anthony (*Caenolestes convelatus* Anthony, Amer. Mus. Novit., No. 120, pp. 1-3, June 21, 1924). This is characterized as "Most like *caniventer* in external appearance but skull lacking facial vacuities and dentition heavier."

with a tendency to become two-rooted through the coalescence of the root of the canine with the anterior root of the premolar.

Rhyncholestes genus novum.

(Plate XXIII, figs. 3, 3a, 3b, 3c.)

Type, *Rhyncholestes raphanurus* sp. nov.

Characters. External appearance much as in *Caenolestes* and *Orolestes*, but tail shortened and (at least periodically in the male) thickened at the base, making it elongate radiceform with a basal diameter of 7 mm. or more. Mammae five, two inguinal on each side and one midventral slightly in advance of them. Skull very narrow and elongate, especially the facial part, the ratio of the zygomatic breadth to the total skull-length being only 41.9* or about equal to that in the peramelids. Antemolar teeth widely spaced; second, third and fourth upper incisors with a broad anterior blade and a blunt, slightly curved, posterior cusp, the two parts separated by a deep cleft; upper canines single-rooted, strictly caniniform in the male, but somewhat premolari-form in the female, in which there is a faint anterior cusp and a marked posterior one; principal molariform teeth much as in *Caenolestes* and *Orolestes*, but (at least in the two known specimens) without any apparent development of metaconules; fourth molar minute, even more reduced than in *Orolestes*. Posterior palatine vacuities without any median bony division except for slightly pointed prolongations of the maxillary and the palatine, thus leaving the deeply-channeled ventral surface of the vomer exposed for a considerable part of its length. Anteorbital vacuities much as in *Caenolestes*; infraorbital foramen opening over middle of first molar, divided by a bony bar forming a double foramen, the lower and larger division of which opens forward as usual and the upper and smaller, elliptical in shape, opens laterally in the exposed wall of the maxillary. Zygomata relatively broad and flattened throughout and channeled anteriorly on the exterior surface; a downwardly directed knoblike process at the junction of the jugal and the maxillary.

Skeletal characters mostly as in *Caenolestes*. Humerus relatively shorter, the ratio of its length to the ulna 67 as against 79 in *Caenolestes*; deltoid ridge occupying fully half the length of the humerus; outer epicondylar or supinator ridge not meeting the shaft of the humerus gradually but abruptly with a tendency to the development of a hooked process; metapodials relatively short; first and second caudal

*This ratio in *Caenolestes* is 47.7; in *Perameles nasuta* it is 39.7 and in *P. bougainvillei* 43.2; in other marsupials it is much greater (see antea, p. 99).

vertebrae with spinous processes; total number of caudals reduced to 22 (27 in *Caenolestes*).

***Rhyncholestes raphanurus* sp. nov.**

Type from forests at mouth of Rio Inio, south end of Chiloe Island, Chile. S. lat. $43^{\circ} 20'$. No. 22422 Field Museum of Natural History—skin and skeleton. Adult female. Collected Jan. 12, 1923 by Wilfred H. Osgood. Original No. 5500.

Characters. Color and character of pelage practically as in *Caenolestes fuliginosus* and *C. obscurus*, perhaps slightly darker; rich dark brown above and below without markings of any kind; tail nearly black all around, darker and less brownish than in any of the northern forms examined. In the type, which is a female, the tail is self-colored, but in the male, the skin of which could not be preserved, it was noted in the field that the tail was bluish black except on the terminal third which was mottled black and white above and below and with the tip entirely white. General characters as given in the generic diagnosis.

Measurements. Adult male and female, respectively; Total length 204, 175; tail vertebrae 78, 65; hind foot 21, 19.5; ear from notch 11, 11; diameter of tail at base 7, 5. Skulls: Greatest length 34, 30.3; basal length 33.3, 29.6; zygomatic breadth 14.7, 12.5; mastoid breadth 11.1, 10.2; length of nasals 18, 15.8; greatest breadth of nasals 3, 2; least interorbital breadth 6.9, 7; length of palate from gnathion 20.3, 17.5; anterior palatal foramina 9.7, 7.3; palatal vacuities 6, 5.8; front of upper canine to back of last molar 13.9, 11.9; combined length of four upper molars 5.4, 5.2; combined length of three lateral incisors 3.9, 3.5; length of bone of mandible from condyle 20.18; exposed length of median lower incisor 6.8, 6.1.

Remarks. This genus is well characterized in numerous ways. In the dentition, the most striking peculiarity is seen in the unequally bifid lateral upper incisors, the interpretation of which has many interesting possibilities. The canines also are of especial interest since they provide at least a hypothetical intermediate stage between those of *Orolestes* and *Caenolestes*. In view of the diversity in the dentition of the three genera of caenolestids now living, it is evident that considerable changes may have taken place in a relatively short time, and it is perhaps idle to speculate on the history of particular teeth over long geological periods. However, if the bifid lateral incisors of *Rhyncholestes* and the premolariform canines of *Orolestes* (and *Rhyncholestes*) be regarded as primitive, they fall into a long list of other primitive characters possessed by these forms. Taking them together, incisors

and canines, the presumption that they are primitive is very strong. The incisors have no parallel among modern marsupials and if they were present in any of the Miocene caenolestids, it is not suggested by the scanty material yet available; but they could be derived easily from some types of early insectivores or from the triconodont teeth so common among Mesozoic mammals. The simple three-cusped type of tooth now seen in the anterior premolars of caenolestids could produce the lateral upper incisor of *Rhyncholestes* by reflexing the principal cusp, bringing the anterior face of this cusp toward the horizontal and approximating the apices of the central and the posterior cusps thus deepening the cleft between them and giving it a less regular form, meanwhile elevating the front of the tooth and reducing the anterior cusp to a mere angle—in other words, by rotating the whole tooth backwards. Continuing this process and eliminating the posterior cusp might produce the simple hatchet-shaped blade seen in the upper lateral incisors of *Caenolestes obscurus*. The slight sulcus near the posterior angle of these teeth, noted by Anthony in certain Ecuadorean examples of *Caenolestes*, may be a vestige of the process, but this is uncertain. The roots of the lateral incisors are broad and heavy and also suggest derivation from a two-rooted premolariform antecedent. Examination of the teeth by transmitted light shows a longitudinal deficiency of dentine above the center of the root such as might be expected after the coalescence of two roots.

The condition of the canines in *Rhyncholestes* leaves little room for doubt that the two-rooted premolariform canine seen in *Orolestes* is the forerunner of the single-rooted scalpriform canine of *Caenolestes*. In *Rhyncholestes* both types are represented, one in the female, the more primitive, and the other in the male, more specialized. The tooth in the female has the two roots partly coalescent and its apex is somewhat elevated, but it retains a well-defined posterior cusp and a trace of an anterior one. In the male it has full single-rooted caniniform development. Viewing this tooth in the female in connection with the two premolars behind it and the three incisors in front of it, the essential similarity of the whole series is striking, and the belief that all have been derived from the same type of tooth is scarcely to be avoided. Two-rooted premolariform canines, it is interesting to note, are found in other marsupials only in *Choeropus*, one of the peramelids.

An unexpected external character of *Rhyncholestes* is the possession of five mammae instead of four. The fifth is in median position somewhat in advance of the others, which are paired as in *Caenolestes*. This constitutes a similarity to the didelphids, the only other group

previously known to have mammae in odd numbers, but doubtless it is not especially significant of relationship.

The discovery of this new genus of caenolestids at sea level, in the forests of southern Chile, enormously extends the range of the group and renders it probable that further forms will be found in the great areas intervening between the relatively few spots at which specimens have been taken. Only two specimens of *Rhyncholestes* were obtained, fortunately male and female, both fully adult. The locality was in heavy temperate forests near the mouth of the Rio Inio at the extreme southern end of Chiloe Island. Here it was only after much trapping that the first specimen, the female, was taken. The vicinity was thereupon carefully re-examined and additional traps placed in all promising situations with the result that at the end of a week's time, the male was secured. An enforced absence had made it impossible to visit the traps for several days, and the male specimen was found in somewhat decomposed condition so that, although color notes and measurements could be taken, the skin could not be saved.

During some five weeks of constant trapping on Chiloe Island no other specimens of *Rhyncholestes* were taken, so it is evidently rare and difficult to obtain. Doubtless it is not confined to Chiloe Island, but will be found also on the forested mainland of southern Chile. Various localities in this region were visited, however, and although conditions seemed favorable, all efforts to secure additional specimens were unsuccessful.

REFERENCES

ANTHONY, H. E.

1921. Preliminary Report on Ecuadorean Mammals. No. 1. Amer. Mus. Novit., No. 20, p. 6, Nov. 3, 1921.

Original description of *Caenolestes caniventer*.

1923. Preliminary Report on Ecuadorean Mammals. No. 3. Amer. Mus. Novit., No. 55, pp. 1-3, Jan. 31, 1923.

Original description of *Caenolestes tatei*.

1924. Preliminary Report on Ecuadorean Mammals. No. 5. Amer. Mus. Novit., No. 120, pp. 1-3, June 21, 1924.

Original description of *Caenolestes convelatus*.

GREGORY, W. K.

1922. On the "Habitus" and "Heritage" of *Caenolestes*. Journ. Mamm., 3, pp. 106-114, May 1922.

LONGMAN, HEBER A.

1924. The Zoogeography of Marsupials. With notes on the origin of the Australian fauna. Mem. Queensland Mus., 8, Part 1, pp. 1-15, Jan. 30, 1924. Includes discussion of relationships of *Caenolestes*.

LÖNNBERG, EINAR.

1921. A Second Contribution to the Mammalogy of Ecuador with some Remarks on *Caenolestes*. Arkiv. för Zool., 14, No. 4, pp. 1-104, pl. 1, 1921.



1



2



3




1a




2a




3a



2b



1b



3b

3c

Carl F. Gronemann, del.

1, 1A, 1B. *Caenolestes obscurus*.

F. M. N. H. No. 18602, Male.

3, 3A, 3B, 3C. *Rhyncholestes raphanurus*.

F. M. N. H. No. 22423, Male; No. 22422, 3C Female.

2, 2A, 2B. *Orolestes inca*.

U. S. N. M. No. 194428, Male.

(X 1 $\frac{1}{2}$.)

173
FIELD MUSEUM OF NATURAL HISTORY

PUBLICATION 224

ZOOLOGICAL SERIES

VOL. XIV, No. 3

THE BRAINS OF THE SOUTH AMERICAN
MARSUPIALS CAENOLESTES
AND OROLESTES

BY

JEANNETTE BROWN OBENCHAIN

Hull Laboratory of Anatomy, University of Chicago

WILFRED H. OSGOOD
Curator, Department of Zoology



CHICAGO, U. S. A.

January 26, 1925

THE BRAINS OF THE SOUTH AMERICAN MARSUPIALS CAENOLESTES AND OROLESTES

BY JEANNETTE BROWN OBENCHAIN

CONTENTS

	PAGE
Introduction	175
External anatomy of the brains of <i>Caenolestes</i> and <i>Orolestes</i>	178
Internal anatomy of the cerebral hemisphere of <i>Caenolestes</i>	188
Primary olfactory area	189
Secondary olfactory areas	189
Anterior olfactory nucleus	191
Tuberculum olfactorium	195
Septal region	199
Pyriform lobe	200
Hippocampal formation	206
Corpus striatum	220
Neopallium	221
General considerations	222
Summary	225
Acknowledgments	227
Bibliography	228
Abbreviations	231

INTRODUCTION

Caenolestes and *Orolestes* are tiny shrewlike marsupials five inches in length of head and body exclusive of the slender tail. They are natives of high Andean forests from Venezuela southward into Peru, and although discovered in 1863, have not been known to science, except by imperfect material, until recent years. Their peculiar dentition has made their assignment to one or the other of the marsupial suborders (Diprotodontia and Polyprotodontia) a matter of much uncertainty. This question remains even yet, perhaps, in suspense, since the one brain character of evidential value seems to reverse the latest conclusion, based on careful sifting of all non-nervous characters (Osgood, 1921; Obenchain, 1923).

Aside from its bearing upon classification, the study of the brains of these species is warranted by many other considerations. These small marsupials, surviving members of an ancient American group, are characterized by the retention of many primitive features, by the absence of any great degree of specialization, by few non-marsupial characters, by numerous resemblances to the modern peramelids (Australian polyprotodonts), and by lack of affinity to the other living American marsupials, the polyprotodont *Didelphidae* (Osgood, 1921, pp. 150-151).

Material.—The material which forms the basis of the present study consists of the brains of three female specimens, one of *Caenolestes obscurus* and two of *Orolestes inca*. The first brain came from an adult female of *Caenolestes* collected in 1911 in the Colombia-Venezuela boundary region by Dr. Wilfred H. Osgood of the Field Museum of Natural History; it was examined by Dr. C. Judson Herrick, who described and figured its external surface in a short paper (1921, *antea*, pp. 157-162, 3 figures) appended to Dr. Osgood's long monograph on the anatomy and zoological position of *Caenolestes*. Dr. Herrick later offered me the opportunity of studying this brain, if I could convert it into serial sections. The other two brains, from adult females of *Orolestes inca*, were collected by Mr. E. Heller of the Yale National Geographic Society Peruvian Expedition of 1914-15, and were loaned by the U. S. National Museum to the Field Museum to be transferred to me for the purpose of this study. The first specimen alone (*Caenolestes*) has been sectioned and stained. The description of the internal anatomy is therefore based entirely on this one transverse series, which, although somewhat imperfect because of incomplete fixation within the unopened skull, has stained with unexpected brilliancy by the iron-haematoxylin method. This series, owing to the fact that both cells and fibers have been stained, serves remarkably well, all things considered, for the study here undertaken. Since the brain was sectioned whole, its medial surface is known only by means of wax and linear reconstructions made from the sections. These are believed to be reasonably trustworthy, but since they are made from an imperfect series, additional corroboration was greatly to be desired. Therefore the two brains of *Orolestes* were especially welcome. I take this opportunity to thank Mr. W. J. Owen of the Anatomical Institute of Melbourne, Australia, for his kind assistance in the attempt to remove them as nearly intact as possible from the skulls. The smaller of the two brains (No. 194948) was divided by a sagittal cut, the right hemisphere was detached from the brain stem, and drawings

of the medial surface of both the whole brain and the detached hemisphere were made by Mr. Kenji Toda, of the University of Chicago, to whom I am indebted for the five beautiful and accurate air brush pictures of the brain of *Orolestes* (Figs. 1-5). The larger brain (No. 194921) was used for the external views, with correction and supplement from the other where necessary, as in the case of the parafloccular lobes, which the larger brain had lost. The business of clearing these small and fragile brains from membranes and coagulated cerebro-spinal fluid was both tedious and difficult, but fairly successful in the end, and the figures are offered as faithful representations of the actual specimens.

Three air brush drawings of the brain of *Caenolestes* by Mr. A. B. Streedain, formerly of the Department of Anatomy, also may be consulted (Herrick, *antea*, pls. XXI-XXII). This brain, owing to the difficulty of removing membranes and débris, was in somewhat less favorable condition for drawing than were the other two. This applies particularly to the ventral surface, which was further marred by a considerable hole due to faulty fixation.

All three specimens were originally preserved whole, with only a ventral longitudinal opening of the body, in 10% formalin, and subsequently transferred to alcohol. In the case of *Caenolestes*, the brain, when freed from the skull, ballooned out with a glistening white surface, but later shrunk and darkened. The other two brains (*Orolestes*) were quite rigid when uncovered, and presented a dull, light brown surface. All three completely filled the skull. It is hoped that the two brains of *Orolestes* may be successfully converted into sagittal and horizontal series.

Method.—The brain of *Caenolestes*, infiltrated in 42 degrees paraffine and blocked in 56 degree paraffine (to minimize brittleness), was cut 10 micra anteriorly and 15 micra farther back, and made about 1200 sections. About 850 sections were drawn at a magnification of 25 diameters with the aid of the Edinger projection apparatus. These have proved a very useful close series for annotation. A wax model made from these drawings, while somewhat disappointing, especially in the midregion, owing to the emptiness of the greatly expanded ventricles (the fragility of the specimen precluded aspiration) and consequent wrinkling and uneven spreading of the thinner portions of the walls, has been an invaluable aid to study. Linear reconstructions drawn to scale have also proved indispensable. An additional series of 31 Edinger drawings with a magnification 85 was made from the precommissural hippocampal region.

EXTERNAL ANATOMY

Dr. Herrick, in the first account ever given of the brain of *Caenolestes* (or of any caenolestid brain), emphasized its extreme simplicity, the enormous development of its rhinencephalon, the smoothness and limited extent of its neopallium, and called attention to the fact that it most closely resembles the brains of the lowly Australian polyprotodonts, *Notoryctes* and *Perameles*. To this it will be necessary to add the description of the median section of the whole brain and median surface of the hemisphere, as well as a discussion of the other surfaces in the light of knowledge of the internal structure of *Caenolestes*, with reference to changes of nomenclature, or in comparison with the brain of *Orolestes*. This will of course involve some unavoidable repetition.

The two types, *Caenolestes* and *Orolestes*, exhibit only minor differences in external anatomy. Although all three brains are supposed to have come from adult female specimens, they differ somewhat in size and proportions, as may be observed in the table of measurements. *Caenolestes* is the smallest of all, but it has the longest hemisphere. Its cerebellum is smaller and less plump than the other two, and exhibits an extra furrow and convolution—not a matter of great importance, as will be seen below under the description of the cerebellum. It is, however, to be remembered that the three brains represent two genera of caenolestids.

The *olfactory bulbs* are truly enormous, almost half the total length of the hemisphere at the medio-ventral angle. They are of the "sessile" type, with no visible olfactory peduncle in the intact brain. Following Livini (1908), we have called the deep constricting sulcus which marks the posterior border of the olfactory formation, the *fissura circularis* (*fs.circ.*, Figs. 6, 12, 15, 25-28; unlabelled, Fig. 1-5). The circular fissure is a compound structure whose component parts are homologous to portions of fissures described in the brains of other animals. Medially it represents the anterior portion of the fissura prima of His, above the antero-medial tip of the rhinal arcuate fissure (*fs.rh.acr.*, Figs. 12, 28); dorsally, after meeting the medial prolongation of the rhinal fissure (*fs.rh.a.*, Figs. 15, 25) at its medial end, it diverges from the latter to leave the interval for an exposed triangle of dorsal olfactory peduncle, and then meets the rhinal fissure again as the latter turns caudad on the lateral surface of the brain (Fig. 14); meets also the anterior end of the endorhinal fissure (*fs.erh.*, Fig. 13) just below, and then drops downward across the wide lateral olfactory tract (*tr.*

ol.l., Fig. 13) overlying the pyriform lobe, to continue as a deep constriction across the base of the brain between the olfactory bulb (*b.ol.* and the olfactory tubercle (*t.ol.*, Fig. 15), finally joining the anterior end of the medial portion of the rhinal arcuate fissure (Fig. 15), and thus completely defining the caudal border of the olfactory formation. Dorsally the olfactory bulb is overhung by the projecting frontal pole of the hemisphere, which conceals the exposed olfactory peduncle from view. The rest of the olfactory peduncle (see *anterior olfactory nucleus*, page 191) is thrust deep into the heart of the olfactory bulb, as recognized long ago in this type of brain by Elliott Smith and others. This condition is partly the result of the great caudal extension of the olfactory bulb itself, partly one of the numerous expressions of the very considerable fore-and-aft "telescoping" characteristic of this brain, which puts it in rather marked contrast to more elongated brains like that of the Virginia opossum (*Didelphis virginiana*), for example. *Didelphis marsupialis*, however, resembles the caenolestid brain in this respect (Beccari, 1910, Fig. 18).

The great hemispherical mass of the tuberculum olfactorium (*t.ol.*) is also delimited by an encircling sulcus, the *fissura rhinalis arcuata* (*fs.rh.arc.*, an adaptation of one of Retzius' names; see Herrick, 1924a). Medially the fissure is homologous to the caudal portion of the fissura prima of His. Its posterior more transverse portion has been called the fissura diagonalis by Beccari (1910), but his median continuation of it upward in the septum is not followed here (see tuberculum olfactorium, page 196). Laterally the rhinal arcuate fissure is usually taken to be homologous to the endorhinal fissure, but this is not strictly accurate, since the endorhinal fissure lies wholly within the pyriform lobe and therefore above the tuberculum, as has been noted in other forms by Smith, Johnson and others. The endorhinal fissure really marks the dorsal border of the massive part of the lateral olfactory tract, which is very wide rostrally, but rapidly diminishes caudally through the loss of fibers to the pyriform lobe and tuberculum olfactorium, so that in its caudo-ventral course it gradually approaches the rhinal arcuate fissure and finally meets it near the posterior limit of the tuberculum (Figs. 13, 15, 33-34). The greatly reduced massive part of the lateral olfactory tract lies at that level within the endorhinal fissure, where the latter fuses with the rhinal arcuate fissure and so comes to an end. In its caudo-ventral course the lateral olfactory tract leaves a thin film of fibers (*tr.ol.l.d.*, Figs. 25-33) covering the pyriform cortex almost to the height of the rhinal fissure and far toward the caudal pole of the pyriform lobe.

Another important fissure visible on the lateral surface of the hemisphere of these brains is the *rhinal fissure* (*fs.rh.*, to be seen in the majority of the figures), which separates the dorsal or neopallial convexity from the latero-ventral pyriform lobe. It traces an almost exactly horizontal course backward over the lateral surface of the brain at the very high level characteristic of the lowly marsupials *Notoryctes* (where it is internal only) and *Perameles* and the insectivore *Erinaceus*, forms whose neopallial development is the least extensive among mammals. Anteriorly the rhinal fissure is a continuation of the deep groove between the projecting frontal pole of the neopallium and the exposed dorsal part of the olfactory peduncle (*fissura rhinalis medialis* of some writers, but here called the *fissura rhinalis anterior*, *fs.rh.a.*, Figs. 12-15), which bends sharply on to the lateral surface of the hemisphere just above the rostral end of the endorhinal fissure and runs horizontally backward at this high level to round the caudal pole of the hemisphere, where it appears upon the median surface as the median rhinal fissure running obscurely forward into the subicular border of the hippocampus (*fs.rh.m.*, Figs. 6, 12, 42-43; *rh.*, Fig. 5, where it probably coincides only caudally with the diagonal groove shown—there has been no control with sections in *Orolestes* so far). Even where imperfectly developed externally the position of the *fissura rhinalis* is unmistakably discernible throughout, and internally it is in *Caenolestes* always clearly defined.

The dorsal convexity or *neopallium* contains only one other externally visible fissure. In *Caenolestes*, "about one-fifth of the distance from the frontal to the posterior pole of the hemisphere there is a distinct, though shallow, transverse sulcus which probably represents the sulcus orbitalis of Elliot Smith's descriptions (Herrick, 1921, *antea*, p. 158). Ventrally the orbital fissure is "obscurely confluent" with the rhinal fissure. In the smaller of the two brains of *Orolestes* this fissure seems to be absent as an external groove, but its position is apparently marked by blood vessels. In the reconstructions of *Caenolestes* also it does not show up, and a cursory examination of the sections has failed to reveal it. A more careful study might disclose its position.

The *pyriform lobe* comprises the larger part of the lateral and ventral aspects of the hemisphere. Its dorsal boundary is the rhinal fissure, which is at the same time the latero-ventral boundary of the neopallium. Antero-laterally the pyriform lobe contains the very sharp endorhinal fissure (as explained above, page 179) and below this fissure its antero-lateral boundary is the lateral portion of the rhinal

arcuate fissure (sharply incised in *Orolestes* but not depressed at all in *Caenolestes*), a fissure marking the *ventral* border of the massive portion of the lateral olfactory tract, as the boundary of the tuberculum, since the latter appears just ventral to this line.

The greatly widened caudal portion of the pyriform lobe is divided into two distinct parts by a very important fissure so faintly indicated externally that its existence was unsuspected before the brain was sectioned. This is the *fissura amygdaloidea* (1913, 1923) of Johnston (*fs.amg.*, Figs 2, 3, 6, 12, 13, 15, 36-43). In the sections it branches quite clearly from the rhinal arcuate fissure a short distance in front of the caudal pole of the tuberculum (between Figs. 35 and 36), and very soon disappears as a recognizable fissure. Internally it is quite distinct throughout its course, marking the quasi-horizontal line of junction between the sharply defined ventral edge of the pyriform cortex and the lateral border of the amygdaloid complex which occupies practically the entire base of the hemisphere behind the tuberculum. The rather pronounced sculpturing of this region is due to the state of development of the different amygdaloid nuclei, which have been so thoroughly worked out in the Virginia opossum by Johnston (1923). These will be more fully described further on, in connection with the internal anatomy of this brain (page 201).

In *Caenolestes* a very salient caudo-ventro-lateral prominence corresponds to the "natiform eminence" of Elliot Smith (1895b) in *Notoryctes* (*em.nat.*, Figs. 37-41; unlabelled, Fig. 2, pl. XXI, antea, Herrick; cf. *Orolestes*, figure 3 here, which shows it much less distinctly). It is due both to an actual thickening of the lateral wall of the hemisphere in this region and, in greater degree perhaps, to the lateral cupping of the hemisphere to accommodate the wide midbrain. Elliot Smith thinks that the temporal bending of the hemisphere has been a main factor in the formation of this eminence. This is much more strongly suggested in *Notoryctes* than in *Caenolestes* (see page 213).

The median section of the brain of *Caenolestes* is known only in wax and linear reconstructions, but the smaller of the two brains of *Orolestes* was divided sagittally, and figures 4 and 5 show respectively the left median section of the whole brain and the median surface of the right hemisphere. Figure 6 is a linear reconstruction of the left hemisphere (reversed) of *Caenolestes*.

The third ventricle and external medial surface of the brain of *Orolestes* were so filled or coated with a hard coagulum of the same color as the tissue and so tightly adherent to it that it was not found

possible to remove it without some laceration of the tissue. Thus the edges of the massa intermedia (*m.i.*, Fig. 4) were somewhat frayed, the aqueduct was probably widened, and the epiphyseal attachments and choroid roof of the third ventricle were torn. But these defects, regrettable as they are, do not greatly impair the value of the section for present purposes. At least the sacrifice of the other brain with the prospect of no better success did not in the circumstances seem to be justified.

The large anterior or ventral commissure of *Orolestes* (*v.*, Figs. 4 and 5) is slightly oval and considerably darker than the rest of the surface, standing out sharply. That of *Caenolestes* (*v.*, Fig. 6) is similar in size, shape and position. The dorsal commissure (*d.*) of *Orolestes* appears as a moderately darker and distinctly bilaminar mass occupying the dorsal and posterior borders of this portion of the lamina terminalis (lamina supraneuroporica of Johnston), enclosing a small whitish triangular space which gives the impression of being without transverse fibers. The dorsal commissure of *Orolestes* (counting it only as the darkened bilaminar mass) thus exhibits macroscopically the typical marsupial form, while that of *Caenolestes* (*d.*, Fig. 6) as reconstructed from sections, shows only the merest hint of bilaminarity in the slightly reentering anterior angle and the dorso-caudal prolongation ("splenium"), suggesting an intermediate type between the solid rounded type of the monotremes and the bilaminar type of the marsupials. The reconstruction of *Caenolestes* was made from the exact midline of the sections, the outline enclosing only the area through which commissural fibers were coursing. These fibers are much less dense in the ventral than in the dorsal region of the commissure, owing to the intermingling of many cells (bed or nucleus of the dorsal commissure) with the fibers in its ventral region.

In *Orolestes* the third ventricle (Fig. 4), as seen from the median surface, has the form of a tall parallelogram tipped slightly backwards and downwards. Its walls are formed anteriorly by the lamina terminalis, whose upper half is greatly thickened by the two commissures, while the lower half remains very thin; posteriorly, by the almost vertical mammillary body and tuberculum posterius in line with the rostral opening of the aqueduct (*aq.*) and the thick anterior end of the tectum mesencephali (*tect.*); dorsally by the choroid roof and the epiphysis; ventrally by the thin floor plate containing the small chiasma (*ch.*) rostrally, and the infundibular lumen more caudally; laterally, by the median surfaces of the thalami. The upper half of the third ventricle is almost filled by the enormous oval massa intermedia, which

so nearly approaches the neighboring walls as to leave on three sides of it only two very narrow passages leading into the aqueduct. The lower half of the ventricle is quite open, and the thalamic lateral wall displays a diagonal sulcus running backward and downward from the middle of the lower border of the massa intermedia to the posterior wall of the ventricle just above the floor. The ventricle is also deepened laterally just above the floor in two places—the preoptic recess and the mammillary recess. The interventricular foramen opens into the narrow canal between the dorsal commissure and the massa intermedia.

A very sharp midbrain flexure results in the formation of a deep and narrow vertical cleft across the base of the brain which folds a section of it between the pons and the mammillary region together like the pages of a book. The tectum mesencephali (*tect.*) is entirely concealed from above by overlying structures, the rostral three-quarters by the cerebral hemispheres and the caudal quarter by the anterior tip of the median lobe of the cerebellum. (Figure 4 shows a gap between the hemisphere and the cerebellum, due to a mass of coagulum afterwards cleared away; the relations in *Caenolestes* and *Orolestes* are the same.) "This is in contrast to the usual marsupial arrangement, for the corpora quadrigemina are in most cases well exposed dorsally. (*Petaurus* is another exception; see Elliott Smith, 1895, p. 188)" (Herrick, 1921, *antea*, page 158). The condition in *Orolestes* and *Caenolestes* is apparently one of the many expressions of the fore-and-aft compression already mentioned. Partly in response to the crumpling of the brain base as described above, the tectum displays a marked caudal prolongation, emphasized by its recurved keel, as indicated by the forward point of attachment of the anterior medullary velum (*v.m. a.*, Fig. 10). Laterally the posterior colliculus is even more caudally extended (*col. p.*, Figs. 7, 8, 10). The cerebellum, which will be described below, fits snugly into and around these structures, behind an almost perpendicular anterior medullary velum, and below, behind and above the tectum.

The median surface of the cerebral hemisphere of *Orolestes* (Fig. 5) and of *Caenolestes* (Fig. 6) exhibits a flattened surface anteriorly, in close apposition with its fellow, and a deeply concave postero-median surface, hollowed out to accommodate the bulky midbrain. It is traversed by four prominent arcuate fissures, concentrically arranged and different in extent and curvature. These are, in order from without in, the hippocampal, fimbrio-dentate, fimbrio-alvear (to coin a term), and choroid fissures.

The outermost or *hippocampal fissure* (*fs.hip.*, Figs. 5 and 6) apparently describes almost three-quarters of a circle in *Orolestes* (Fig. 5), stretching from the fissura circularis (*fs.circ.*) upward, backward, downward and forward nearly to the posterior end of the tuberculum olfactorium. The same fissure in the reconstruction of *Caenolestes* (*fs.hip.*, Fig. 6) falls short of this at both ends. It does not reach the fissura circularis rostrally and it ends more briefly caudally near the medial prolongation of the fissura amygdaloidea, as the latter goes forward towards the posterior end of the choroid fissure. Perhaps also in *Orolestes* what appears to be the strongly recurved temporal end of the hippocampal fissure is really the fissura amygdaloidea medialis. Indeed this interpretation is suggested by the sudden pronounced shallowing and narrowing of the fissure at the exact point where a slight diagonal fissure runs into it from behind. The posterior end of the diagonal fissure coincides with the apparent level of the caudal end of the amygdaloid fissure on the lateral surface, which on the figured lateral view of the left hemisphere (Fig. 3) is much less clearly indicated than on the right (Fig. 5). The sections of this particular right hemisphere will, when made, clear up this point, which now rests partly on the external configuration of the hemisphere of *Orolestes* and partly on the internal configuration of that of *Caenolestes*.

The second fissure, the *fimbrio-dentate fissure* (*fs.fim.d.*) extends also in *Orolestes* (Fig. 5) to the fissura circularis, just below the fissura hippocampi. In *Caenolestes*, however, it drops sharply downward away from the hippocampus entirely, and ends briefly just in front of the dorsal commissure (Fig. 6). This portion of the fissure seems to be purely a response to the pressure of a blood vessel which is lodged in the canal formed by the corresponding fissures of the two hemispheres (Figs. 6, 17b). The postcommissural and main portion of the fimbrio-dentate fissure separates the gyrus dentatus (*gy.dent.*) not from the massive fimbria, but from the extraventricular ammon's horn or inverted hippocampus, with its thin coating of subpial alveus fibers, and it is therefore really an *alveo-dentate* fissure in this region. Posteriorly it is both less extensive and less recurved than the hippocampal fissure.

The third fissure seems to have no name, but it is nevertheless quite distinct as the line of demarcation between the massive fimbria and the alveus-coated extraventricular hippocampus (ammon's horn), defining its rolled ventral limit. It might be designated as the *fimbrio-alvear fissure* (*fs.fim.al.*, Figs. 6, 17a; unlabelled, Figs. 5, 35-38). It begins rostrally just behind the dorsal commissure as an offshoot of

the fimbrio-dentate fissure, dropping rapidly away from the fimbrio-dentate fissure as the extraventricular hippocampus widens, and it runs out caudally as the massive fimbria tapers down to nothing at the posterior end of the choroid fissure (just in front of figure 39).

The innermost or *choroid fissure* (unlabelled, Fig. 5; *fs.ch.*, Figs. 6, 17a, 35-39; Fig. 39 is really just behind the caudal end of the choroid fissure) extends from the ventral tip of the dorsal commissure backwards and downwards in a smooth curve outlining the ventral border of the fimbria and the thalamo-hemispheric junction. In *Caenolestes*, as noted above, the internal medial prolongation of the fissura amygdaloidea can be quite clearly traced forward to the neighborhood of the caudal end of the choroid fissure.

Other features of the medial surface of the hemisphere are the *tuberculum olfactorium* in profile behind the olfactory bulb, followed immediately in *Orolestes* by the small ventro-median eminence (Fig. 5, unlabelled) which is probably due to the nucleus of the lateral olfactory tract (*nuc.tr.ol.l.*, Figs. 36-38). This small tubercle appears in *Notoryctes* (Elliot Smith, 1895b) and in a number of other marsupials and lowly eutherian mammals, but is apparently absent or not very prominent in *Caenolestes*, although the nucleus, as will appear below, is well developed. It is, however, partly covered by the *tuberculum olfactorium*, and this masks it to some extent. The *amygdaloid complex* (*amg.*, Fig. 6; unlabelled, Fig. 5) extends upward as high as the position of the medial portion of the amygdaloid fissure (*fs.amg.m.*) already described, and the posterior pyriform cortex (*cx.pir.p.*) curves around the caudal pole of the hemisphere to form the more temporal subicular border between the medial extensions of the rhinal and amygdaloid fissures, in the familiar mammalian pattern. From the postero-medial extension of the rhinal fissure (*fs.rh.m.*) forward to the antero-medial extension of the same fissure (*fs.rh.a.*) the space above the hippocampal fissure is occupied by neopallium. In front of the anterior rhinal fissure the hippocampal cortex is in cellular continuity with the dorso-medial portion of the anterior olfactory nucleus (*nuc. ol. ant.d.*, Fig. 26), the peduncular or postbulbar gray matter which laterally merges without interruption into the pyriform cortex (see page 192). Save for these rostral and caudal junctions the lateral and medial olfactory cortices are split apart dorsally by the wedge-like neopallial cap of the hemisphere.

The entire length of the fissura prima of His appears upon the median surface of the hemisphere as the visible portions of the fissura circularis rostrally and fissura rhinalis arcuata caudally (see page

178). Between it and the lamina terminalis lies the precommissural area (paraterminal body) of Elliot Smith, the parolfactory region of Johnston, the septum of most neurologists (*a. prcom.*, Fig. 6; unlabelled, Fig. 5). The nucleus parolfactorius medialis, the nucleus of the diagonal band of Broca, the precommissural fornix, olfacto-hippocampal fibers, and the medial forebrain bundle (fasciculus medialis telencephali), are the more superficial structures lying above the posterior part of the fissura prima (medial part of the fissura rhinalis arcuata). The sharply grooved dorso-medial margin of the hemisphere is the imprint of the longitudinal sinus, which is seen in position in figure 4.

The enormously expanded lateral ventricle of *Caenolestes* (*v.l.*, Figs. 28-42) is probably not pathological, but the result of defective fixation. As a cursory inspection of the more caudal sections will show, a recurved temporal horn has not even begun to form. The posterior horn, present in the Virginia opossum, is also absent in *Caenolestes*. In the lower rostral wall of the ventricle a flaring opening leads into a narrow crooked canal, which in turn expands into a widened terminal sac. These constitute the olfactory ventricle (*v.ol.*, Figs. 12-15, 17, 23-27).

Cerebellum.—(Figs. 1, 3, 4, 7-10, 10a and 10b, 41-44; also Herrick, 1921, *antea*, pl. XXI.) The cerebellum of *Caenolestes* and *Orolestes* (the two forms are practically identical in general structure) very neatly fills the gap between those of *Notoryctes* and *Perameles*, the two simplest mammalian cerebella hitherto described. (Elliot Smith, 1902b, 1903c, d, e). The cerebella of the insectivores *Macroscelides* and *Erinaceus* (also described by Elliott Smith, 1902c, 1903c) probably occupy the fourth and fifth places in the series. The cerebellum of the Virginia opossum is considerably more complex than any of those mentioned.

Of the three fundamental cerebellar lobes separated by the fissura prima (*fs.pr.*) and fissura secunda (*fs.s.*), the median and posterior lobes differ little in the first four forms named above. But the anterior lobe grades very clearly from the simple unfissured lobe of *Notoryctes* (10a), through *Caenolestes* (Figs. 4, 7-10) or *Orolestes*, with two lobules separated by the deep fissura preculminata (*fs.prcul.*), and a third well developed lobulus, the lingula (*lg.*), which is not separated from the rest by a fissure, to *Perameles* (Fig. 10b), with an anterior lobe crossed in the midline by four fissures of varying depth. The anterior lobe and the fissura prima are entirely concealed in the intact

brain. The visible portion of the cerebellum is formed by the exposed parts of the median and posterior lobes. It is crossed in the midline by a ventrally concave fissure, the fissura secunda (*fs.s.*), separating a small postero-medial convolution, the uvula (*uv.*) belonging to the posterior lobe, from the larger median lobe above. In *Caenolestes* but not in *Orolestes*, the secondary fissura suprapyramidalis (*fs.spyr.*), shallow but sharp, divides the median lobe into a large suprapyramidal region (*p.spyr.*) and a slender curved ventral convolution, the pyramis (*pyr.*), concentric with the uvula, which fills the ventral concavity of the fissura secunda. The other lobule of the posterior lobe, the nodulus (*nod.*), lies entirely hidden on the ventral surface of the cerebellum, being separated from the uvula by the concealed fissura postnodularis (*fs.pnod.*). The cerebellar ventricle (*v.cb.*) is seen between the nodulus and the lingula.

The large mushroom-shaped pedunculate lateral masses projecting beyond the lateral lobes (these indicated only by a very slight partial constriction) are the paraflocculi (*pfloc.*), the dorsal components of the floccular lobes. The ventral components are the tiny flaplike flocculi, concealed from view, being plastered against the medulla beneath the paraflocculi, from which they are separated by the fissura floccularis (*fs.floc.*). The fissura parafloccularis (*fs.pfloc.*), visible laterally, separates the paraflocculus from the median lobe. It cuts down to the floccular peduncle between these structures.

All the cerebellum save the floccular lobe and the lateral lobes (area pterioidea, *a.pt.*, in part) corresponds probably to the vermis of higher cerebella. The lateral lobes similarly are probably homologous with the cerebellar hemispheres.

The deep nuclei (*deep nuc.*) form a pair of large oval masses almost meeting in the midline, with a large area of their ventral surface exposed in the roof of the fourth ventricle. They exhibit the mammalian characteristic of complete separation from Deiters' nucleus, as van Höevell (1916) found in some other marsupials. There are only slight indications of differentiation into separate nuclei (dentate and roof nuclei).

Since the cerebellum of *Caenolestes* and *Orolestes* will form part of the next report on these brains, it need not be further discussed at this time. The intra- and extracerebellar relations will be partially clarified by the diagrammatic reconstructions, figures 7-10, together with the sections, figures 41-44. The nomenclature follows the usage of Elliot Smith.

Measurements.—The dimensions of these three brains as measured on the alcoholic specimens are as follows:

	<i>C. obs.*</i> 18507 F.M.	<i>O. inca</i> 194948 194921 U.S.N.M. U.S.N.M.	
1. Total length, tip of olfactory bulb to first spinal nerve	14.1 mm.	16.5 mm.	19.1 mm.
2. Length, tip of olfactory bulb to end of cerebral hemisphere	2.6	3.0	3.6
3. Length of cerebral hemisphere	10.0	9.3	8.8
4. Length of cerebellum on longitudinal axis of brain in median plane	3.0	5.0	6.2
5. Greatest width of olfactory bulbs	7.6	7.8	7.9
6. Greatest width of both cerebral hemispheres	11.8	11.8	12.6
7. Total width of cerebellum and floccular lobes	11.0	11.7	
8. Width of cerebellum exclusive of floccular lobes	8.8	11.0	11.0
9. Maximum vertical height of cerebral hemispheres	...	7.9	8.9
10. Distance of orbital fissure behind rostral tip of hemisphere	(2.0)*	2.3*	2.4
11. Distance of rhinal fissure below vertex at orbital fissure		2.7*	4.2
12. Length, head and body	107.0	(89.0)	(102.0)
13. Length, tail	118.0	115.0	116.0
14. Length, total	225.0	204.0	218.0
15. Weight of brain, immediately out of 80% alcohol	...	760.9 mg.	961.0 mg.

* *C. obscurus*, measurements 1-11, from Herrick (1921); 12-14, from Dr. Osgood.

* Position of orbital fissure in *C. obscurus* is "about one-fifth of the distance backward from the frontal to the posterior pole of the hemisphere".

* Orbital fissure not very clear in *O. inca*, No. 194948.

O. inca No. 194921 had probably all of cervical cord attached, but lost parafloccular lobes on removal from the skull.

O. inca No. 194948 was broken off just behind cerebellum, in removing it from the skull.

INTERNAL ANATOMY

The twenty-two simplified cross sections from the hemisphere of *Caenolestes obscurus* (Figs. 23-44) and the linear reconstructions of the hemisphere or of parts of it (Figs. 12-15, 17a and b) may perhaps, with the aid of the figures and descriptions of the external surfaces, render a brief and incomplete account of the internal anatomy comprehensible. The following table of critical levels and their section numbers may also assist in the orientation of internal structures.

Sec. 1—Anterior tip of the olfactory bulb.

Sec. 225—Anterior tip of the frontal lobe of the neopallium (Fig. 23).

Sec. 400—Caudal (medio-ventral) limit of the olfactory bulb, adjoining the tuberculum olfactorium (Fig. 25).

Sec. 710—Caudal limit of the tuberculum, adjoining the amygdaloid complex (Fig. 36).

Sec. 935—Posterior tip of the cerebral hemisphere (Fig. 44).

PRIMARY OLFACTORY AREA

The description of the histological pattern of the *bulbar formation*, both ordinary and accessory, may be dispensed with here, in view of its similarity to that of the Virginia opossum, already described in detail by McCotter (1912). Contrary to the condition in the opossum, however, the horizontal diameter of the olfactory bulb in *Caenolestes* is greater than the vertical diameter. The usual medial displacement of the olfactory ventricle is emphasized here by the considerable dorso-ventral flattening of the olfactory bulb. This displacement may perhaps be a condition of interest in connection with the disposition of the olfactory tract fibers.

The well developed *accessory olfactory bulb* (*b.ol.ac.*, Figs. 12, 14, 23-24) is embedded in the postero-dorsal olfactory formation beneath the overhanging frontal pole of the neopallium. It is outlined by a slight fissure. Its peripheral nerve, the *vomeronasal nerve* (*n.vn.*, Figs. 12, 14, 23-24) is quite clear as it curves up over the medio-dorsal angle of the olfactory bulb and spreads out to cover the surface of the accessory bulb. The *secondary tract* of the accessory bulb passes laterally and superficially and helps to make up the pars dorsalis of the lateral olfactory tract (*tr.ol.l.d.*, Figs. 23 ff.) The olfactory tracts, which, as Cajal pointed out (1911), arise indiscriminately from all parts of the olfactory formation and are without specificity (save for the fibers from the accessory bulb), first condense in the center of the bulb some distance in front of the olfactory ventricle (*tr.ol.*, Figs. 12-15). Since they so quickly begin to distribute to the secondary olfactory areas and even in part (intermediate olfactory tract) to receive fibers from them, their description will be continued below under that head.

SECONDARY OLFACTORY AREAS

The olfactory fibers spin a whorl around the olfactory ventricle, much thicker on the lateral side than elsewhere. A second independent half-whorl forms within the first, between its thick lateral portion and the ventricle. This mass of fibers, composed of secondary (direct) and tertiary olfactory fibers, soon rounds up in the lateral wall of the ventricle as the intermediate olfactory tract (*tr.ol.i.*, Figs. 23-28), largely forming the rostral limb of the anterior commissure (*c.a.*, Figs. 24-28). Meanwhile, between it and the lateral part of the first whorl (the massive portion of the lateral olfactory tract, *tr.ol.l.*) appear the most anterior cells of the peduncular grey or anterior olfactory nucleus, pars lateralis (*nuc.ol.ant.l.*, Figs. 12-15, 17, 23-25). This growing

cell mass rapidly increases the widening interval between the lateral part of the big whorl externally and the half-whorl internally, and immediately begins to contribute tertiary olfactory fibers to the inner half-whorl, the intermediate or commissural olfactory tract, which also continues to receive secondary olfactory fibers from all parts of the olfactory formation. Still farther back the tuberculum olfactorium (*t.ol.*, Figs. 13, 28) cuts the outer whorl into two parts, the *lateral olfactory tract*, much the largest of all the olfactory tracts, and the slender *medial olfactory tract* (*tr.ol.m.*, Fig. 28). The *intermediate tract*, which retains its subependymal position back to the point where the head of the caudate nucleus (*nuc. caud.*, Fig. 28) overlies it, soon contains perhaps more tertiary than secondary fibers, and is therefore labelled only anterior commissure (*c.a.*) from figure 24 back.

The *lateral olfactory tract*, the most widely distributed of the three, becomes superficial (*tr.ol.l.*, Figs. 3, 12, 25) behind the fissura circularis, forming the medullated external fiber layer of the lateral surface of the hemisphere over most of its extent below the rhinal fissure. It may be divided into several parts: anteriorly, two peduncles, a sub-ventricular ventral peduncle (*tr.ol.l.p.v.*, Figs. 24-27) a part of which caudally becomes independent and is then called the medial olfactory tract (*tr.ol.m.*, Fig. 28) as noted above, and a supraventricular dorsal peduncle (*tr.ol.l.p.d.*, Figs. 23-24) with two roots, a ventral root, below the accessory olfactory bulb, and a dorsal superficial root which is continued caudally as the pars dorsalis of the lateral olfactory tract (*tr.ol.l.d.*, Figs. 25 ff.); a clubshaped massive portion (*tr.ol.l.*, Figs. 23-35) filling the interval between the endorhinal and the rhinal arcuate fissures, wide anteriorly but decreasing to the vanishing point as these fissures meet (*fs.erh.*, *fs.rh.arc.*, Figs. 13, 33); posteriorly a pars ventralis (*tr.ol.l.v.*, Figs. 29 ff.), distributing to the tuberculum, nucleus of the lateral olfactory tract (*nuc.tr.ol.l.*, Figs. 12, 36-38) and other amygdaloid nuclei, and perhaps also to the diagonal band nucleus (*nuc.d.b.*, Figs. 32-33). The small *medial olfactory tract* (*tr.ol.m.*, Fig. 28) can be seen turning sharply downward into the plexiform layer of the tuberculum at its medial border, where some observers (*e.g.*, Livini, 1908) recognize a terminal nucleus of the median olfactory tract in a large median rolled portion of the tubercular formation. Such a structure is present here (*nuc.ol.m.*, Fig. 31). Beccari (1910) inclines to doubt the distribution of secondary olfactory fibers to any extent to the tuberculum, but such evidence as these sections give, while not at all conclusive, seems to point to it.

Anterior olfactory nucleus.—The less differentiated postbulbar gray matter of the rhinencephalon has been termed by Herrick (1910, pp. 191-2, Figs. 9 and 10) the anterior olfactory nucleus. In *Caenolestes* it is almost entirely intrabulbar and in direct continuity with the rest of the more ventral gray subjacent to the lateral olfactory tract. Rostrally therefore it completely surrounds the olfactory ventricle. It corresponds essentially to the peduncular gray of Cajal, and is not included in his pyriform lobe. It is here excluded from the pyriform lobe, though earlier, following Johnston's definition of the pyriform lobe as all the gray underlying the lateral olfactory tract, I counted it a part of the pyriform lobe (Obenchain, 1923). Since its lateral border is farthest advanced rostrally, this periventricular rhinencephalic ring appears incomplete in the more rostral sections (Figs. 23-26), and since the pyriform cortex is also advanced rostrally beyond some portions of the anterior olfactory nucleus, the whole of the latter is never seen in any transverse section. The anterior olfactory nucleus, on both topographical and histological grounds, may be subdivided into several parts, all in cellular continuity rostrally. I have, as far as seemed advisable, conformed to the terminology applied to the different parts of the anterior olfactory nucleus in the Virginia opossum (Herrick, 1924a). Divergences will be noted as they occur.

The *lateral part* (*nuc.ol.ant.l.*, Figs. 12-15, 17, 23-25), rostrally most advanced, passes caudally without interruption into true pyriform cortex, and therefore has no really definite posterior limit. The *pars lateralis* is extended medially above the olfactory ventricle as the *pars dorsalis* (*nuc.ol.ant.d.*, Figs. 12-15, 17, 25-26); it corresponds to Cajal's superior peduncular nucleus. Just in front of the antero-median extension of the rhinal fissure (*fs.rh.a.*) it comes to the surface, as noted above, and just behind it fuses with the cell mass of the overlying neopallial frontal pole. More caudally and below the ventricle the anterior olfactory nucleus is also prolonged medially by another hooklike extension, the *pars latero-ventralis* (*nuc.ol.ant.l.v.*, Figs. 12, 15, 17, 26), and beyond that the *pars posterior* (*nuc.ol.ant.p.*, Figs. 12, 15, 17, 27-29), which fills the wedgelike interval in front of the fusion of the head of the caudate nucleus and the tuberculum olfactorium. The *pars posterior* extends further back than other portion of the anterior olfactory nucleus, but it is neither in continuity with the caudate nucleus—forming the rostral portion of its head, as in the turtle, *Cistudo carolina* (Johnston, 1915)—nor with the deeper layer of the tuberculum, as in the alligator (Crosby, 1917). It stops bluntly just rostral to the fusion of the caudate head and the tuberculum. This part

of the anterior olfactory nucleus never shows any appreciable degree of differentiation, probably owing to the too purely olfactory character of its connections. The latero-ventral and posterior parts of the anterior olfactory nucleus as here described do not exactly coincide with the divisions in the Virginia opossum. More caudally the dorsal and posterior parts meet medially, thus completing the anterior periventricular rhinencephalic ring by means of a *pars medialis* (*nuc.ol.ant.m.*, Figs. 12, 15, 17, 26).

At the dorso-medio-rostral border of this ring there is a small superficial condensation of cells. This is the anterior tip of the *hippocampal formation* (*hip.a.*, Figs. 12, 17; *cx.hip.a.*, Fig. 26). It corresponds to a similar dorso-lateral *pyriform* condensation (*cx.pir.a.*, Fig. 25), which, although it appears in more rostral sections, does not quite reach the rostral border of the ring. The lateral and medial olfactory cortices (pyriform and hippocampal) are thus only *indirectly* continuous rostrally by a double bond (supra- and infraventricular) through the agency of the anterior olfactory nucleus, while, as we have seen, they are directly continuous caudally. There are thus two periventricular rhinencephalic rings, a smaller transverse and a larger horizontal one, which are partially fused anteriorly.

The medial part of the anterior olfactory nucleus soon passes obscurely backward into the lateral parolfactory nucleus (*nuc.pol.l.*, at a level between figures 26 and 27). The latero-ventral part, which is transitional between the lateral and posterior parts, continues backward into the ventral pyriform cortex underlying the massive portion of the lateral olfactory tract along the lateral border of the tuberculum (*pars ventralis*, *lobus piriformis*, Gray, 1924; *ex.pir.v.*, Figs. 27-33), and still more caudally perhaps into the diffuse region of the anterior perforated space (Johnston, 1923) (*l.perf.a.*, Figs. 35-38), along the anterior portion of the amygdaloid fissure (*fs.amg.*, Figs. 12, 15, 36-38).

One other—and the most interesting—part of the anterior olfactory nucleus remains to be described, the *pars externa* (*nuc.ol.ant.ex.*, Figs. 13, 15, 24-26). It is apparently the result of the doubling of the intrabulbar portion of the lateral part of the anterior olfactory nucleus, which is the only portion to exhibit this phenomenon. The external nucleus appears rostrally as a vertical plate of smaller cells between the *pars lateralis* and the massive lateral olfactory tract. It is separated from the subjacent *pars lateralis* by a wide plexiform layer narrowing ventrally to suggest cellular continuity between the two nuclei. Rostrally a very narrow but definite external plexiform layer separates it

from the fibers of the lateral olfactory tract. Caudally the external nucleus, or external part of the lateral part of the anterior olfactory nucleus (to give it its full designation) slides downward along the internal surface of the lateral olfactory tract, gradually diminishing and rounding up into a small mass of cells which, as it slips medialward around the inside of the "elbow" of the lateral tract, becomes counter-sunk in a space due to the clearing away of some of the blue tract fibers. It is here opposite to the pars latero-ventralis, but its main bulk lies opposite the pars lateralis—there is no definite boundary between these two parts. The general form of the external nucleus is that of a long and slender pennant (Fig. 13) whose caudo-ventrally extended tail curves far medialward to end almost beneath the olfactory ventricle (Fig. 15). The external nucleus of *Caenolestes* is thus entirely intrabulbar, so that it faces the granular olfactory formation across the olfactory tract. Dr. Herrick finds that the Virginia opossum also possesses an essentially similar nucleus, save that it lacks an external plexiform layer, and is caudally unevenly swallow-tailed. In *Caenolestes* a small dorsal group of cells tends to be separated from the rest by the fibers of the ventral root of the dorsal peduncle of the lateral olfactory tract. Röthig (1910, Fig. 1) figures but does not name a small vertical cell plate in *Didelphis marsupialis*, which is plastered against the inner aspect of the dorsal thin part of the lateral olfactory tract *behind* the bulbar formation, a dorsal shift which would bring it nearer to the remaining bulbar formation at this level. The entire width of the plexiform layer separates it from the underlying pyriform cortex. As this is the most anterior section figured by Röthig, the more rostral extent and relations of this cell plate are unknown to me. Livini (1908, Fig. 2) shows in *Hypsiprymnus rufescens* a doubling of the intrabulbar portion of his anterior pyriform lobe (which corresponds to the lateral part of the anterior olfactory nucleus as described here) into two nearly equal vertical plates of cells only narrowly separated, the outer of which seems to be composed of somewhat smaller cells. In a Nissl series of a white rat brain I find the anterior olfactory nucleus also doubling laterally in the olfactory peduncle, in the gap between the dorsal and ventral edges of the bulbar formation, which in cross section has just broken in two—in other words, just behind the fissura circularis. The external portion is a more condensed vertical plate of cells, immediately beneath the massive lateral olfactory tract, bridging the gap in the olfactory formation and therefore in contact with it at its dorsal and ventral margins. More caudally, with the widening of the gap between the receding edges of the olfactory

formation, the external nucleus separates into two parts, dorsal and ventral, which diminish as they separate more and more, and finally end as tiny cell masses at the upper and lower borders of the bulbar formation near its caudal limits. Its general form is that of an evenly swallow-tailed pennant. The Winkler-Potter rabbit and cat atlases (1911, 1914) give no figures showing such a nucleus. Cajal (1911) neither mentions nor figures it, a circumstance which is perhaps to be ascribed to its failure to impregnate in his rich collection of Golgi series of the brain of the mouse. His only figure showing the lateral peduncular gray (1911, II, Fig. 431, p. 696, mouse, 5 days old, after Calleja) is apparently a horizontal section, which may miss the level of the external nucleus, if there be such a nucleus in the mouse. I have so far found no reference in the literature to this curious cell mass beyond the Röthig and Livini figures cited above, in which it was unlabelled.

The interpretation of this nucleus offers apparently no great difficulty. It is, probably, some sort of reenforcing or stepping-up device for olfactory stimuli, an *accessory olfactory nucleus*, discharging its afferent fibers into the subjacent lateral part of the anterior olfactory nucleus. It is a regulatory response probably provoked in part by the antero-posterior compression of this highly macrosomatic type of brain at the point where the neurobiotactic attraction of accumulated secondary fibers is strongest. Does this nucleus represent the morphological anterior end of the anterior olfactory nucleus, detached and carried back by the more anterior collaterals of the lateral olfactory tract, either actually or by being held fast by them during the progress of the compression which telescopes bulb and anterior end of the extrabulbar rhinencephalon? Or is it delaminated *in situ* from the subjacent cell mass, being composed of the cells which possess no basilar dendrites and are therefore unable to resist the unopposed neurobiotactic influence of the overlying fibers? It is not inconceivable that both delamination and dislocation may have operated in its formation. Although Cajal neither pictures nor describes the external nucleus, and gives but one copied figure of the lateral peduncular formation in connection with which it arises, his figure and description of the region immediately *behind* its due position furnish apparently unmistakable clues, in the light of Kappers' concept of neurobiotaxis, of the mode and causes of its formation further forward (1911, II., Fig. 433, p. 680, rabbit, aged 25 days). In the cortex of his "frontal lobe" (anterior area of pyriform cortex, Gray, 1924) the third layer (outer cell layer, superficial polymorphs or medium pyramids) contains cells without basilar den-

drites, all their dendrites being directed outward toward the fibrillar layer (lateral olfactory tract). These cells form, according to Cajal, a rather precise wavy band, not however separated off from the cells below, and they are extremely variable in form. The deeper pyramidal cells and the still deeper polymorphic cells are provided with both ascending and descending dendrites, and are therefore doubly anchored and not subject to extreme outward displacement by unbalanced olfactory stimuli of the lateral olfactory tract, as are the more superficial cells without descending dendrites. The balancing attraction resides, probably, in the heterolateral secondary and perhaps in the tertiary fibers in the rostral limb of the anterior commissure, which lies beneath the lateral portion of the anterior olfactory nucleus. Whether any other fibers reach it from this direction, as, for example, homolateral secondary fibers of the intermediate olfactory tract, I do not know. At any rate this external part of the anterior olfactory nucleus is slung like a hammock between two opposing neurobiotactic forces, and the doubling or splitting of the nucleus in this region expresses the resolution of the situation. The deep and main portion of the nucleus in this region lies in *Caenolestes* closer to the anterior commissure than to the lateral olfactory tract. Cajal (1911, II., p. 678) says that the cortex of the olfactory peduncle and of the "frontal lobe" (which lies next behind it) are essentially the same in structure. We should indeed expect to find the lateral peduncular gray less differentiated, with perhaps shorter axons and fewer descending dendrites, and with only a slight tendency toward cortical lamination. If the situation is as sketched we have a clear and exquisite illustration of the two-fold activity of neurobiotaxis at work. Since the lateral olfactory tract is always in macrosomatic animals an exceedingly heavy mass of fibers it is not improbable that the external olfactory nucleus is well developed in at least all those forms whose olfactory peduncle (anterior olfactory nucleus) is jammed forward and largely enclosed within the bulbar formation, as in the forms mentioned here. We should expect to find vestigial traces of it widespread among mammals in general, and it has most likely already found its way into the literature in some form or other.

Tuberculum olfactorium.—The enormous tuberculum olfactorium begins in these sections latero-ventrally (*t.ol. (i.C.)*, Figs. 2-5, 12, 13, 15, 28-36), instead of medio-ventrally as in the Virginia opossum. It rapidly expands medialward to occupy the entire width of the base of the brain, and caudally to a point beyond the middle of the hemisphere (section 710 in a hemisphere numbering 935 sections). Antero-medially it turns up on the medial surface for a short distance, where it is de-

limited by the medial portion of the fissura rhinalis arcuata (*fs.rh.arc.*), which does not as here described quite agree with Beccari's account (1910, Fig. 18); he extends the anterior and posterior parts of the encircling fissure of the tuberculum upward in the medial wall to meet at the ventral border of the hippocampal formation; but I have thought it simpler (and more in keeping with corresponding conditions of the lateral wall) to carry it across the base of Beccari's median triangle, in a fissure which is present and which delimits the *cellular* formation characteristic of this region, just as the arcuate fissure does laterally. In the more rostral sections it will be seen that the median part of the rhinal arcuate fissure is much shallower than the very sharp one lying immediately below it within the tuberculum. The two fissures define a rather prominent rolled portion of the tuberculum which is probably to be identified with the *nucleus of the medial olfactory tract* of Livini (1908) and others. Above the rhinal arcuate fissure the characteristic formation of the tuberculum falls away from the surface and runs up along the median border of the nucleus accumbens (*nuc. ac.*, Fig. 30), almost if not quite to the ventricular ependyma beneath the more rostral portion of the body of the anterior commissure, where the latter breaks across the ventricle to reach the septum. It thus intervenes between the sharply defined nucleus accumbens and the ventral portion of the precommissural body or septal formation. Its own medial boundary is also sharply defined from the septal formation. The tuberculum apparently receives secondary olfactory fibers from both the lateral and medial olfactory tracts (Fig. 28), whose fibers may be seen bending down into the plexiform layer of the tuberculum on its lateral and medial borders. Beccari (1910) questions this (see page 190 above), finding evidence of other sources of origin (the pyriform lobe in particular) for the fibers of the external plexiform layer of the tuberculum. He thinks that the olfactory tract fibers, if present, exist only rostrally in this layer. It looks otherwise here, but this is not really decisive material.

In front of the strio-tubercular fusion (see Figs. 29-30), which takes place behind the caudal end of the posterior part of the anterior olfactory nucleus (see page 191) essentially in the fashion so clearly described by Livini (1908) in *Hypsiprymnus*, a rather wide *deep* plexiform layer, continuous with the superficial encircling plexiform of the section, intervenes between the tuberculum and the rest of the cellular formation of the section. This layer is crowded with a wealth of fibers of diverse origin and destination, whose adequate analysis is impossible in this series. They include fibers from the lateral olfactory tract, an-

terior olfactory nucleus, intermediate olfactory tract (probably both crossed and uncrossed), from the tuberculum, which are destined for the septal nuclei, hippocampus, frontal neopallial pole. Probably all of these systems are to some extent doubly oriented. And some of them also apparently contribute to the median forbrain bundle, fasciculus medialis telencephali (*f.med.t.*, Figs. 29-35), which begins collecting in the base of the septum quite far forward; these fibers are projection fibers to stem centers and probably from them also. It is the multitudinous *ascending* precommissural fibers which are mainly responsible for the differentiation of the rostral portion of the hippocampus in lower mammals.

The histological development of the tuberculum is in *Caenolestes* spectacular in the highest degree. The external cell layer of medium darkstaining pyramidal or polymorphic cells is thrown into battlement-like folds, interrupted irregularly by islands of Calleja (*i. C.*, Figs. 29-32), composed of extremely small, pale, round cells densely crowded, glomerulus-like roundish or vermiculate areas, and including sometimes a few pale giant cells. These masses vary greatly in size and shape, and they are so sharply delimited and so different from their surroundings as to suggest pathological growths. They probably correspond to Beccari's (1910) Type 2 islands, while deeper ones of the same general character belong to his Type 3 islands. The Type 1 islands consist of thickenings of the crenulations of the external cell layer; these are sometimes fringed with pale granules like those of the other islands. While these types are sharply differentiated, there are, as Becarri found, intermediate types. Small isolated cell masses in the external plexiform layer of the more caudal sections especially are all traceable into the main mass of the tuberculum. None of Beccari's figures show so great a histological complexity as *Caenolestes*. The conditions in the Virginia opossum are much less complex (Gray, 1924).

What is the function of these highly elaborated and integrated *organs within an organ*? They suggest some sort of elaborate rehandling and sorting of incoming stimuli—a physiological analysis by means of differential thresholds, effecting a secondary specificity from mass stimuli dissociated, reenforced and more or less independently projected? A searching study of the tuberculum at its most bizarre stage of development in the lower mammals ought surely to discover valuable clues to modes of nervous organization.

The deep layers of the tubercular "cortex" are best described in connection with the immense fiber and giant cell stream which, like a great diagonally slung hammock, extends from the ventral pyriform cortex

and lateral anterior commissure region across the base of the hemisphere to the ventro-medial "regio innominata", the vestibule to the thalamus and lower stem centers. This great and complex system, composed of an immense number and variety of fibers, both projection and associational, is certainly one of the striking features of the middle subventricular region of the hemisphere. It is generally called the basal olfactory bundle (of Wallenberg) in these lower mammalian brains. It forms a part of the practically unanalyzable mass of fibers (in this series) traversing the basal region of the hemisphere, and including such systems as the olfactory projection of Cajal, striatal systems (*ansa lenticularis*), etc., the whole mass, save the association fibers, drifting ventro-medially to join the medial forebrain bundle and continue spinalward with it. Therefore, following some writers, I have called it the *lateral limb of the medial forebrain bundle* (*fasciculus medialis telencephali, pars lateralis, f.med.t.l.*, Figs. 29-33). It is largely, in all probability, a doubly oriented system. Its more rostral portion is characterised by the presence of a great multitude of pale giant cells strewn among the fibers (nucleus of the basal olfactory bundle, Figs. 29-33). A similar condition occurs in the more median fiber tangle, whose giant cells correspond probably to the "border nucleus" of Völsch (1906). The *ansa lenticularis* component contains few cells. But for many of its fibers the cells of origin are the giant cells characteristic of the globus pallidus (*gl.p.*, Figs. 12, 15; *glob. p.*, Fig. 34), and these are just like those strewn so thickly among the fibers of the more rostral stream of the medial forebrain bundle, especially in its lateral limb and in the more medial portion, where they seem to correspond to the "border nucleus" of Völsch (1906). The globus pallidus cells are certainly motor projection cells, and very likely the other giant cells mentioned also send long axons to stem centers. The fact that the red nucleus is also mainly composed of pale giant cells of the same type further tends to support this supposition.

With the reduction of the lateral and basal olfactory centers in higher mammalian brains, the more posterior *ansa lenticularis* complex becomes so preponderant as to throw the more anterior olfactory complex into the shade, and so the name "*ansa lenticularis*" comes to be applied to the whole stream. But in lower mammalian brains it is the more anterior component, mediating correlated olfactory stimuli, which seems more conspicuous. Many longitudinal fibers and fiber bundles (association tracts between rostral and caudal regions of the base of the hemisphere) further complicate the situation.

Septal region.—The precommissural area of the median surface of the hemisphere is the external surface of the thick paraterminal body of Elliot Smith or septum of ordinary terminology. This region contains the ascending olfactory systems already enumerated (olfacto-septal, olfacto-cortical) and the descending precommissural fornix, septo-amygdaloid (Johnston's stria terminalis bundle 4), septo-habenular (of stria medullaris system) and median forebrain bundles. The bundle labelled olfacto-frontal (*tr.ol.fr.*, Fig. 27), extending from the frontal pole of the neopallium into the septum, where it mingles with the septal fibers which sweep laterally beneath the ventricle or below the lateral mass of the anterior commissure, cannot itself be disentangled and followed with precision in these sections. Arising in the frontal pole of the neopallium just in front of the laterally directed corona radiata fibers destined for the internal and external capsules, these fibers inevitably suggest the possibility of part of the coronal fibers being diverted, or rather persisting, medially into the septum as projection fibers to stem centers by way of the median forebrain bundle, as a vestige of what Edinger has called the septomesencephalic tract of submammalia. In this connection Pedro Cajal's findings (1917, 1919) in *Varanus* and *Lacerta*, of a septal passage for descending fibers from the entire cortex, including the depressed portion between the medio-dorsal hippocampal cortex and the lateral pyriform cortex, which he considers the "general cortex", seem highly significant. One is led to recall also that the only cortical projection tract in birds is a septal one, and that it does not proceed from what seems to be olfactory cortex. It is not inconceivable that in the neopallial frontal pole of the lowly mammalian type of brain, at a level where the hippocampal formation is still rather insignificant, the more medial neopallial fibers might have preserved the shorter septal path.

Two well developed nuclei lie in the septum—the *lateral* and *medial parolfactory nuclei* (Herrick). The lateral parolfactory nucleus (*nuc.pol.l.*, Figs. 28-30) occupies the septal wall lateral to the pre-commissural fornix fibers. It corresponds to Johnston's (1913) primordium hippocampi and not to his lateral parolfactory nucleus, which is here the nucleus accumbens (*nuc.ac.*), the medial portion of the head of the caudate nucleus. The medial parolfactory nucleus (*nuc.pol.m.*, Figs. 30-31), located medially and ventrally, corresponds to Johnston's nucleus of the same name, and, as he finds in the Virginia opossum and in other forms (Johnston, 1923), it passes back insensibly into the *nucleus of the diagonal band of Broca* (*nuc.d.b.*, Figs.

32-33). Apparently it grades also into the *preoptic nucleus* of the telencephalon medium (*nuc. prop.*, Fig. 34).

The *dorsal (d.)* and *anterior (c.a.) commissures* have already been briefly described in connection with the median section of the brain. The dorsal commissure will be considered also below in connection with the hippocampus. The *composition* of the anterior commissure offers in *Caenolestes* nothing of unusual interest. But in view of the disagreement among zoologists with regard to the assignment of *Caenolestes* to one or the other of the two marsupial subgroups—Diprotodontia and Polyprotodontia—with its bearing on marsupial distributional problems, the *arrangement* of the fibers of the anterior commissure was, since it seemed to be the only remaining anatomical evidence to be expected on the question, a matter of great interest (Obenchain, 1923a, 1923b). After an examination of the brains of every marsupial genus except *Caenolestes* (which was not available) Elliot Smith (1902a and b) found that all diprotodont brains exhibit one feature never found in any polyprotodont brain, or indeed in any other vertebrate brain. This exclusive diprotodont character he named the *aberrant bundle (fasciculus aberrans)* of the anterior commissure, considering it a true diagnostic character of diprotodont brains. It is merely the dorsal portion of the anterior commissure which in diprotodonts splits off from the rest of the commissure to pass upward by way of the internal instead of the external capsule, the common route in all other brains. The aberrant bundle is absent in *Caenolestes*, a fact which, if this feature be decisive, would ally it with the polyprotodonts. The brains of the fossil caenolestids, however, can never be known, but the presumption of the absence of the aberrant bundle in them also would perhaps be justifiable. The effect of this would occasion no further disturbance of Dr. Osgood's marsupial "family tree" than the lengthening of the polyprotodont bracket to include *Caenolestes*, leaving it still in place between the generalised polyprotodont *Perameles* and the diprotodonts of Australia; or at most, in view of the intermediate position of the exposed gyrus dentatus of *Caenolestes*, the latter might be shifted to a position between *Perameles* and *Notoryctes*.

*Pyriform Lobe**.—The pyriform lobe consists of two distinct parts: (1) the pyriform cortex, mostly confined to lateral wall of the hemi-

*The limits here assigned to the pyriform lobe do not exactly coincide with those assigned either by Cajal or Johnston, although the actual descriptions of the areas involved vary little or not at all. Cajal (1911) restricts the pyriform lobe to include, besides the amygdala, only the median pyriform cortex

sphere, between the rhinal fissure above and the rhinal arcuate and amygdaloid fissures below; and (2) the amygdaloid complex, mostly in the ventral wall between the amygdaloid and choroid fissures.

The *amygdaloid complex*, thanks to Johnston's illuminating comparative analysis (1923), which builds upon and completes Völsch's intensive study (1906), is no longer the mysterious territory it has been. Since it is, strictly speaking, a subcortical center, it will be described first here, leaving the pyriform cortex to precede the hippocampus. Owing to its enormous extent, the amygdaloid complex is perhaps the most spectacular part of the brain—unless it share this distinction with the tuberculum olfactorium. The superficial extent of the amygdaloid complex has already been described. Internally it exceeds this both in length and width, overlapping the posterior portion of the tuberculum anteriorly and the pyriform cortex laterally, and, as Johnston (1923) points out in the Virginia opossum, stretching rostrally far towards the anterior commissure. Behind the caudate nucleus and the putamen (*put.*) it floors the ventricle, while the cellular bed of its great fiber system, the stria terminalis (*st.t.*, Figs. 32 ff.) forms the median strip of the floor behind the anterior commissure. I am not, however, able to follow the stria bed in these sections into the anterior olfactory nucleus, as Johnston does in the opossum. It is clear anteriorly here only as it lies upon the anterior commissure and more caudally upon the internal capsule or cerebral peduncle.

The nuclei of the amygdaloid complex in *Caenolestes* comprise the six described by Johnston (1923), and include also the extra seventh one he found in the Virginia opossum: the nucleus of the lateral olfactory tract, the central, medial, lateral, basal, accessory basal, and cortical amygdaloid nuclei.

(his *anterior* pyriform cortex), receiving terminals of the lateral olfactory tract, and the posterior pyriform cortex, receiving no direct olfactory fibers (his superior temporal cortex). This makes it coincide approximately with the gyrus hippocampi of primates. Johnston (1915) defines the pyriform lobe as the gray matter underlying the lateral olfactory tract, which would include the entire anterior olfactory nucleus, but not the posterior pyriform cortex. By this definition the pyriform lobe and hippocampus are *directly* continuous anteriorly, but not in contact posteriorly, since the posterior pyriform cortex would be included in the neopallium. Since this area is the field of origin of the temporo-ammonic tract (Cajal), the main afferent tract to the hippocampus in mammals, it seems more logical to include it within the pyriform lobe. Its exclusion by Johnston would explain the complete caudal separation of the pyriform lobe and hippocampus in the turtle, *Cistudo carolina* (1915). The anterior pyriform cortex here coincides with Cajal's "frontal lobe", which he excludes from the pyriform lobe, on the basis that it, like the peduncular gray (anterior olfactory nucleus) receives mainly *collaterals* of the lateral olfactory tract.

The well delimited *nucleus of the lateral olfactory tract* (*nuc.tr.ol.l.*, Figs. 12, 15, 36-38) is a duplex structure, consisting of a large-celled dorsal part and a small-celled ventral part, separated by a narrow plexiform layer. It is tilted up rostrally, and shifted forward for about half its length above the caudal portion of the tuberculum, and therefore only its posterior half lies beneath the medio-ventral tubercle which in *Caenolestes* is apparently not so pronounced as in *Orolestes*, of which no sections are now available for comparison. The nucleus of the diagonal band (*nuc.d.b.*, Figs. 32, 33) curves around from the median wall here, and the diagonal band fibers pass laterally in the diffuse region above the posterior end of the nucleus of the lateral tract towards the pyriform cortex, probably also effecting connections with the intermediate region. The ventral small-celled portion of the nucleus is so sharply delimited as to appear almost encapsulated, an appearance heightened at its caudal pole, which extends beyond the dorsal part as a sort of island of cells in the plexiform layer of the hemisphere. The corresponding nucleus of the Virginia opossum exhibits neither the forward shifting above the tuberculum nor the histological differentiation seen in *Caenolestes* (Gray, 1924). The nucleus of the lateral olfactory tract receives secondary olfactory fibers from the pars ventralis of the lateral olfactory tract (unlabelled, Figs. 36-38), and dorsally a great fan of fibers from the stria terminalis (*st.t.l.*, Figs. 33-36), its most anterior contingent. Johnston (1923) identifies this with the commissural bundle of the stria, but these sections do not actually permit this.

The *central nucleus of the amygdala* (*c.*, Figs. 12, 15; *nuc.amg.c.*, Figs. 35-37) lies above the nucleus of the lateral olfactory tract. It is confluent with the strongly developed "intercalary plate" (*int.plate*, Fig. 36) of Johnston (1923), which is the most hypertrophied part of the stria bed. The central nucleus advances farther forward than any other amygdaloid nucleus, but apparently not so far as Johnston found in the Virginia opossum, up to the region of the anterior commissure. Its limits are not well defined nor its cellular structure striking.

The *medial amygdaloid nucleus* (*m.*, Figs. 12, 15; *nuc.amg.m.*, Figs. 36, 37) occupies the ventro-medial angle of the hemisphere, and also lacks well defined limits or strikingly marked cell structure. It is impossible to fix its caudal limit here. It also is continuous with the intercalary plate and with the central nucleus, as Johnston found.

The *lateral amygdaloid nucleus* (*l.*, Figs. 12, 15; *nuc.amg.l.*, Figs. 35-39) is the "poststriatum" of earlier writers. It is a large and ex-

extremely well defined antero-posteriorly elongated nucleus, oval in section and characterized by good-sized pyramidal or polymorphic cells. It lies in the concavity of the external capsule (*cap.e*), between it and the ventral part of the putamen (*put.*), and behind the latter it rises to form the more lateral part of the ventricular floor. It clearly receives external capsule fibers, more probably originating in the pyriform cortex, and it seems also to receive a large number of stria terminalis fibers, coursing horizontally just above the basal amygdaloid nucleus (*st.t.* 3, Fig. 38). This last is contrary to Johnston's observations.

The large-celled well defined *basal* nucleus of the amygdala (*b.*, Figs. 12, 15; *nuc.amg.b.*, Figs. 37-42) arises medial to the more caudal portion of the lateral nucleus, near its ventral border, and increases to large proportions as the lateral nucleus diminishes. Behind the flattened tail of the caudate nucleus it occupies a large part of the floor of the ventricle, extending far medialward and lying caudally directly beneath the ependyma. Both it and the lateral nucleus are in cellular continuity with the deep cells of the ventral border of the pyriform cortex at the level of the amygdaloid fissure. Johnston (1923) considers them to have been derived from the pyriform cortex by a process of infolding along this line, and the situation in *Caenolestes* seems to support this view.

The *accessory basal* nucleus of the amygdala (*b.ac.*, Fig. 15; *nuc.amg.b.ac.*, Figs. 37-38) is a less clearly defined nucleus of medium dark cells lying among the ventral fibers of the external capsule as they fan out in the postero-ventral amygdaloid region, below the caudal ends of the lateral and basal nuclei and obscurely confluent with them. It is not only present in *Caenolestes*, but it would be only too easy to subdivide the heterogeneous caudal amygdaloid region into further nuclei.

The *cortical* amygdaloid nucleus (*amg.*, Fig. 12; *nuc.amg.cort.*, Figs. 38-43) forms almost the entire superficial portion of the amygdaloid complex. Laterally it approaches or adjoins the ventral edge of the pyriform cortex and medially the ventral edge of the hippocampus behind the choroid fissure (*fs.ch.*) along an internally well marked and an externally partially obvious medial continuation of the amygdaloid fissure (*fs.amg.m.*). Its histological structure assumes in places a structure similar to that of the pyriform cortex. Its deeper portion is very heterogeneous. Some confusion, however, results from the tangential nature of the posterior sections of the hemisphere. The cortical amygdaloid nucleus receives secondary fibers by way of the pars ventralis of the lateral olfactory tract. The hippocampal-amygda-

loid junction (*subic. (amg.)*, Figs. 40-42) along the median extension of the amygdaloid fissure is also evidence of fiber connections, and Johnston (1923) has demonstrated them in the Virginia opossum.

The *stria terminalis* (*st.t.*, Figs. 32-38), the great fiber system related to the amygdaloid complex, has also been carefully analyzed by Johnston (1923). Practically its entire course is quite clear in these sections, and rostrally and caudally its five component bundles as identified by Johnston, can easily be recognized, but in the compact medial portion of the tract, where it lies in the stria bed between the cerebral peduncle and the ependyma of the lateral ventricle they cannot be individually identified. Johnston's numbers have been affixed to these bundles as traced by him in the Virginia opossum. Rostrally they are certainly correctly applied, but caudally they are applied without direct evidence from the brain of *Caenolestes* of continuity with the respective rostral bundles.

The lateral part of the anterior olfactory nucleus passes without break directly over into the least differentiated anterior pyriform cortex behind it. The *cortex of the pyriform lobe* covers most of the lateral surface of the pyriform lobe. Antero-laterally it may be divided into three regions: the anterior pyriform cortex (*cx.pir.a.*, Figs. 13, 15, 26-31; area piriformis anterior, Gray, 1924; frontal lobe of Cajal, 1911), which passes more caudally by very gradual transition into still more differentiated medial pyriform cortex (*cx.pir.m.*, Figs. 13, 15, 32-34; area piriformis medialis, Gray; anterior pyriform cortex of Cajal; this in turn merges more abruptly into the most specialized posterior pyriform cortex (*cx.pir.p.*, Figs. 12, 13, 15, 42-44; area piriformis posterior, Gray; superior temporal or angular nucleus or center of Cajal. Since the lateral part of the anterior olfactory nucleus merges insensibly with the anterior pyriform cortex, and the latter merges insensibly into the medial pyriform cortex, no definite boundaries can be made out between them. The second cortical layer (not counting the external fibrillar layer) becomes progressively more and more condensed, and any lines of demarcation are, in these sections, merely arbitrary. But the posterior pyriform cortex (*cx.pir.p.*, Figs. 42-44), which is on a different plane functionally from the remainder of the pyriform cortex because (by Cajal's definition) it receives no secondary olfactory fibers, also differs histologically. Its histological development is connected not only with the absence of secondary olfactory fibers, but perhaps even more with the increase of non-olfactory fibers, and its consequent elevation into an associational area second

in rank only to the neopallium. It displays a very striking anatomical character. This is mainly due to the development within it of a wide and very dense plexus in the third layer of the cortex. In *Caenolestes* this plexus, which according to Cajal is in the mouse of extraordinary density, almost fills the pyriform wall dorso-caudally. Its broad, rounded head rises slightly above the level of the rhinal fissure. It contains many pale giant cells, found nowhere else in the pyriform cortex of *Caenolestes*. The posterior pyriform region is the field of origin of the great temporo-ammonic or angular bundle of Cajal (1911, speno-ammonic, 1906), which delivers a huge stream of highly correlated olfacto-somatic impulses to both the ammon's horn and the gyrus dentatus, for almost their entire length—certainly reaching as far forward in *Caenolestes* as the anterior level of the commissural region, where its structural influence is suddenly and strikingly felt. Since it mingles with other pyriform and with neopalial association fibers to form the cingulum limitans (*ci.lin.*, Figs. 30-42) at the inner dorsal angle of the lateral ventricle, and the cingulum ammonis (*ci.am.*, Figs. 30-42) in the outer plexiform layers of the ammon's horn and gyrus dentatus, it is not separately named in the sections given here. In the posterior region of the hemisphere this avalanche of fibers may be seen pouring above and behind the ventricle into the presubicular and subicular regions and through them, either directly or indirectly, as perforating fibers to the ammon's horn and gyrus dentatus.

Anteriorly the lateral olfactory nucleus certainly sends tertiary olfactory fibers above the olfactory ventricle into the anterior hippocampal formation (Figs. 26-27). The sections suggest that the dorsal path between the pyriform and hippocampal cortices might perhaps be patent for practically the entire length of the neopallium, in the deep layer of the corona radiata. Owing to the height of the rhinal fissure the distance is nowhere very great, and this would certainly be the shortest path for the more dorsal pyriform cortex. The anterior pyriform cortex must also contribute very largely to the subventricular systems, both to the olfacto-septal, olfacto-hippocampal and olfacto-frontal systems anteriorly, and to the more posterior and extensive complex included under the head of the lateral limb of the medial forebrain bundle, as explained above (see page 198). The great efferent pyriform path to stem centers is of course the olfactory projection path of Cajal, probably, like so many of these fiber systems, a doubly oriented one. The stria terminalis component of this system (Johnston's bundle 2, 1923) arises from the amygdaloid complex (so far as it is a descending bundle). The fiber connections between the

pyriform cortex and the amygdaloid complex have already been mentioned (page 203).

In addition to the antero-posterior divisions of the pyriform cortex, Gray (1924) has described several narrow longitudinal bands along the boundary fissures in the Virginia opossum. Three such areas accompany the rhinal fissure: the area perirhinalis, the area piriformis dorsalis, and the area piriformis fissuralis, in order from above downwards. The extreme shallowness of the rhinal fissure in *Caenolestes* is not conducive to great development of these areas. The area perirhinalis, the transition area between neopallial and pyriform cortex, is probably individualized, but the other two are feebly developed and not distinct from one another. Ventrally also the area piriformis ventralis and the area subpiriformis of Gray tend to fuse into one band quite well marked, the ventral pyriform cortex (*cx.pir.v.*, Figs. 27-34), especially anteriorly where it describes a deep reentering angle subjacent to the massive lateral olfactory tract. This band is here regarded as following caudally upon the latero-ventral part of the anterior olfactory nucleus (*nuc.ol.ant.l.v.*, Fig. 26). Behind the tuberculum, and even more rostrally, where the diffuse anterior perforated space (*l.perf.a.*, Figs. 35-38) of Johnston (1923) intervenes between it and the nucleus of the lateral olfactory tract, the pyriform cortex exhibits a very sharp ventral margin back to the point where it gives evidence of infolding at the level of the lateral and basal amygdaloid nuclei. Still further back it becomes more or less continuous with the outer layer of the cortical amygdaloid nucleus, and histological differentiation tends to fade out. The tangential nature of the sections in this caudal region is somewhat confusing, and control by sections in other planes is unfortunately lacking.

HIPPOCAMPAL FORMATION

The hippocampal formation of *Caenolestes* (Figs. 6, 12, 17, 26-43) begins with a small patch of cells condensed upon the outer aspect of the rostral border of the anterior olfactory nucleus, at its dorso-medial angle, that is to say, immediately caudal to the bulbar formation (*hip.a.*, Figs. 12, 17; *cx.hip.a.*, Figs. 26-29). It thus takes part in the formation of the rhinencephalic ring surrounding the olfactory ventricle, by means of which the two olfactory cortices are indirectly continuous with one another through the anterior olfactory nucleus. It is to be noted that while the hippocampal formation begins at the *very* rostral margin of this ring, in its intrabulbar portion, just caudal to the olfactory formation itself, a similar lateral (pyriform) condensation falls

short of this, being separated from the bulbar formation by uncorticated anterior olfactory nucleus. The explanation of this precocity of hippocampal differentiation might conceivably be tied up with the prevailing latero-ventral drift of the secondary olfactory fibers toward the lower olfactory correlation centers of the lateral wall and base of the hemisphere—pyriform lobe and tuberculum olfactorium. This trend is clearly brought out in the bulb, where there is never any considerable accumulation of secondary fibers in the dorso-medial region, those arising there apparently for the most part making as rapidly as possible for the laterally and ventrally situated lateral, intermediate and medial olfactory tracts. The only exception to this is the presence of the small more median portion of the pars dorsalis of the lateral olfactory tract (*tr.ol.d.m.*, Fig. 26), which may, and probably does (as silver preparations indicate in the Virginia opossum, Herrick, 1924a), give off some collaterals to the hippocampal formation, which is, however, already strongly under the influence of tertiary olfactory fibers. It seems but natural that such a region, receiving only a minimum of direct olfactory fibers and beginning almost, if not quite, at once to receive tertiary olfactory fibers from the more lateral portions of the anterior olfactory nucleus (these can be seen crossing above the olfactory ventricle) and presently still more highly correlated olfactory stimuli from the tuberculum by way of the septum, should undergo accelerated differentiation in comparison with a region preponderantly invaded by secondary olfactory fibers (see page 219).

Dorsally the continuity between the hippocampal formation and the pars dorsalis of the anterior olfactory nucleus becomes transformed at the junction between the neopallium and the rhinencephalon into a continuity between the hippocampus and the neopallium on the one hand, and between the pyriform cortex and the neopallium on the other (Figs. 26-27). Ventrally the hippocampal formation, at first continuous with the pars posterior of the anterior olfactory nucleus (*nuc.ol.ant.p.*, Figs. 27-28), breaks away from the latter and becomes separated from it by an increasing interval of diffusely scattered small cells through which some fine pale fibers pass from the deeper septal region to the plexiform layer (Fig. 29). This area is replaced more caudally by the medial parolfactory nucleus (*nuc.pol.m.*, Figs. 30-31).

The ventral end of the precommissural hippocampus (Fig. 29) thins to a slender needle point which, as it recedes upward in the septal wall, approaches the pial surface. This slender column of cells widens

above very quickly as it merges with the neopallial gray. It soon begins to show a slight concavity toward the pial surface (*fs.hip.*, Fig. 29), which more caudally comes to expression as the external hippocampal fissure (*fs.hip.*, Fig. 30). The cells diminish in size and increase in density from above downward, to the sharply pointed ventral end of the hippocampal formation, which curves upwards and backwards in line with the ventral (medial) edge of the definitive gyrus dentatus at the fimbrio-dentate fissure near its rostral end at the antero-dorsal angle of the hippocampal commissure (*fs.fim.d.*, *c.d.*, *cx.dent.*, Figs. 30-31). As the hemisphere increases in height the hippocampal formation lengthens, the upper thicker and larger-celled portion more rapidly than the lower, so that it comes to exceed the lower smaller-celled portion in length. It becomes at the same time thinner and denser. It is interesting to note that rostrally the hippocampal formation differentiates more rapidly at its peripheral (dentate) border, and that differentiation travels upward, the orderly condensed files of definitive ammon's pyramids appearing tardily. Caudally also we find the cortex dentatus apparently leading the way in the formation of the recurved temporal pole of the hippocampus (see page 215).

When I was making the first Edinger drawings I noted a curious breaking up or disorganization in the precommissural hippocampal cell plate, due to the loosening up and paler staining of some of the cells of the upper region (*, Figs. 12, 17a). This rift or line of fracture I interpreted as the locus of the interpositio medialis, the break between the definitive gyrus dentatus and the ammon's horn. But to my great surprise the interpositio medialis formed rather suddenly at a distinctly lower (more ventral) level at the anterior border of the dorsal commissure (*ip.m.*, Fig. 31). A provisional explanation of this puzzling rift in the cell plate suggested itself later, however, in connection with Cajal's statement (1911, II., p. 754) that the gyrus dentatus and the more ventral half of the ammon's horn (his extraventricular ammon's horn or region of grand pyramids) seem to form an indissoluble anatomical and functional unit, owing to the exclusive distribution of efferent gyrus dentatus axons (the "mossy fibers") to the extraventricular ammon's horn. The line of fracture in question might presumably be interpreted, then, as the upper limit of the distribution of the "mossy fibers", the boundary between intra- and extraventricular ammon's horn. (For a provisional explanation of its absence in the supra- and postcommissural ammon's horn, see page 209 below). The material available here does not furnish proof of this hypothesis, which is offered merely as a suggestion.

At the level of the anterior face of the dorsal commissure the pre-commissural hippocampus, which, as Elliot Smith (1896c) long ago pointed out, exhibits in marsupials all stages of development to be seen in the reptilian hippocampus, suddenly assumes the definitive mammalian appearance of double interlocked arcs by the abrupt formation of the interpositio medialis separating the ammon's horn and the gyrus dentatus, and the synchronous humping up of the latter into a horse-shoelike form into whose ventral concavity the terminal lamina (Levi, 1904; nucleus fasciae dentatae, Elliot Smith, 1896c) of the ammon's horn is displaced. The explanation of these sudden changes is probably three-fold: external pressure, internal pressure and neurobiotactic influence. External pressure is exerted on three sides: from above by the downward pressure of the neopallium (as pointed out by Elliot Smith, Levi and many other neurologists), medially by the opposite hemisphere, and ventrally by the rigid barrier of the dorsal commissure and subpallial structures, beneath the choroid fissure (the atrophic choroid barrier so strongly emphasized by Levi, 1904). The internal pressure arises from the intrinsic growth of the hippocampus, chiefly the ammon's horn, which must crumple or roll to adjust itself to the space allotted. The neurobiotactic influence is in the first place due to the "perforating fibers" of the great temporo-ammonic tract and fibers associated with it in the cingulum limitans and cingulum ammonis (see pages 201 and 205). These fibers distribute equally to the gyrus dentatus and the ammon's horn. Those fibers which discharge into the granule cells of the gyrus dentatus break across the hippocampal fissure as it deepens and cause a more or less partial obliteration of the fissure by a secondary fusion of its two lips. They pull the sheet of granules upward through its entire length along a longitudinal axis (nearly median in *Caenolestes*), so that in cross section the gyrus dentatus describes a horseshoe curve. The efferent fibers of the gyrus dentatus (axons of the granules, "mossy fibers"), converging upon the terminal lamina of the ammon's horn, cause a break at the position of the interpositio medialis (*ip.m.*, Fig. 31), and the displacement of the lamina terminalis into the ventral concavity of the gyrus dentatus. Figure 11a, b, c illustrates diagrammatically the mode of formation of the definitive mammalian "hippocampal figure" in *Caenolestes* under the three influences named. The arrows indicate the directions in which the force is applied and its relative strength.

It might be possible to invoke also Kappers' ever useful concept of neurobiotaxis to explain the obliteration of the precommissural rift or fracture in the hippocampal formation (*, Figs. 12, 17a) noted above

page 208) as the possible boundary line between the intra- and extraventricular ammon's horn; the extraventricular ammon's horn alone receives "mossy fibers" from the gyrus dentatus, a circumstance that might be held to account for the rift or fracture line in question; but the grand pyramids of this region, according to Cajal, send recurrent collaterals into the external plexiform layer of the ammon's horn (stratum lacunosum) which rake the whole extent of the ammon's pyramids, both intra- and extraventricular, and so tend to close any gap in them. Furthermore the afferent commissural fibers in the alveus might contribute to the same result, as well as to the formation of the interpositio *lateralis*, by a general neurobiotactic compacting of the whole line of pyramids. More than this, the temporo-ammonic fibers which distribute to the ammon's pyramids either by way of the superficial plexiform layer external to the stratum lacunosum, or by way of the alveus (Cajal's temporo-alvear tract), engage, according to Cajal, not only the intraventricular but also some of the nearer extraventricular pyramids, perhaps most of them. At any rate it would seem as though the initial agent in the production of such dislocations as those considered is always the functional activity of the intrinsic structures concerned, and not the mechanical action of the fibers which may later not only occupy the break, but enlarge it, even to the extent, conceivably, of interfering with the function which originally produced it.

The *dorsal or hippocampal commissure* (*c.d.*, Fig. 31) exhibits a narrower and denser dorsal portion, the *psalterium dorsale* (*ps.d.*), which Cajal (1911) considers the commissural path of the crossed portion of the temporo-ammonic system, and the wider and much more diffuse *psalterium ventrale* (*ps.v.*) the commissure of the ammon's axons. Many cells are mingled with these latter fibers (the nucleus of the commissure), so that on the whole the dorsal commissure in comparison with the much denser anterior commissure seems larger than it really is. The precommissural fornix fibers (*f.prcm.*, Fig. 30) may be seen passing vertically downward in front of the commissures, and above the psalterium dorsale the dense short mass of the fornix longus, bending downward and partly interweaving with the psalterium dorsale, forms the "knieförmiges Bündel" (Koelliker; Livini, 1908); they belong to the median striae Lancisii (Johnston, 1913). The *descending columns* of the *fornix* (*c.for.*, Fig. 31) collect in the usual way close to the midline as deeply stained oval bundles above the anterior commissure and pass down behind it into the hypothalamic region. Two similar oval bundles of pale fibers located between the

fornix bundles and the anterior commissure on either side of the inferior recess (*r.i.*) may be the supracommissural bundle of the stria terminalis (Johnston's bundle 4, 1923). They seem to be receiving more lateral fibers disposed in the same diagonal direction as some of the darker fibers entering the fornix bundles. I cannot tell whether they are of septal or hippocampal origin, or both. These bundles cannot in my sections be definitely followed laterally into the stria bed above the anterior commissure. There is also the possibility that they belong to the stria medullaris system, but I cannot follow them to the habenula.

As Johnston (1913) was the first to point out, marsupials possess well developed *medial striae Lancisii* above the dorsal commissure, a level at which full-bodied hippocampus also exists, proving that the lateral and not the medial stria of the indusium are the vestigial remnants of degenerate hippocampus in "callosal" brains. These median striae are perfectly clear in *Caenolestes*, but they are omitted from the reduced sections figured here. They are seen as cross cut fibers upon the dorsal surface of the hippocampal commissure and as vertical fibers cutting across the anterior dorsal commissure fibers and partly interwoven with them (see knieförmiges Bündel, page 210).

In connection with the structures under discussion it may be well to describe the form taken by an interesting ventral diverticulum of the superior recess (of Elliot Smith), which has been described and figured in sagittal section of Johnston (1913) for the Virginia opossum (Fig. 16b). In *Caenolestes* this subcommissural pouch (Fig. 16a) runs its course between section 510 (Fig. 30 shows the beginning of the glial mass in which it is embedded) and section 528. The dorsal recess or sac bends down around the rostral surface of the dorsal commissure, carrying with it membranous roof tissue which hypertrophies in the septum between the commissures and in the pial side of the precommissural area to form a rather thick mass of glial tissue between the medial fornix fibers. This mass, composed of densely crowded fine pale granules, is sharply separated from the nervous tissue adjacent to it. Rostrally it is bifurcated and caudally it comes to a median point in the septum beneath the dorsal commissure. It contains a lumen of similar shape, lined with ependyma, which ends caudally as a median recess in front of and above the inferior recess and bifurcates rostrally to end in two smaller diverticula in the precommissural septal walls. In *Caenolestes* the communicating canal to the dorsal recess or pouch is collapsed and obscured in dark-stained membranes and blood vessels. Figure 16a shows a reconstruction of

the ventral diverticulum of the dorsal recess in *Caenolestes* in the horizontal plane below the entrance of the communicating canal; figure 16b the sagittal section of this diverticulum in the Virginia opossum from Johnston (1913, part of figure 35, redrawn).

The middle portion of the hippocampus (Figs. 32-37) is naturally less interesting than the two extremities, which offer more developmental and adaptational clues, and it therefore calls for but few remarks here. With respect to the amount of gyrus dentatus exposed upon the median surface of the hemisphere, *Caenolestes* is intermediate between *Perameles* and *Notoryctes*, a fact which indicates that the neopallial pressure, apparently the chief varying factor here, exceeds that in the latter and falls below that in the former. According to Elliot Smith (1895b, Fig. 6) *Notoryctes* has the least extensive neopallium found among mammals, and "in no other animal does one find the simplicity of arrangement which the hippocampus of *Notoryctes* presents, an appearance which recalls the foetal hippocampus of *Perameles* or *Macropus*." (This statement refers only to what I have called the "hippocampal figure" as seen in cross section and not to the development of the temporal pole of the hippocampus, which will be discussed below.) The intermediate condition of the exposed gyrus dentatus in *Caenolestes* is especially interesting in view of the similarly intermediate condition of the cerebellum in *Caenolestes*.

It should be held in mind that the more caudal sections of the hippocampus become progressively more tangential to the structure itself and present an increasing distortion of the "hippocampal figure" of double interlocked arcs, as well as an exaggeration of its size. We should have a true picture of the hippocampus only if we could make all sections vertical to the hippocampal axis, which, since that is curved, would make them radial. The sections show that the hippocampus as a whole has in *Caenolestes* not yet "turned the corner"; that is to say, the typical "hippocampal figure" does not appear twice, like an object and its inverted image, in any one cross section, as it does, for example, in the rabbit's brain (Fig. 20a, b, c, Winkler-Potter, redrawn); or once, the "mirror image" only, as, for example, in the lion (Fig. 21, Elliot Smith, redrawn) and in man. In *Caenolestes* the gyrus dentatus alone has recurved and therefore it occurs in two places, one dorsal and one ventral, in some sections (Fig. 39). Neither the fimbria nor the ammon's horn has, however, recurved, and hence the ventral hippocampal figure of double interlocked arcs is incomplete and the temporal hippocampus is practically unrecognizable at first glance. The Virginia opossum has advanced one step farther in

the recurving process by the formation of a small forwardly directed pouch on the rostral face of the ammon's horn at its ventral border below the caudal end of the fimbria (Fig 18a-g, cross sections, from unpublished drawings, Streeter, redrawn; Fig. 18h, reconstruction made from same). This pouch is evidently formed in response to the neurobiotactic attraction of the gyrus dentatus, which has here grown forward beyond the field of the unrecurved ammon's horn. This change still does not result in a complete "hippocampal figure". But in higher marsupials, like *Hypsiprymnus* (Fig. 19a, b, c, Livini, 1908, redrawn), the temporal development has advanced to a point which allows the typical hippocampal figure to appear twice in the same section.

In *Notoryctes* (Fig. 22, Dart, 1920, Fig. 13, redrawn), to my great surprise, the temporal pole of the hippocampus is apparently quite as well developed as in *Hypsiprymnus*, and considerably better developed than in *Caenolestes* and the Virginia opossum. *Notoryctes* is a sightless form without external ears, and the elongation of the hippocampus and great amount of recurving of its temporal pole is perhaps to be regarded as compensatory, in view of the absence or extreme reduction of the visual and auditory systems. The median surface of the hemisphere (Elliot Smith, 1895b, Fig. 1) has a peculiar peaked appearance, which may perhaps be due to the elongation of the hippocampus and its consequent caudal bowing. The pressure of the slightly developed neopallium is not sufficient to cause a great degree of inrolling, not so much as in *Caenolestes*. *Ornithorhynchus* also has an elongated slender but well formed temporal pole of the hippocampus (Elliot Smith, 1896b, Figs. 4, 5, 5", pp. 472-3—the section given through the "tail" of the hippocampus is not very near its temporal end, however). In this case the elongation of the hippocampus and its temporal recurving, which is pronounced, is probably to be correlated with the very *great* size of the neopallium and consequent compression of the hippocampus. I do not know the condition in *Echidna*, but should expect a similar situation from similar conditions. The temporal pole of *Perameles* is also unknown to me.

In the rabbit's brain, a "callosal" one, in which the hippocampal commissure is caudally displaced by the corpus callosum, there are also two typical hippocampal figures, posed in opposite directions, joined by the tangentially cut fimbria (Fig. 20a, b, c, Winkler-Potter, 1911, pls. XII, XIII, XIV, redrawn). Here we have two exactly identical and completely detached images in front of the middle of the fimbria (Fig. 20a and b). I do not know whether this is

the case also in *Hypsiprymnus* (Fig. 19b). Livini does not give a section showing the fimbria in two parts. The more rostral section (Fig. 19a) shows, however, a little ventral "island" which may be the projecting tip of the uncus, and between this and the next caudal section given the fimbria may appear in two parts; but even so it would probably be less recurved than in the case of the rabbit.

In the case of the higher and more microsmatic mammals, as in the lion (Fig. 21), and in man, the dorsal "figure" disappears as the corpus callosum elongates and the supracallosal hippocampus (lateral striae Lancisii and the accompanying gray) stretches, and only the ventral figure persists completely. We can thus easily assemble a short series which will adequately illustrate the remarkable complete reversal of the hippocampus in the course of phylogeny by the agency of the combined action of callosal growth, tremendous neopallial hypertrophy, and the anchoring of the hippocampus in a rostral position by the pyriform cortex, which is itself strongly anchored to the bulbar formation. The tuberculum and the amygdala, as they diminish, retain their old places one behind the other between the bulb and the temporal end of the hippocampus, and medial to the lateral peduncular gray (lateral olfactory gyrus) and pyriform cortex. The neopallium expands enormously in the caudal direction, further accentuating the hippocampal reversion. The pyriform cortex loses its distinctive histological character and assumes a progressively more neopallial appearance, until the primate condition is attained, where, as "gyrus hippocampi" it becomes histologically practically identical with the neopallium (save in the presence of the external fibrillar layer in its anterior portion). The posterior pyriform region, that devoid of secondary olfactory fibers and richly supplied with non-olfactory fibers, begins, even in the Virginia opossum, (Gray, 1924) to resemble neopallial cortex rather closely along the border for the caudal prolongation of the rhinal fissure. In the mouse, as Cajal's (1911) intensive studies show, this area has developed an exceedingly characteristic histological structure of its own, and assimilation to the neopallial cortical pattern has not apparently made so much progress. The same is true in *Caenolestes*, in which the boundary between the neopallium and the posterior pyriform area is pretty definite (Figs. 41-42), so far as can be construed from frontal sections. The conditions existing at the temporal extremity of the hippocampus in *Caenolestes*, the Virginia opossum, *Hypsiprymnus* and the rabbit certainly stress the dynamic character of the temporal displacement of the hippocampus—"the brain is not a rigid mosaic of morphological units which were laid down in the primordial vertebrate

and thereafter preserved inviolate" (Herrick, 1922, p. 199). The little, rostrally directed ammon's horn pouch in the Virginia opossum is clear evidence of the regulative nature of the changes going on. In *Caenolestes* the forward advance of the temporal gyrus dentatus has not removed it from the unrecurved ammon's field, but in the Virginia opossum this would have happened if the ammon's horn had not responded by the formation of the pouch.

It is to me rather surprising to find the gyrus dentatus leading in the reversal of the hippocampus when rostrally it appears in *definitive form* tardily. But this tardiness is perhaps more apparent than real, since the lower portion of the precommissural hippocampus is in mammals, as Elliot Smith has stated for the same region of the posterior part of the reptilian hippocampus, clearly "on the way" towards differentiation into definitive gyrus dentatus. In the mammalian precommissural hippocampus it is the ventral portion, that directly continuous with the definitive gyrus dentatus, which takes the lead in differentiation. Differentiation apparently travels from below upwards, and the definitive ammon's horn or rather, perhaps, its "intraventricular" portion, is perhaps the last to develop its distinctive structure and extent. It is at least the last to *begin* differentiating. Pedro Cajal's illuminating studies on the reptilian brain (*Varanus* and *Lacerta*, 1917, 1919) by the silver methods show that in the ventral or medial small-celled portion of the hippocampal formation (*quasi* gyrus dentatus) there is a clear transition in cell type from the deeper to the outer cell ranks. The cells of the outermost cell ranks (nearer the pia) are practically true granules, the innermost true pyramids of the ammon's type. This is very significant, and it would not, I think, be surprising to find in full-bodied precommissural hippocampus of lower mammals an analogous state of affairs. The situation in this *Caenolestes* series strongly suggests, far in advance of the interpositio medialis and the inrolling of the gyrus dentatus, that the ventral cells of the more ventral region of the hippocampus are rapidly verging towards the true granule type, so that the transition to definitive gyrus dentatus involves apparently no sudden cytological changes. The hippocampus is apparently affected by two distinct waves of differentiation, both starting at the ventral border and traveling upward. The first develops cells of pyramidal type: the second, following in its wake, transforms pyramids into granules (phylogenetically speaking). The first wave affects the entire width of the hippocampal formation, reaching its dorsal border (locus of the interpositio lateralis), the second stops at the level of the interpositio medialis, affecting the "dentate" region but

never the ammon's region above it. The two main types of cells are perhaps to be interpreted as an expression of the enhancement of function in the hippocampus by its organization into a highly integrated duplex structure, the ammon's horn from which spring the long projection axons (which alone form the efferent path of the hippocampus), and the gyrus dentatus, whose shorter axons of a specialized type ("mossy fibers") deliver stimuli to part of the ammon's pyramids directly and to many others indirectly (recurrent collaterals of grand pyramids). We can never perhaps get the complete phylogenetic story in any one form, certainly not in the lowest and highest, where either later or earlier steps fail to appear.

Although the general form of the ammon's horn and the gyrus dentatus cell sheets in *Caenolestes* is really simple, it is difficult to show them both in the same diagram. Therefore dissected reconstructions (Figs. 17a and 17b) attempt to show them separately, one or the other being cut away at the commissural level of the hippocampus. The two parts are shown in place together in figure 12. It is seen that the gyrus dentatus takes the form of a sort of helmet or hood with the top of the crown or apex pointing caudally; the front or visor of the helmet corresponds to the elongated dorsal portion of the gyrus dentatus extending above the commissure where it ends in a little pouched thickening (Fig. 30); the back or neck portion corresponds to the ventral or temporal recurved part of the gyrus dentatus. The ammon's horn sheet takes the form of a loose scroll, greatly widened posteriorly in the sections (Figs. 38-40), so that it is really more like a cornucopia, with the wide end caudo-ventral. The upturned median flap of this scroll, covered with a thin coating of alveus fibers, is exposed upon the median surface of the hemisphere, save where its caudally directed corner is inserted into the hooded portion of the gyrus dentatus. This upturned ammon's horn forms Elliot Smith's extraventricular or inverted hippocampus ("dorsales Blatt" of Koelliker). It is seen in the sections to diminish in length as its lower edge recedes upward between the approaching dorsal and ventral portions of the gyrus dentatus (Figs. 34-40), which still more caudally unite to form an oval ring (the "crown" of the hood) containing the caudal tip of the inverted hippocampus (a few scattered cells, Fig. 41). At the caudal end of the choroid fissure the continuity of the two parts of the ammon's horn (in sections) is dissolved (Figs. 40-42), and the lower end of the extraventricular ammon's horn begins to recede upward, while that of the intraventricular ammon's horn seems to turn laterally under the median angle of the ventricle (Völsch's

subventricular hippocampus, 1906), where it comes in contact with the amygdaloid complex (stria bed and cortical amygdaloid nucleus). This contact extends backward along the line of the medial continuation of the amygdaloid fissure (*fs.amg.m.*, Figs. 40-43), always very strongly marked internally and sometimes externally apparent. As the sections show, the more temporal extremity of the gyrus dentatus (Fig. 39) is contiguous with the ammon's horn and perhaps with the amygdala where the two structures adjoin. The fiber connection between the hippocampus and the accessory basal nucleus of the amygdala (Johnston, 1923, Fig. 55) is not apparent in this series. This by no means proves its absence.

The emphasis here placed upon the amount of recurving of the hippocampus is intended to apply specially to mammals, where it is to be considered in connection with the inrolling of the hippocampus under the double necessity of enhancement of function and economy of space. Even mammals, as we have seen above (*Notoryctes*), do not present a quite orderly series in this respect. The degree of recurving *alone* is not a criterion of advancement in hippocampal development. In some reptiles (*e.g.*, *Cistudo carolina* (Johnston, 1915, Figs. 6, 12, 13, 45) the recurving of the temporal pole is more pronounced than in *Caenolestes*—in single cross sections two separate "hippocampal figures", complete for this brain, appear. The factors provoking the recurving in such cases are easily recognisable and need not be detailed here. The factors operating in mammals to bring about inrolling have apparently provoked a method of increase for the hippocampus, which, in concert with other existing conditions, may result in less recurving of the temporal pole than in lower forms. In higher forms the caudal displacement of the hippocampus due to callosal elongation, reverses or recurses the *inrolled* hippocampus exactly as in the case of simpler brains alluded to. What we seem to have always before us is the structural record of the solution of various problems of regulative behavior, from which we may attempt to reconstruct a phylogeny of function. We should remember that there are always a great number of factors working in concert or as more or less independent variables, and that the outstanding potency of any particular factor does not mean its exclusive activity.

Above the medial amygdaloid fissure (*fs.amg.m.*, Figs. 6, 12, 17, 40-43) the dorsal or subicular border of the ammon's horn adjoins, as it curves upward, the posterior pyriform cortex along a line reaching upward to the medial extension of the rhinal fissure (*fs.rh.m.*) thus completing the great horizontal rhinencephalic cortical ring, as de-

scribed in the earlier part of this paper dealing with the external form of the brain (page 185). In *Caenolestes* this line is longer than in the Virginia opossum, because the position of the rhinal fissure is considerably higher. The exact point at which the rhinal fissure hits the subicular edge of the hippocampus is not determinable, owing to the fading out of histological differentiation, and therefore the extended portion of the fissure is shown as a broken line (*fs.rh.m.*, Figs. 6, 12, 17, 42).

Elliot Smith (1895b, p. 183), in discussing the hippocampus of *Notoryctes*, makes the following remarks: "The hippocampus is equally convoluted in all mammals, because it reaches its maximum development quite early in the phylogenetic history of the individual. Thus in *Platypus* it possesses a histological differentiation quite as complex and fine as is found in the highest mammals. Like the pyriform, it is developed early both in phylogeny and ontogeny in accordance with the development of the olfactory apparatus. Because part of the smell center should reach a high state of development, when the pallium is not proportionately intricate, is no argument that the cortex of the smell center does not behave like the rest of the cortex in similar circumstances. It should be noted, however, that, intimate as is the connection between the hippocampus and the olfactory lobe, the relative sizes of the two parts are by no means constant. Thus, in spite of the marked difference in the sizes of the olfactory bulb in *Ornithorhynchus* and *Perameles*, there is no appreciable difference in the sizes of their hippocampi. In *Notoryctes* the size of the hippocampus is relatively small, considering its huge olfactory. *What determines the size of the hippocampus is hard to say.*" (Italics mine.)

These statements are very suggestive, and in my study of the cerebral hemisphere of *Caenolestes* they have often recurred to me. Are we, however, warranted in saying that the hippocampus reaches its "maximum development" in the lower mammals, and that in *Ornithorhynchus* the histological differentiation is "quite as complex and fine as is found in the highest mammals"? It is true, as explained above, that the definitive mammalian "hippocampal figure" appears suddenly and at a constant level in the lowest mammals. But the amount of inrolling and the development of the temporal pole of the hippocampus both vary considerably, as we have seen, in the marsupials alone. Now in the marsupials we find also a great variation in the amount of neopallium present, without, however, a greatly elongated eutherian type of corpus callosum to account for the temporal pole by mere displacement of the entire supra- and postcommissural hippocampus. Fur-

thermore, in considering the histological differentiation of the hippocampus, we now know that if we take into account not merely the simple ranks of ammon's pyramids and the granules of the gyrus dentatus as they are stained by ordinary cell methods, such as Nissl, but also the many types of cells in the other layers of the hippocampal cortex, as they were demonstrated by Cajal (1911) in the mouse by the silver methods, we get a tremendously suggestive picture of the possibilities for progressive complication within the hippocampus. I attempted to count the different types of intrahippocampal reenforcing and stepping-up devices as described and figured by Cajal in the mouse, and found some twenty-five or thirty of them. Only a comparative study by the silver methods (such as that being carried on by del Rio-Hortega, 1919) of the whole range of mammalian brains could reveal the stage at which the maximum development (maximum histological differentiation) of the hippocampus was attained. I should rather expect to find it nearer the upper than the lower end of the mammalian phylum, among those brains in which the discrepancy between the olfactory bulbs and the neopallium is marked, but in which smell is still an active function. As Elliot Smith remarks, the hippocampus does not, like the pyriform lobe and the tuberculum olfactorium, decrease *pari passu* with the olfactory bulbs. On the contrary, it holds its own (disregarding the amosmatic brains, and even they manage to retain some recognizable hippocampus), perhaps because it *can* so efficiently combine enhancement of function with economy of space (see page 217) as the neopallium increases, and this, apparently because of the increasing activity back and forth between the two. In this connection it is to be noted (see Herrick, 1922, page 196) that the primordium hippocampi has as early as the preganoidean stage no direct somatic connections, and in the ganoidean stage begins to lose direct olfactory connections, and very soon begins to develop a characteristic structure in the presence of indirect correlation and associational connections from various sources, "resulting in a topographical rearrangement which prepared the way for the differentiation within this area of true hippocampal cortex in higher forms" (Herrick, 1922, page 196). The ability of the hippocampus to preserve a sort of structural constancy or individuality with almost total loss of direct olfactory connections, and in the face of great reduction of primary and secondary olfactory centers, has apparently undergone no abatement in the presence of the developing mammalian neopallium. Its peculiar structural pattern has merely unfolded in its own way; it has not approximated the neopallial structural pattern.

Functionally the increasing neopallial-hippocampal bond is of tremendous import. On reflex and lower psychic levels olfaction is mainly if not exclusively linked with food and sex; on higher psychic levels it becomes increasingly significant esthetically and intellectually, in correspondence with the increase of the neural connections between the hippocampus and the somatic pallium which apparently accompanies the *decrease* of neural connections between the hippocampus and the diminishing olfactory bulbs and their immediately dependent secondary olfactory centers. This does not mean that the hippocampus increases phylogenetically in the same ratio with the somatic pallium, or in any constant ratio with it. Apparently neither is true. But it seems hard to believe that a direct relation does not exist between the neopallium and the hippocampus, a relation which is responsible not only for the remarkable preservation of pattern (not its *initiation*) and maintenance of size of the latter structure in higher and relatively microsmatic brains, but finally for the sublimation of smell into a "nobler" sense, practically as truly subservient to higher psychic life as are vision and audition. Language is a form of behavior, and the rich poetic and spiritual imagery clustering around such words as "fragrance" and "incense" amply testifies at once to the importance of the existing sense of smell in man and to its sublimation. So too does the tremendous evocatory power of smell, when it recreates vanished experiences which arouse high and beautiful thoughts or emotions.

CORPUS STRIATUM

The corpus striatum consists of a caudate nucleus (*nuc.caud.*, Figs. 12, 15, 28-38) projecting into the ventricle and extending farther rostrally and caudally than any other part of the striatum, and a lentiform nucleus lateral and ventral to it, in two parts, the putamen (*put.*, Figs. 29-35) and globus pallidus (*gl.p.*, Figs. 12, 15; *glob.p.*, Fig. 34). The medial or septal part of the head of the caudate nucleus, the nucleus accumbens (*nuc.ac.*, Figs. 12, 20-30; the lateral parolfactory nucleus of Johnston, 1913—see page 199), has already been mentioned in connection with the fusion of the head of the caudate with the tuberculum (page 196). Rostral to this fusion the large head of the caudate is laced with small diagonal fiber bundles, like darning stitches, and heavily fringed on its ventral border with diagonal fibers, all the way from the septum to the lateral arm of the anterior commissure (Fig. 29). It is split dorsally by a small caudally directed ventral diverticulum of the lateral ventricle, roofed by the lower surface of the anterior commissure which here breaks across the ventricle to reach

the median wall (Figs. 30-31). The large and compact internal capsule (*cap.i.*, Figs. 29-34) separates the caudate nucleus from the large putamen, whose lateral boundary is the curve of the external capsule (*cap.e.*, Figs. 28-35). The caudate nucleus, including the nucleus accumbens, and the putamen are in this series characterized by a peculiar dark, gun-metal gray background, seen also in the lateral and basal amygdaloid nuclei, setting all these structures sharply off from neighboring ones. I do not know what it may be worth as a criterion, but it is interesting to note that the nucleus accumbens, which Johnston (1913) considers to be the lateral parolfactory nucleus, is uniform with the rest of the caudate in this respect, being sharply delimited from the septum by this ground color.*

The globus pallidus (*gl.p.*, Figs. 12, 15; *glob.p.*, Fig. 34) is a prominent mass of pale giant cells in a rich tangle of fibers, situated near the center of the sections in which it appears, just below the internal capsule as it passes into cerebral peduncle, and not very far behind the anterior commissure level. Small giant cells trailing irregularly towards the basal forebrain bundle in the innominate region (prethalamus, border nucleus of Völsch) may belong to the basal nucleus of the palacostriatum (De Vries, 1910; Kappers, 1921b).

NEOPALLIUM

No attempt has been made to analyze the neopallium of *Caenolestes* into anatomical regions, although even this inadequate series gives some evidence of its possibility. It is hoped that the two series to be made from the brains of *Orolestes* may, with the aid of the intensive cortical analysis of the Virginia opossum in this laboratory (Gray, 1924), render a fairly adequate analysis of the somatic cortex possible in *Caenolestes* and *Orolestes*.

The *claustrum*, which Brodmann and others consider to be a neopallial derivative, is certainly present in the more dorsal part of the

*I was much interested to read (Kappers, 1923, page 365) of the "Spatz reaction" of sulphur ammonium, based on the presence of iron, which gives a much stronger blue color in the palaeostriatum than in the neostriatum. While not inclined to attribute very great test force to his reaction, Kappers thought it interesting that results on the chick and man corresponded. In this iron-haematoxylin series of *Caenolestes* the dark coloration has a very different distribution. The globus pallidus is conspicuously without it. The caudate nucleus (nearly all but the tail) and parts of the tuberculum fusing with it, the putamen, and the two newest amygdaloid nuclei (according to Johnston, 1923), the lateral and the basal, are the regions strikingly affected by the dark coloration. Very pronounced in the rostral portion of all the structures named, it tends to fade out caudally in each of them. I do not know enough about either the Spatz reaction or the dark coloration here to form an opinion as to any relation between them.

anterior pyriform region, but is not clearly delimited. It apparently forms an irregular cell plate between the external capsule and a cell-poor strip in which some fibers appear which may be an incipient capsule extrema (*clau.*, *cap.ex.*, Figs. 27-34). Elliot Smith (1919a, b) considers the claustrum a derivative of the upturned lower edge of the pyriform cortex. This material affords no real evidence either way, although the claustrum here seems to be continuous with the deeper layers of the neopallial cortex.

The cells of this brain, which are remarkably well stained, considering the method used, may be divided first of all into two groups, which Völsch (1906) has called round and pyramid cells, referring to the hemisphere only. The pyramidal cells (using the term very loosely for any angular cell) are black and show dendritic stumps, sometimes very long. The round cells are pale-stained with the nucleus clearly visible. The variation in size is very much greater for the round cells than for the pyramidal cells. All the giant cells of the hemisphere belong to the former type, those of the basal olfactory region, in the anterior portion of the lateral limb of the median fore-brain bundle (see page 198), those of the regio innominata, prethalamus, etc. The smallest cells also belong to this type, the olfactory granules, those of the gyrus dentatus, the exceedingly fine granules of some of the islands of Calleja in the tuberculum, and the scattered masses of very tiny granules tucked between adjacent structures like packing. Völsch considers these last to be non-nervous, I do not know on what evidence other than their size and resemblance to stained glia nuclei. The largest cells of the pyramidal or dark angular type are probably those found in the basal nucleus of the amygdala, for the hemisphere at least.

GENERAL CONSIDERATIONS

In the beginning of this study of a type of brain totally unknown to me I saw that no rational progress was possible save upon a broad and strictly comparative basis. In the course of parallel studies of the literature and of the sections of this particular brain, some more general considerations arose, first as persistent questions, usually entirely unanswerable by me at least, but sometimes coming halfway to rest in my mind. Two of these are briefly outlined below.

One of the most puzzling aspects of such neurological studies as this comes out in connection with the very great diversity of anatomical pattern of structures whose afferent and efferent connections seem almost or practically identical. How shall we explain the enormously

discrepant anatomical structure? The real functional adequacy of every existing structure must of course be taken for granted as the basis of its existence. But more than this must apparently be called in to explain its particular anatomical type of structure. Phyletic tradition clarifies some things—inherited type of anatomical pattern (or the potency to develop it) retained, embellished, even accentuated, so long as it is functionally utilisable or perhaps not actually disadvantageous. The hippocampus is perhaps the classic example under this head. Even when no longer adequate, as evidenced by reduction and degeneration, it seems very hard to be got rid of entirely. Fortunately for the comparative neurologist, it hangs on, sometimes only a pale and shrunken relic, long after it has been more or less supplanted by other structures, with similar or unlike functions, which have become more important in the action-system of the animal either because of fundamentally more serviceable type of anatomical structural pattern, or because of more useful afferent or efferent connections, or for both reasons. Thus a sort of natural selection (Roux's "struggle of parts"?) is constantly operating among structural patterns. A structural pattern is safe so long as the demands upon it are not too heavy, or so long as a competing pattern with greater possibilities does not outrun it. The superiority and final supremacy of the competing pattern may depend not only upon its intrinsic possibilities, but also upon its topographical position and the character and activity of neighboring structures. Thus the early "physiological isolation" (to use Child's now familiar term, 1915, 1921) of the primordium hippocampi from direct sensory stimuli (see page 219) led to "topographical and physiological relationships [which have] prepared the way for the differentiation within this area of true hippocampal cortex in higher forms" (Herrick, 1922, page 196). The trail of the Law of Neurobiotaxis is to be seen everywhere, and this is apparently the most potent single factor which can be invoked immediately to "explain" structural pattern (i.e., to translate it into functional pattern) within the central nervous system. Acting in concert with phyletic tradition, it might well result in exaggerations of structural pattern which do not seem to parallel the development of functional pattern. We must, of course, in comparing similar lists of afferent and efferent connections of very dissimilar structures, take into account the quantitative and positional differences of all the factors, as well as the selective operation of synapses. On the surface, perhaps the discrepancies seem much more striking on the structural than on the functional side, although sometimes the reverse seems to be true. We must of course believe that no two

structures that are visibly different do have absolutely identical functions, not even right and left members of paired structures in the same brain. On the whole, however, the physiologist's frequently voiced objection to the anatomist's too great emphasis upon structural pattern may not perhaps be an unjustifiable one. The modern anatomist, of course, regards structure as merely a useful and convenient clue to function.

Another question which naturally arises in such a study as this concerns what, if any, light is thrown upon the method of progressive evolution in the central nervous system. Davidson Black (1913, page 366) quotes Roux's definition of the two phases of the development of an active tissue: "self-differentiation", which goes on without regard to functional differentiation, and "dependent differentiation", which cannot proceed normally in the absence of functional connections—in the nervous system, functional continuity of neuron systems, as Bechterew pointed out. The transition point between these two phases of growth Black calls the "critical point". Apparently the essential character or activity of progressive evolution in the central nervous system is the pushing forward of this "critical point", relegating more and more of the period of "dependent differentiation" back into the period of "self-differentiation", where it undergoes compression, abridgement, and, so to speak, distortion, thus building the foundation for newer and higher development in the period of dependent differentiation—like using the capstones of a newer structure to strengthen and enlarge the old foundation and so to fit it for still newer and more ambitious structures. What this "antedating" of structure implies in physiological terms, what metabolic changes or reorganization by which, or accompanying which, dissociations take place, some phases dropping out, others being temporally dislocated with reference to the stimuli formerly necessary to elicit them, or whether the dissociations are more apparent than real, I am not competent to discuss.

In attempting to reconstruct the phyletic history of any form, it is to be remembered that not all the foundation blocks, original or "second-hand", have been retained in the enlarged foundation—some have been rejected entirely, all have been recut, and many have altered their relations with reference to others. The higher the form the greater the compression, abridgment and rearrangement, and therefore the greater the necessity of prudence and hesitation in the reconstruction of the *detail* of phyletic pattern, no matter how simple and obvious, perhaps deceivingly so, the ontogenetic pattern may appear.

SUMMARY

1. The brain of *Caenolestes* (and of *Orolestes*) is of the extreme macromastic type, characterized by the great size of the olfactory bulbs and the great development of the higher rhinencephalic centers (Herrick, 1921).

2. The neopallium, in contrast, forms only the shallow cap of the cerebral hemisphere above the high rhinal and hippocampal fissures, and may perhaps be relatively the least extensive or the second least extensive among mammals (Herrick, 1921).

3. The olfactory cortices—pyriform and hippocampal—form, with the aid of the anterior olfactory nucleus, two periventricular rhinencephalic rings partially united anteriorly: a smaller *quasi*-vertical one rostrally, in which the pyriform and hippocampal cortices are doubly (supra- and infraventricularly) but indirectly united through the anterior olfactory nucleus; and a much larger horizontal ring formed by the additional direct union of the posterior pyriform cortex and the subicular margin of the ammon's horn on the median surface of the hemisphere between the medial prolongations of the rhinal and amygdaloid fissures. The two olfactory cortices are split apart dorsally by the wedgelike neopallium and ventrally by a similar wedgelike formation composed of the tuberculum olfactorium rostrally and the amygdaloid complex caudally.

4. The marked antero-posterior foreshortening of this brain is one of its characteristic features, frequently in evidence. It is probably to be explained, in part at least, by the exaggeration of the subcortical centers.

5. The hippocampus begins immediately behind the olfactory (bulbar) formation, passes through practically all reptilian stages precommissurally, and at the dorsal commissure rather suddenly assumes definitive mammalian form, under the three-fold influence of external pressure, internal pressure and neurobiotactic attraction. Its temporal end has just begun to recurve, the gyrus dentatus alone being involved. In the Virginia opossum the ammon's horn has begun to follow suit. More advanced critical stages can be added to these to form a complete and illuminating series of the phyletic development of the mammalian hippocampus, omitting the monotremes and the lowly marsupial *Notoryctes*, aberrantly advanced in this respect.

6. In the amount of gyrus dentatus exposed upon the median surface of the hemisphere (an index of the degree of inrolling of the hippocampus) *Caenolestes* is intermediate between *Notoryctes* and *Perameles*.

7. The enormous size of the amygdaloid complex is a striking characteristic of this brain, as is also its prolonged contact with the temporal hippocampus, both ammon's horn and gyrus dentatus.

8. The absence of the aberrant bundle of the anterior commissure would on the basis of Elliot Smith's definition of it as a diagnostic character of diprotodont brains, seem to put *Caenolestes* among the polyprotodonts, and even to raise the question of the exclusion of the diprotodonts from America.

9. The cerebellum of *Caenolestes* and *Orolestes* is almost exactly intermediate between those of *Notoryctes* and *Perameles*, the two simplest mammalian cerebella hitherto described.

10. In consideration of the marked specialization and aberrancy of the two monotreme brains, the brain of *Caenolestes* (and of *Orolestes* so far as can be judged from the external anatomy) ranks as one of the three simplest and most generalised mammalian brains known at the present time. In view of the exaggerated development of the temporal pole of the hippocampus in *Notoryctes*, it really takes first rank. But on the whole, it is fairly intermediate between *Notoryctes* and *Perameles*, the two simplest and most generalized mammalian brains hitherto described (Elliot Smith), which it most closely resembles. It should offer—and actually does offer—especially promising clues for the reconstruction of the presumptive phyletic stages involved in the transition from the reptilian to the mammalian type. And since it is at once a mammalian brain and so simple and generalized—almost, indeed, a mammalian brain reduced to lowest terms—it also offers hints of unusual legibility for the verification of the structural activities of the nervous system, viewed as regulatory behavior of a more or less plastic material. The cerebral hemisphere of *Caenolestes*, the only part of the brain yet studied in any detail, fairly swarms with exquisitely clear examples of structural evidences of the operation of neurobiotaxis, the principle whose conception and definition by Dr. Kappers of Amsterdam has transferred the study of brain morphology from a static to a dynamic basis.

ACKNOWLEDGMENTS

I am glad to acknowledge here my indebtedness to Dr. C. Judson Herrick, who set me upon this research and who has aided me with practical advice; to Dr. Wilfred H. Osgood, who procured for me the use of further material and in other ways furthered this work; to Dr. Percival Allen Gray, Jr., who did me the very great service of reading the first draft of this paper *without reference to the figures*, and offered

many suggestions which have greatly contributed to the lucidity of the final draft; to Miss Helen Kates for welcome assistance in the tedious work of labelling the figures. I wish it were possible also to acknowledge fully the pleasure and benefit I have received from all those who were interested enough in this material to examine or discuss it with me.

BIBLIOGRAPHY

- BECCARI, N.
1910. Il lobo parolfattorio nei mammiferi. *Arch. anat. e embriol.*, **9**, pp. 173-220.
- BLACK, DAVIDSON.
1913. The study of an atypical cerebral cortex. *Jour. Comp. Neurol.*, **23**, pp. 351-369.
- CHILD, C. M.
1915. Individuality in organisms. The University of Chicago Press.
1921. The origin and development of the nervous system. The University of Chicago Press.
1924. Physiological foundations of behavior. New York.
- CROSBY, ELIZABETH C.
1917. The forebrain of Alligator mississippiensis. *Jour. Comp. Neurol.*, **27**, pp. 325-402.
- DART, R. A.
1920. A contribution to the morphology of the corpus striatum. *Jour. Anat.*, **55**, pp. 1-26.
- GRAY, P. A., JR.
1924. The cortical lamination pattern of the opossum, *Didelphys virginiana*. *Jour. Comp. Neurol.*, **37**, pp. 221-263.
- HERRICK, C. JUDSON.
1910. The morphology of the forebrain in Amphibia and Reptilia. *Jour. Comp. Neurol.*, **20**, pp. 413-549.
1921a. The brain of *Caenolestes obscurus*. Field Museum of Nat. Hist., Pub. 207, Zool. Ser., **14**, pp. 157-162. 3 figures.
1921b. The connections of the vomeronasal nerve, accessory olfactory bulb and amygdala in amphibia. *Jour. Comp. Neurol.*, **33**, pp. 213-280.
1922. Functional factors in the morphology of the forebrain of fishes. *Libro en honor de S. Ramon y Cajal*, Madrid, pp. 143-244.
1924a. The nucleus olfactorius anterior of the opossum. *Jour. Comp. Neurol.*, **37**, pp. 317-359.
1924b. Neurological foundations of animal behavior. New York.
- VAN HÖEVELL, J. J. L. D.
1916. The phylogenetic development of the cerebellar nuclei. *K. Akad. van Wetenschappen*, Amsterdam, **18**, pp. 1421-1434.
- JOHNSTON, J. B.
1913. The morphology of the septum, hippocampus, and pallial commissures in reptiles and mammals. *Jour. Comp. Neurol.*, **23**, pp. 371-478.
1915. Cell masses in the forebrain of the turtle, *Cistudo carolina*. *Jour. Comp. Neurol.*, **25**, pp. 393-468.
1923. Further contributions to the evolution of the forebrain. *Jour. Comp. Neurol.*, **35**, pp. 337-481.
- KAPPERS, C. U. ARIENS
1917. Further contributions to neurobiotaxis. IX. An attempt to compare the phenomena of neurobiotaxis with the phenomena of taxis and tropism. The dynamic polarization of the neurone. *Jour. Comp. Neurol.*, **27**, pp. 261-298.

- 1921a. On structural laws in the nervous system. The principles of neurobio-taxis. *Brain*, **44**, pp. 125-129.
- 1921b. Die vergleichende Anatomie des Nervensystems der Wirbeltiere und des Menschen. II. Abschnitt: Vergleichende Anatomie des Kleinhirns, des Mittel- und des Zwischenhirns und des Vorderhirns. Haarlem.
1923. Le développement ontogénétique du corps strié des oiseaux en comparaison avec celui des mammifères et de l'homme. *Schweizer Arch. F. Neurol. u. Psychiat.*, **13**, pp. 348-370.
- LEVI, G.
- 1904a. Ueber die Entwicklung und Histogenese der Ammonshornformation. *Schultze's Arch.*, **64**, pp. 389-404.
- 1904b. Sull'origine filogenetica della formazione ammonica. *Arch. ital. di anat. e embriol.*, **3**, pp. 234-247.
- LIVINI, G.
1908. Il proencefalo di un marsupiale. *Arch. ital. di anat. e embriol.*, **6**, pp. 549-584.
- McCOTTER, R. E.
1912. The connections of the vomeronasal nerve and the accessory olfactory bulb in the opossum and other mammals. *Anat. Rec.*, **6**, pp. 299-318.
- OBECHAIN, JEANNETTE B.
- 1923a. The brain of *Caenolestes obscurus*. *Anat. Rec.*, **25**, p. 145.
- 1923b. The brain of *Caenolestes obscurus*. *Trans. Illinois State Academy of Science*, **16**, pp. 100-106.
- OSGOOD, W. H.
1921. A monographic study of the American marsupial, *Caenolestes*. *Field Mus. of Nat. Hist., Pub. 207, Zool. Ser.*, **14**, pp. 1-156.
- RAMON Y CAJAL, P.
- 1917-19. Nuevo estudio del encefalo de los reptiles. *Trabajos del Lab. de Investigaciones biol. de la Univ. de Madrid*, **15**, pp. 83-99; **16**, pp. 309-333.
1922. El cerebro de los batracios. *Libro en honor de S. Ramon y Cajal, Madrid*, pp. 13-58.
- RAMON Y CAJAL, S.
1911. *Histologie du système nerveux de l'homme et des vertébrés, tome II.*, Paris.
- RIO-HORTEGA, P. DEL
1919. Particularidades histológicas de la fascia dentata de algunos mamíferos. *Trabajos del Lab. de Investigaciones de la Univ. de Madrid*, **16**, pp. 291-308.
- RÖTHIG, P.
1910. Riechbahnen, Septum, und Thalamus bei *Didelphys marsupialis*. *Abh. Senckenb. Nat. Ges. Frankfurt a. M.*, **31**, pp. 1-19.
- SMITH, G. ELLIOT
1894. A preliminary communication upon the cerebral commissures of mammalia, with especial reference to the Monotremata and Marsupialia. *Proc. Linn. Soc. New. So. Wales*, **9**, pp. 635-656.
- 1895a. The cerebrum of the marsupial mole, *Notoryctes typhlops*. *Zool. Anz.*, **18**, pp. 480-482.
- 1895b. The comparative anatomy of the cerebrum of *Notoryctes typhlops*. *Trans. Roy. Soc., So. Australia*, **19**, pp. 167-193.
- 1896a. The brain of foetal *Ornithorhynchus*. *Quart. Jour. Micr. Soc.*, **39**, pp. 191-206.
- 1896b. Structure of the cerebral hemisphere of *Ornithorhynchus*. *Jour. Anat. and Physiol.*, **30**, pp. 463-487.
- 1896c. The fascia dentata. *Anat. Anz.*, **12**, pp. 119-126.
- 1896d. The morphology of the limbic lobe, corpus striatum, the reptilian septum pellucidum, and the fornix. *Jour. Anat. and Physiol.*, **30**, pp. 157-167, and pp. 185-203.
- 1897a. The morphology of the indusium and the striae Lancisii. *Anat. Anz.*, **13**, pp. 23-27.


- 1897b. The origin of the corpus callosum; a comparative study of the hippocampal region of the cerebrum of Marsupialia and certain Cheiroptera. *Trans. Linn. Soc. Lond.*, **7**, pp. 47-69.
- 1897c. Further observations upon the fornix, with especial reference to the brain of *Nyctophilus*. *Jour. Anat. and Physiol.*, **32**, pp. 231-246.
- 1897d. The fornix superior. *Jour. Anat. and Physiol.*, **31**, pp. 80-92.
- 1897e. The relation of the fornix to the margin of the cerebral cortex. *Jour. Anat. and Physiol.*, **32**, pp. 23-58.
1899. The brain in the Edentata. *Trans. Linn. Soc. Lond.*, 2nd ser., Zool., **7**, pp. 277-394.
- 1902a. Descriptive and illustrated catalogue of the physiological series of comparative anatomy contained in the Museum of the Royal College of Surgeons of England. Vol. 2. London. 2nd ed.
- 1902b. The primary subdivision of the mammalian cerebellum. *Jour. Anat. and Physiol.*, **36**, pp. 381-385.
- 1902c. Notes on the brain of *Macroscelides* and other Insectivora. *Jour. Linn. Soc., Zool. Ser.*, **28**, pp. 443-448.
- 1902d. On a peculiarity of the cerebral commissures in certain Marsupialia not hitherto recognised as a distinctive feature of the Diprotodontia. *Proc. Roy. Soc.*, **70**, pp. 226-231.
- 1903a. On the so-called "gyrus hippocampi". *Jour. Anat. and Physiol.*, **37**, pp. 324-328.
- 1903b. On the morphology of the cerebral hemisphere in the vertebrata, with especial reference to an aberrant commissure found in the brains of certain reptiles. *Trans. Linn. Soc. Lond.*, 2nd Zool. Ser., **8**, pp. 455-500.
- 1903c. On the morphology of the brain of the mammalia, with especial reference to that of the Lemurs, recent and extinct. *Trans. Linn. Soc. Lond.*, **8**, part 10, pp. 319-432.
- 1903d. Notes on the morphology of the cerebellum. *Jour. Anat. and Physiol.*, **37**, pp. 329-332.
- 1903e. Further observations on the natural mode of subdivision of the mammalian cerebellum. *Anat. Anz.*, **23**, pp. 368-384.
- 1919a. A preliminary note on the morphology of the corpus striatum and the origin of the neopallium. *Jour. Anat.*, **53**, pp. 272-291.
- 1919b. A note on Professor Landau's memoir on "The comparative anatomy of the nucleus amygdalae, the claustrum and the insular cortex." *Jour. Anat.*, **53**, pp. 361-362.
- VÖLSCH, M.
1906. Zur vergleichenden Anatomie des Mandelkern und seiner Nachbargebilde. *Anat. Anz.*, **68**, pp. 573-683; **76**, pp. 373-523.
- DE VRIES, E.
1910. Das Corpus Striatum der Säugetiere. *Anat. Anz.*, **37**, pp. 385-405.
- WINKLER, C. and POTTER, ADA
1911. An anatomical guide to experimental researches on the rabbit's brain. Amsterdam.
1914. An anatomical guide to experimental researches on the cat's brain. Amsterdam.

ABBREVIATIONS

al.	alveus	cx. pir. m.	cortex piriformis medial
a. ln.	area lunata cerebelli	cx. pir. p.	cortex piriformis posterior
amg.	nucleus amygdalae; amygdaloid complex	d. b.	diagonal band of Broca
a. precom.	area precommissuralis	d. v.	diverticulum ventralis (of superior recess)
a. pt.	area pterioidea cerebelli	fila ol.	fila olfactoria
aq.	aqueduct of Sylvius	fim.	fimbria
b.	nucleus basalis amygdalae	floc.	flocculus
b. ac.	nucleus basalis accessorius amygdalae	f. med. t.	fasciculus medialis telencephali (medial forebrain bundle)
b. ol.	bulbus olfactorius	f. med. t. l.	fasciculus medialis telencephali (lateral limb of m. f. b.)
b. ol. ac.	bulbus olfactorius accessorius	form. bul.	formatio bulbaris (olfactorius)
c. (Fig. 8)	brachium conjunctivum	f. precom.	fornix praecommissuralis
c.	nucleus centralis amygdalae	fs. amg.	fissura amygdaloidea
c. a.	commissura anterior sive ventralis; see v.	fs. amg. m.	fissura amygdaloidea medialis
c. am.	cornu ammonis	fs. circ.	fissura circularis
cap. e.	capsula externa	fs. ch.	fissura choroidea
cap. em.	capsula extrema	fs. di-tel.	fissura di-telencephalica
cap. i.	capsula interna	fs. erh.	fissura endorhinalis
ch., ch. op.	chiasma opticum	fs. f. d.	fissura fimbrio-dentata
ci. am.	cingulum ammonis	fs. fim. al.	fissura fimbrio-alvearis
ci. lim.	cingulum limitans	fs. fim. d.	fissura fimbrio-dentata
clau.	claustrum	fs. h., fs. hip.	fissura hippocampi
c. c.	corpus callosum	fs. pln.	fissura postlunata
c. d.	commissura dorsalis; see d., ps. d., ps.v.	fs. pnod.	fissura postnodularis
c. g. l.	corpus geniculatum laterale	fs. orb.	fissura orbitalis
c. g. m.	corpus geniculatum mediale	fs. pr.	fissura prima cerebelli
col. a.	colliculus anterior	fs. precul.	fissura preculminata
col. p.	colliculus posterior	fs. rh.	fissura rhinalis
cort.	nucleus corticalis amygdalae	fs. rh. a.	fissura rhinalis anterior
cul.	culmen, pars culminata cerebelli	fs. rh. arc.	fissura rhinalis arcuata
dec. pyr.	decussatio pyramidalis	fs. rh. m.	fissura rhinalis medialis
deep nuc.	deep nuclei of cerebellum (undivided)	fs. s.	fissura secunda cerebelli
em. nat.	eminentia natiformis	fs. spyr.	fissura suprapyramidalis
cx. am.	cortex ammonis	fs. tr. cb.	fissura transversa cerebelli
cx. am. iven.	cortex ammonis intraventricularis	g. bas. op.	ganglion basale opticum
cx. am. xven.	cortex ammonis extraventricularis	g. 5	ganglion semilunaris
cx. dent.	cortex dentatus	gl. p., glob. p.	globus pallidus
cx. hip. a.	cortex hippocampalis anterior (precommissuralis)	gy. dent.	gyrus dentatus
cx. pir.	cortex piriformis		
cx. pir. a.	cortex piriformis anterior		


hab.	habenula	nuc. ol. ant. l. v.	nucleus olfactorius anterior, pars latero-ventralis
hip. a.	hippocampus anterior (prae commissuralis)	nuc. ol. ant. m.	nucleus olfactorius anterior, pars medialis
i. C.	island of Calleja	nuc. ol. ant. p.	nucleus olfactorius anterior, pars posterior
inf.	infundibulum	nuc. pol. l.	nucleus parolfactorius lateralis
interpos. l.	interpositio lateralis	nuc. pol. m.	nucleus parolfactorius medialis
interpos. m.	interpositio medialis	nuc. tr. ol. l.	nucleus tracti olfactorii lateralis; see n. t. o. l.
int. plate	intercalary plate of Johnston (1923)	ped. floc.	pedunculus flocculi
ip. m.	interpositio medialis	pfloc.	paraflocculus
l.	nucleus lateralis amygdalae	pl. ch. 3 (4)	plexus choroidea ventriculi tertii (quarti)
lg.	lingula cerebelli	p. preul.	pars preculminata cerebelli
lob. a. cb.	lobus anterior cerebelli	p. spyr.	pars suprapyramidalis cerebelli
lob. m. cb.	lobus medialis cerebelli	put.	putamen
lob. p. cb.	lobus posterior cerebelli	pyr.	pyramis cerebelli
l. perf. a.	locus perforatus anterior	r.	corpus restiforme
m.	nucleus medialis amygdalae	rec. i.	recessus inferior
med.	medulla oblongata	rec. s.	recessus superior
m. i.	massa intermedia	rad. thal.	radiatio thalamica
neop.	neopallium	st. glom.	stratum glomeruli
n. 2	nervus opticus	st. gr.	stratum granulare
n. 5	nervus trigeminus	st. m. c.	stratum cellularum mitralium
nod.	nodulus cerebelli	st. med.	stria medullaris, see st. t. 5
n. t. o. l.	nucleus tracti olfactorii lateralis	st. t.	stria terminalis
nuc. ac.	nucleus accumbens	st. t. 1	stria terminalis, bundle 1 (Johnston, 1923) = com. bundle
nuc. amg. b.	nucleus basalis amygdalae	st. t. 2	stria terminalis, bundle 2 (Johnston, 1923) = ol. proj. tr. (amg.)
nuc. amg. b. ac.	nucleus basalis accessorius amygdalae	st. t. 3	stria terminalis, bundle 3 (Johnston, 1923) = subcom. bundle
nuc. amg. c.	nucleus centralis amygdalae	st. t. 4	stria terminalis, bundle 4 (Johnston, 1923) = supracom. bundle
nuc. amg. cort.	nucleus corticalis amygdalae	st. t. 5	stria terminalis, bundle 5 (Johnston, 1923) = st. med. bundle
nuc. amg. l.	nucleus lateralis amygdalae	st. t. bed	stria terminalis bed
nuc. amg. m.	nucleus medialis amygdalae	tect.	tectum mesencephali
nuc. caud.	nucleus caudatus		
nuc. d. b.	nucleus of diagonal band of Broca		
nuc. ip.	nucleus interpeduncularis		
nuc. ol. ant.	nucleus olfactorius anterior (Herrick)		
nuc. ol. ant. d.	nucleus olfactorius anterior, pars dorsalis		
nuc. ol. ant. ex.	nucleus olfactorius anterior, pars externus		
nuc. ol. ant. l.	nucleus olfactorius anterior, pars lateralis		

t. ol.	tuberculum olfacto- rium	tr. t. a.	tractus temporo-am- monis of Cajal (1911; sphenotemp., 1906); see ci. lim. (cingulum limitans) and ci. am. (cingu- lum ammonis)
tr. ol.	tractus olfactorius		
tr. ol. d. m.	tractus olfactorius dorso-medialis		
tr. ol. fr.	tractus olfacto-fronta- lis	t. v. 4 uv.	taenia ventriculi quarti uvula cerebelli
tr. ol. i.	tractus olfactorius in- termedius (commis- suralis)	v. cb. v. l. v. m. a.	ventriculus cerebelli ventriculus lateralis velum medullare ante- rius
tr. ol. l.	tractus olfactorius la- teralis (massive part)	v. ol. v. 3 (4)	ventriculus olfactorius ventriculus tertius (quartus)
tr. ol. l. d.	tractus olfactorius la- teralis, pars dorsa- lis	xven. alv. xven. cx. am.	extraventricular alveus extraventricular cor- tex ammonis
tr. ol. l. p. d.	tractus olfactorius la- teralis, pedunculus dorsalis	*	line of fracture sup- posed to mark boundary between extra- and intraven- tricular precommis- sural ammon's cor- tex.
tr. ol. l. p. v.	tractus olfactorius la- teralis, pedunculus ventralis	—>	arrows to show direc- tion and approxi- mate degree of force operating in the formation of the definitive mamma- lian "hippocampal figure", figure 11
tr. ol. l. v.	tractus olfactorius la- teralis, pars ventra- lis		
tr. ol. m.	tractus olfactorius me- dialis		
tr. ol. s. + c.	tractus olfacto-septa- lis et corticalis	1	line of amputation of cortex dentatus, fig- ure 17a
tr. op.	tractus opticus	2	line of amputation of cortex ammonis, figure 17b
tr. st. thal.	tractus strio-thalami- cus		




1

OROLESTES INCA.
DORSAL VIEW OF BRAIN.
Five times natural size.




2

OROLESTES INCA.
VENTRAL VIEW OF BRAIN.
Five times natural size.



3




4

OROLESTES INCA.


FIG. 3. LATERAL VIEW OF BRAIN.

FIG. 4. MEDIAN SECTION OF BRAIN.

Five times natural size.



5






6

MEDIAN SURFACE OF HEMISPHERE.

FIG. 5. OROLESTES INCA.

FIG. 6. CAENOLESTES OBSCURUS. RECONSTRUCTION.

Five times natural size.




FIGS. 7 TO 10. CAENOLESTES OBSCURUS. CEREBELLUM. RECONSTRUCTIONS. X 8.


FIGS. 10a-10b. CEREBELLA OF NOTORYCTES AND PERAMELES.

FIG. 11. DEVELOPMENT OF MAMMALIAN "HIPPOCAMPAL FIGURE."

12. MEDIAL



13. LATERAL




A.A.T. →




16

a

14. DORSAL



15. VENTRAL



FIGS. 12 TO 15. CAENOLESTES OBSCURUS. RECONSTRUCTIONS OF HEMISPHERES.

FIG. 16. VENTRAL DIVERTICULUM OF RECESSUS SUPERIOR. a. HORIZONTAL-CAENOLESTES OBSCURUS X 30 b. SAGITTAL-DIDELPHIS VIRGINIANA.







FIG. 17. CAENOLESTES OBSCURUS. DISSECTED DIAGRAM OF HIPPOCAMPUS. X 12.


FIG. *8. DIDELPHIS VIRGINIANA. TEMPORAL HIPPOCAMPUS. X 5.




19



20



21



22


TEMPORAL RECURVING OF HIPPOCAMPUS.

FIG. 19. HYPISPRYMNUS (LIVINI) X 3.


FIG. 20. RABBIT (WINKLER-POTTER).

FIG. 21. FELIS LEO (ELLIOT-SMITH) X 1/2.


FIG. 22. NOTORYCTES (DART).




23 (SEC 243)




24 (SEC 271)




25 (SEC 307)



26 (SEC 343)




27 (SEC 382)




28 (SEC 410)

CAENOLESTES OBSCURUS.


x 8.




29 (SEC. 432)



30 (SEC. 510)




31 (SEC. 542)




32 (SEC. 581)

CAENOLESTES OBSCURUS.

X 8.




33. (SEC 599)



34. (SEC 642)




35. (SEC 668)




36. (SEC 704)

CAENOLESTES OBSCURUS.


X 8.




37 (SEC 730)



38 (SEC 760)



39 (SEC 804)



40 (SEC 832)

CAENOLESTES OBSCURUS.


X 8.




41 (SEC 831)



42 (SEC 879)



43 (SEC. 900)



44 (SEC. 920)

CAENOLESTES OBSCURUS.

X 8.

INDEX

Names of new family, genus and species are printed in **bold-faced** type.

Abductor caudae externus.....	38	Epitrochlearis	40
Abductor digiti indicis.....	47	Ethmoid	101
Abductor digiti minimi.....	47	Extensor brevis digitorum.....	54
Abductor pollicis.....	47	Extensor carpi radialis brevis.....	44
Abductor pollicis longus.....	44	Extensor carpi radialis longus.....	44
Abductors.....	46-47	Extensor carpi ulnaris.....	45
Aberrant bundle.....	200	Extensor caudae lateralis.....	39
Accessory olfactory bulb.....	189	Extensor digitorum communis.....	45
Accessory olfactory nucleus.....	194	Extensor digitorum lateralis.....	45
Acromio-deltoidaeus.....	40	Extensor hallucis longus.....	55
Acromio-trapezius.....	24	Extensor indicis.....	46
Adductor brevis.....	48	Extensor longus digitorum.....	55
Adductor longus.....	47	External Characters.....	19
Adductor magnus.....	47		
Adductors.....	46-57	Fasciculus aberrans.....	200
Alimentary Canal.....	69	Femoro-coccygeus.....	48
Amygdaloid complex.....	185-201	Fimbrio-alvear fissure.....	184
Anconeus.....	42	Fimbrio-dentate fissure.....	184
Anterior olfactory nucleus.....	191	Fissura amygdaloidea.....	181
Arm.....	89	Fissura rhinalis arcuata.....	179
Auditory ossicles.....	107	Flexor accessorius.....	52
Auriculo-occipitalis.....	23	Flexor carpi radialis.....	42
		Flexor carpi ulnaris.....	42
Biceps brachii.....	41	Flexor digitorum fibularis.....	51
Biceps femoris.....	49	Flexor digitorum profundus.....	43
Biventer cervicis.....	25	Flexor digitorum sublimis.....	43
Brachialis anticus.....	41	Flexor digitorum tibialis.....	52
Brachioradialis (Supinator longus).....	44	Flexors.....	46
Brachase.....	99	Foot.....	95
Brain stem.....	160	Forearm.....	89
Buccinator.....	31	Frontals.....	103
Canines.....	118	Gastrocnemius.....	50
Caudal vertebrae.....	84	Gemelli.....	37
Caudofemoralis.....	48	Genioglossus.....	33
Cerebellum.....	159-186	Genio-hyoideus.....	32
Cerebral hemispheres.....	158	Glands.....	73
Cervical vertebrae.....	77	Gluteus maximus.....	36
Cervico-auricularis.....	23	Gluteus medius.....	36
Choriod fissure.....	185	Gluteus minimus.....	37
Circulatory System.....	77	Gracilis.....	50
Claustrum.....	221		
Cleido-mastoideus.....	33	Habits.....	17
Cleido-occipitalis.....	34	Hand.....	92
Complexus.....	25	Hippocampal commissure.....	210
Coracobrachialis brevis.....	41	Hippocampal fissure.....	184
Cremaster.....	35	Hippocampal formation.....	206
		Hyoglossus.....	33
Dentition.....	111	Hyoid.....	106
Digastricus.....	32		
Diprotodonty.....	128	Iliacus.....	37
Distribution.....	15	Iliococcygeus.....	39
Dorso-cuticularis.....	23		

- Iliocostalis.....29
 Incisors.....116
 Infracoccygeus.....39
 Infraspinatus.....40
 Interossei.....46
 Intestines.....72
 Ischiococcygeus.....39

 Lacrymals.....103
 Larynx.....77
 Latissimus dorsi.....24
 Leg.....93
 Levator scapulae.....24
 Liver.....73
 Longissimus.....28
 Longissimus capitis.....25
 Longus colli.....27
 Lumbar vertebrae.....84
 Lumbricales.....56
 Lungs.....76

 Mammae.....75
 Mandible.....108
 Mandibulo-auricularis.....31
 Masseter.....30
 Maxillae.....102
 Measurements.....21
 Molars.....102
 Mouth.....69
 Muscles of the abdomen.....35
 Muscles of the anterior limb.....40
 Muscles of the back.....24
 Muscles of the forefoot.....46
 Muscles of the head.....30
 Muscles of the hind foot.....56
 Muscles of the hind limb.....47
 Muscles of the neck.....32
 Muscles of the sacrum.....36
 Muscles of the skin.....22
 Muscles of the tail.....38
 Muscles of the thorax.....34
 Mylo-hyoideus.....33
 Myrmecoboides.....139

 Nasals.....101
 Neopallium.....180

 Obliquus capitis inferior.....26
 Obliquus capitis superior.....26
 Obliquus externus abdominis.....35
 Obliquus internus abdominis.....35
 Obturator externus.....38
 Obturator internus.....37
 Occipitals.....104
 Oesophagus.....71
 Olfactory areas.....189
 Olfactory bulbs.....158-178
 Omo-cleido-transversarius.....34
 Omo-hyoideus.....33
 Orolestes.....168-178

 Palatine.....105
 Palmaris longus.....42
 Pancreas.....74
 Parietals.....103
 Parolfactory nuclei.....199
 Pectineus.....48
 Pectoral girdle.....87
 Pectoralis major.....34
 Pectoralis minor.....34
 Pelvic girdle.....88
 Periotic.....106
 Peroneus brevis.....52
 Peroneus longus.....53
 Peroneus tertius.....53
 Phylogeny.....140
 Plantaris.....51
 Popliteus.....55
 Premaxillae.....102
 Premolars.....119
 Pronator quadratus.....44
 Pronator teres.....42
 Psoas magnus.....30
 Psoas parvus.....30
 Pterygoideus externus.....31
 Pterygoideus internus.....31
 Pubococcygeus.....39
 Pyriformis.....37
 Pyriform lobe.....180-200

 Quadratus femoris.....37
 Quadratus lumborum.....29

 Rectus abdominis.....35
 Rectus capitis anticus major.....27
 Rectus capitis anticus minor.....27
 Rectus capitis lateralis.....27
 Rectus capitis posterior major.....26
 Rectus capitis posterior minor.....26
 Rectus capitis posterior superficialis.....26
 Rectus femoris.....49
 Respiratory System.....76
 Retractor naris.....32
 Rhinal fissure.....180
 Rhomboideus.....24
Rhyncholestes.....169
Rhyncholestes raphanurus.....170
 Ribs.....86

 Sacral vertebrae.....84
 Sacrococcygeus.....39
 Sacrospinalis.....29
 Salivary glands.....74
 Sartorius.....50
 Scalenus.....35
 Semimembranosus.....50
 Semispinalis cervicis.....28
 Semitendinosus.....49
 Septal region.....199
 Serratus magnus.....34
 Serratus posterior inferior.....25

Serratus posterior superior.....	25
Skeleton	77
Skull	99
Sphenoid bones.....	105
Spinalis dorsi.....	27
Spino-deltoides.....	40
Spino-trapezius	24
Spleen	74
Splenius	25
Squamosal.....	103
Sterno-hyoideus.....	32
Sterno-mastoideus.....	33
Sterno-thyroideus	32
Sternum	86
Stomach	71
Stria terminalis.....	204
Styloglossus	33
Subclavius.....	34
Subcrureus	49
Subscapularis	40
Supinator (brevis).....	45
Supracostalis.....	34
Supraspinatus.....	40
Taxonomy.....	140
Temporalis	30
Teres major.....	41
Thigh	93

Thoracic vertebrae.....	82
Thymus	75
Thyrohyoideus.....	33
Thyroid	75
Tibialis anticus.....	33
Tibialis posticus.....	54
Tongue	70
Trachea	77
Transversus abdominis.....	35
Triceps brachii.....	41
Tuberculum olfactorium.....	185, 195
Turbinals	101
Tympanic.....	106

Urinogenital System.....	61
--------------------------	----

Vacuities	107
Vastus externus.....	49
Vastus internus	49
Vomer	105

Wynyardia	136
Wynyardiidae	138

Zygomata	103
Zygomaticus	32

QL

Fieldiana: zoology

1

F4

v.14

Biological
& Medical
Serials

PLEASE DO NOT REMOVE
CARDS OR SLIPS FROM THIS POCKET

UNIVERSITY OF TORONTO LIBRARY

STORAGE

